AQA

Please write clearly in block capitals.

Centre number | | | | | | Candidate number | | | | |

Surname StemUp Model Solution

Forename(s) _____

Candidate signature _____

A-level
FURTHER MATHEMATICS

Paper 3 Mechanics

Thursday 13 June 2019 Afternoon Time allowed: 2 hours

Materials
- You must have the AQA formulae and statistical tables booklet for A-level Mathematics and A-level Further Mathematics.
- You should have a scientific calculator that meets the requirements of the specification. (You may use a graphical calculator.)
- You must ensure you have the other optional Question Paper/Answer Book for which you are entered (**either** Discrete **or** Statistics). You will have 2 hours to complete **both** papers.

Instructions
- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer each question in the space provided for that question. If you require extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do **not** write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work you do not want to be marked.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
TOTAL	

Information
- The marks for questions are shown in brackets.
- The maximum mark for this paper is 50.

Advice
- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

Do not write outside the box

Answer **all** questions in the spaces provided.

1 A spring has natural length 0.4 metres and modulus of elasticity 55 N

Calculate the elastic potential energy stored in the spring when the extension of the spring is 0.08 metres.

Circle your answer.

[1 mark]

0.176 J (0.44 J) 0.88 J 1.76 J

2 A particle has an angular speed of 72 revolutions per minute.

Find the angular speed in radians per second.

Circle your answer.

[1 mark]

$\frac{6\pi}{5}$ $\left(\frac{12\pi}{5}\right)$ 12π 24π

Do not write outside the box

3 A disc, of mass m and radius r, rotates about an axis through its centre, perpendicular to the plane face of the disc.

The angular speed of the disc is ω.

A possible model for the kinetic energy E of the disc is

$$E = km^a r^b \omega^c$$

where a, b and c are constants and k is a dimensionless constant.

Find the values of a, b and c.

[3 marks]

$[E] = [\frac{1}{2}mv^2] = ML^2T^{-2}$

$[m] = M, \quad [r] = L, \quad [\omega] = T^{-1}$

$ML^2T^{-2} = M^a L^b (T^{-1})^c = M^a L^b T^{-c}$

Comparing powers: M: $a = 1$

L: $b = 2$

T: $-c = -2 \Rightarrow c = 2$

$\therefore a = 1, \; b = 2, \; c = 2$


Turn over for the next question

Do not write outside the box

4 **In this question use $g = 10\,\text{m}\,\text{s}^{-2}$**

An inelastic string has length 1.2 metres.
One end of the string is attached to a fixed point O.
A sphere, of mass 500 grams, is attached to the other end of the string.

The sphere is held, with the string taut and at an angle of 20° to the vertical, touching the chin of a student, as shown in the diagram below.



The sphere is released from rest.

Assume that the student stays perfectly still once the sphere has been released.

4 (a) Calculate the maximum speed of the sphere.

[3 marks]

Height dropped to lowest point: $h = 1.2 - 1.2\cos 20° = 1.2(1 - \cos 20°)$

Max speed at lowest point. KE gained = GPE lost:

$\frac{1}{2} \times 0.5 \times v^2 = 0.5 \times 10 \times 1.2(1 - \cos 20°)$

$v^2 = 1.447...$

$v = \sqrt{1.447...} = 1.203...$

Maximum speed = 1 m s^{-1} (1 s.f., since g = 10)

Do not write outside the box

4 (b) Find the maximum tension in the string.

[3 marks]

Max tension at the lowest point (string vertical, speed greatest).

Resolving vertically, towards O: $T - 0.5 \times 10 = 0.5 \times v^2/1.2$

$T = 5 + 0.5 \times 1.447.../1.2$

$T = 5.603...$

Maximum tension = 6 N (1 s.f.)

4 (c) State, with a reason, whether or not the sphere touches the student's chin again after it has been released.

[2 marks]

No, it will not touch her chin again.


Air resistance acts on the sphere, so some energy is lost

and it cannot rise back up to its original position.

Turn over for the next question

Do not write outside the box

5 The triangular region shown below is rotated through 360° around the x-axis, to form a solid cone.



The coordinates of the vertices of the triangle are (0, 0), (8, 0) and (0, 4).

All units are in centimetres.

5 (a) State an assumption that you should make about the cone in order to find the position of its centre of mass.

[1 mark]

The cone is uniform (it has constant density).

5 (b) Using integration, prove that the centre of mass of the cone is 2 cm from its plane face.

[5 marks]

Line through (0, 4) and (8, 0): $y = 4 - \frac{1}{2}x$

Let density = ρ. Mass = $\rho \times \frac{1}{3}\pi \times 4^2 \times 8 = 128\pi\rho/3$

Moments about the y-axis: $(128\pi\rho/3)\,\bar{x} = \int_0^8 \rho\pi y^2 x\,dx$

$(128\pi\rho/3)\,\bar{x} = \pi\rho \int_0^8 x(4 - \frac{1}{2}x)^2\,dx$

$= \pi\rho \int_0^8 (16x - 4x^2 + \frac{1}{4}x^3)\,dx$

$= \pi\rho \left[8x^2 - (4/3)x^3 + (1/16)x^4\right]_0^8$

$= \pi\rho\,(512 - 2048/3 + 256) - 0$

$= 256\pi\rho/3$

$(128/3)\,\bar{x} = 256/3$

$\bar{x} = 2$

The plane face of the cone lies on the y-axis ($x = 0$),

so the centre of mass is 2 cm from its plane face, as required.

Do not write outside the box

5 (c) The cone is placed with its plane face on a rough board. One end of the board is lifted so that the angle between the board and the horizontal is gradually increased. Eventually the cone topples without sliding.

5 (c) (i) Find the angle between the board and the horizontal when the cone topples, giving your answer to the nearest degree.

[2 marks]

Topples when G is vertically above the lowest edge of the base.

G is 2 cm above the base on the axis; base radius = 4 cm

$\tan \alpha = 4/2 = 2$

$\alpha = 63.43...°$

$\alpha = 63°$ (nearest degree)

5 (c) (ii) Find the range of possible values for the coefficient of friction between the cone and the board.

[3 marks]

The cone does not slide before $\alpha = 63.4...°$, so up to this angle

friction is not limiting: $F < \mu R$

$R = mg \cos \alpha$, $F = mg \sin \alpha$

$mg \sin \alpha < \mu\, mg \cos \alpha$

$\mu > \tan \alpha = 2$

$\therefore \ \mu > 2$

Do not write outside the box

6 A ball moving on a smooth horizontal surface collides with a fixed vertical wall. Before the collision, the ball moves with speed $7\,\text{m}\,\text{s}^{-1}$ and at an angle of 40° to the wall.

After the collision, the ball moves with speed $5\,\text{m}\,\text{s}^{-1}$ and at an angle of 26° to the wall.

Model the ball as a particle.

6 (a) Find the coefficient of restitution between the ball and the wall, giving your answer correct to two significant figures.

[3 marks]

Velocity components perpendicular to the wall:

before: $7\sin 40^\circ$ after: $5\sin 26^\circ$

Newton's law of restitution: $e \times 7\sin 40^\circ = 5\sin 26^\circ$

$e = 5\sin 26^\circ / (7\sin 40^\circ)$

$e = 0.4871\ldots$

$e = 0.49$ (2 s.f.)

Do not write outside the box

6 (b) Determine whether or not the wall is smooth.

Fully justify your answer.

[3 marks]

Velocity components parallel to the wall:

before: $7\cos 40° = 5.362... = 5.36\ m\ s^{-1}$

after: $5\cos 26° = 4.494... = 4.49\ m\ s^{-1}$

$5.36 \neq 4.49$, so the parallel component has been reduced.

A smooth wall exerts no force parallel to it, so this component would be unchanged if the wall were smooth.

As the parallel components are not equal, the wall is not smooth.

Turn over for the next question

Do not write outside the box

7 A particle of mass 2.5 kilograms is attached to one end of a light, inextensible string of length 75 cm. The other end of this string is attached to a point A.

The particle is also attached to one end of an elastic string of natural length 30 cm and modulus of elasticity λ N. The other end of this string is attached to a point B, which is 60 cm vertically below A.

The particle is set in motion so that it describes a horizontal circle with centre B. The angular speed of the particle is $8\,\text{rad}\,\text{s}^{-1}$

Find λ, giving your answer in terms of g.

[9 marks]

Let θ = angle between string AP and the horizontal radius BP.

AP = 0.75 m, AB = 0.6 m vertical, BP horizontal

$\sin\theta = 0.6/0.75 = 4/5$, $\cos\theta = 3/5$

Radius $r = BP = \sqrt{(0.75^2 - 0.6^2)} = 0.45$ m

T_1 = tension in AP (inextensible), T_2 = tension in BP (elastic)

Resolving vertically (no vertical acceleration):

$T_1 \sin\theta = 2.5g$

$(4/5)T_1 = 2.5g \Rightarrow T_1 = 25g/8$

Resolving horizontally towards B, acceleration $r\omega^2$:

$T_2 + T_1\cos\theta = 2.5 \times 0.45 \times 8^2$

$T_2 + (25g/8)(3/5) = 72$

$T_2 = 72 - 15g/8$

Extension of elastic string = 0.45 − 0.30 = 0.15 m

Hooke's law: $T_2 = \lambda \times 0.15/0.3 = \lambda/2$

$\lambda/2 = 72 - 15g/8$

$\lambda = 144 - 15g/4$

Do not write outside the box

Turn over for the next question


Do not write outside the box

8 **In this question use $g = 9.8\,\text{m s}^{-2}$**

A 'reverse' bungee jump consists of two identical elastic ropes. One end of each elastic rope is attached to either side of the top of a gorge.

The other ends are both attached to Hannah, who has mass 84 kg

Hannah is modelled as a particle, as shown in the diagram below.



The depth of the gorge is 50 metres and the width of the gorge is 40 metres.

Each elastic rope has natural length 30 metres and modulus of elasticity 3150 N

Hannah is released from rest at the centre of the bottom of the gorge.

8 (a) Show that the speed of Hannah when the ropes become slack is $30\,\text{m s}^{-1}$ correct to two significant figures.

[6 marks]

Initially each rope runs from the top edge down to Hannah:

length = $\sqrt{(50^2 + 20^2)} = \sqrt{2900} = 53.85...$ m

extension = $53.85... - 30 = 23.85...$ m

Initial EPE (two ropes) = $2 \times 3150 \times 23.85...^2 / (2 \times 30)$

$= 59735$ J

Ropes become slack when each rope is 30 m long:

depth below top = $\sqrt{(30^2 - 20^2)} = \sqrt{500} = 22.36...$ m

height risen = $50 - 22.36... = 27.639...$ m

GPE gained = $84 \times 9.8 \times 27.639... = 22753$ J

Conservation of energy: EPE lost = GPE gained + KE gained

$59735 - 22753 = \frac{1}{2} \times 84 \times v^2$

$36982 = 42v^2$

Do not write outside the box

$v^2 = 880.5...$

$v = 29.67...$

$v = 30 \text{ m s}^{-1}$ (2 s.f.) as required

8 (b) Determine whether Hannah is moving up or down when the ropes become taut again.

[5 marks]

Once the ropes are slack, only gravity acts on Hannah.

The ropes become taut again when she is 22.36... m above the top:

$h = 50 + \sqrt{(30^2 - 20^2)} = 72.36...$ m above the bottom of the gorge.

If the ropes stayed slack, all the initial EPE becomes GPE at the highest point H: $59735 = 84 \times 9.8 \times H$

$H = 72.56...$ m

72.36... m < 72.56... m, so the ropes become taut just below Hannah's highest point, before she has stopped rising.

Hence Hannah is moving upwards when the ropes become taut again.

END OF QUESTIONS

Do not write outside the box

There are no questions printed on this page

DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third-party copyright material are published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2019 AQA and its licensors. All rights reserved.