

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

Forename(s)

Candidate signature

A-level MATHEMATICS

Paper 1

Wednesday 6 June 2018

Morning

Time allowed: 2 hours

Materials

- You must have the AQA Formulae for A-level Mathematics booklet.
- You should have a graphical or scientific calculator that meets the requirements of the specification.

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer each question in the space provided for that question. If you require extra space, use an AQA supplementary answer book; do **not** use the space provided for a different question.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 100.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
TOTAL	



Answer **all** questions in the spaces provided.

1 $y = \frac{1}{x^2}$

Find an expression for $\frac{dy}{dx}$

Circle your answer.

[1 mark]

$$\frac{dy}{dx} = \frac{0}{2x}$$

$$\frac{dy}{dx} = x^{-2}$$

$$\frac{dy}{dx} = -\frac{2}{x}$$

$$\frac{dy}{dx} = -\frac{2}{x^3}$$

2 The graph of $y = 5^x$ is transformed by a stretch in the y -direction, scale factor 5

State the equation of the transformed graph.

Circle your answer.

[1 mark]

$$y = 5 \times 5^x$$

$$y = 5^{\frac{x}{5}}$$

$$y = \frac{1}{5} \times 5^x$$

$$y = 5^{5x}$$



3 A periodic sequence is defined by $U_n = \sin\left(\frac{n\pi}{2}\right)$

State the period of this sequence.

Circle your answer.

[1 mark]

8

2π

4

π

4 The function f is defined by $f(x) = e^{x-4}$, $x \in \mathbb{R}$

Find $f^{-1}(x)$ and state its domain.

[3 marks]

Turn over for the next question

Turn over ►



5 A curve is defined by the parametric equations

$$x = 4 \times 2^{-t} + 3$$

$$y = 3 \times 2^t - 5$$

5 (a) Show that $\frac{dy}{dx} = -\frac{3}{4} \times 2^{2t}$

[3 marks]

5 (b) Find the Cartesian equation of the curve in the form $xy + ax + by = c$, where a , b and c are integers.

[3 marks]



- 6 (a)** Find the first three terms, in ascending powers of x , of the binomial expansion of $\frac{1}{\sqrt{4+x}}$

[3 marks]

- 6 (b)** Hence, find the first three terms of the binomial expansion of $\frac{1}{\sqrt{4-x^3}}$

[2 marks]

Question 6 continues on the next page

Turn over ►



- 6 (c)** Using your answer to part **(b)**, find an approximation for $\int_0^1 \frac{1}{\sqrt{4-x^3}} dx$, giving your answer to seven decimal places.

[3 marks]

- 6 (d) (i)** Edward, a student, decides to use this method to find a more accurate value for the integral by increasing the number of terms of the binomial expansion used.

Explain clearly whether Edward's approximation will be an overestimate, an underestimate, or if it is impossible to tell.

[2 marks]



6 (d) (ii) Edward goes on to use the expansion from part **(b)** to find an approximation

for $\int_{-2}^0 \frac{1}{\sqrt{4-x^3}} dx$

Explain why Edward's approximation is invalid.

[2 marks]

Turn over for the next question

Turn over ►



7 Three points A , B and C have coordinates $A(8, 17)$, $B(15, 10)$ and $C(-2, -7)$

7 (a) Show that angle ABC is a right angle.

[3 marks]

7 (b) A , B and C lie on a circle.

7 (b) (i) Explain why AC is a diameter of the circle.

[1 mark]



7 (b) (ii) Determine whether the point $D(-8, -2)$ lies inside the circle, on the circle or outside the circle.

Fully justify your answer.


[4 marks]

Turn over for the next question

Turn over ►



Angle AOB is θ radians.



[4 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

- 8 (b)** Use the Newton-Raphson method with $\theta_1 = \pi$, to find θ_3 as an approximation for θ .
Give your answer correct to five decimal places.

[3 marks]

- 8 (c)** Given that $\theta = 1.89549$ to five decimal places, find an estimate for the percentage error in the approximation found in part **(b)**.

[1 mark]

Turn over for the next question

Turn over ►



The sum of the first 36 terms of the sequence is equal to the square of the sum of the first 6 terms.

[4 marks]

[illegible]

- 9 (b)** Given that the sixth term of the sequence is 25, find the smallest possible value of a .
[5 marks]

Turn over for the next question

Turn over ►



[illegible]

- 10 (b)** The scientist wants the mass of caffeine in her body to stay below 480 mg
- Use the model to find the earliest time that she could drink another cup of strong coffee.

Give your answer to the nearest minute.

[3 marks]

- 10 (c)** State a reason why the mass of caffeine remaining in the scientist's body predicted by the model may not be accurate.

[1 mark]

Turn over for the next question

Turn over ►



- 11** The daily world production of oil can be modelled using

$$V = 10 + 100\left(\frac{t}{30}\right)^3 - 50\left(\frac{t}{30}\right)^4$$

where V is volume of oil in millions of barrels, and t is time in years since 1 January 1980.

- 11 (a) (i)** The model is used to predict the time, T , when oil production will fall to zero.

Show that T satisfies the equation

$$T = \sqrt[3]{60T^2 + \frac{162\,000}{T}}$$

[3 marks]

- 11 (a) (ii)** Use the iterative formula $T_{n+1} = \sqrt[3]{60T_n^2 + \frac{162\,000}{T_n}}$, with $T_0 = 38$, to find the values of T_1 , T_2 , and T_3 , giving your answers to three decimal places.

[2 marks]



11 (a) (iii) Explain the relevance of using $T_0 = 38$

[1 mark]

11 (b) From 1 January 1980 the daily use of oil by one technologically developing country can be modelled as

$$V = 4.5 \times 1.063^t$$

Use the models to show that the country's use of oil and the world production of oil will be equal during the year 2029.

[4 marks]

Turn over for the next question

Turn over ►



12 $p(x) = 30x^3 - 7x^2 - 7x + 2$

12 (a) Prove that $(2x + 1)$ is a factor of $p(x)$

[2 marks]

12 (b) Factorise $p(x)$ completely.

[3 marks]



$$\frac{30 \sec^2 x + 2 \cos x}{7} = \sec x + 1$$

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.


Turn over ►



13

A company is designing a logo. The logo is a circle of radius 4 inches with an inscribed rectangle. The rectangle must be as large as possible.

The company models the logo on an x - y plane as shown in the diagram.



Use calculus to find the maximum area of the rectangle.

Fully justify your answer.

[10 marks]



Do not write
outside the
box

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

Turn over for the next question


Turn over ►



14

Some students are trying to prove an identity for $\sin(A + B)$.

They start by drawing two right-angled triangles ODE and OEF , as shown.



The students' incomplete proof continues,

Let angle $DOE = A$ and angle $EOF = B$.

In triangle OFR ,

Line 1 $\sin(A + B) = \frac{RF}{OF}$

Line 2 $= \frac{RP + PF}{OF}$

Line 3 $= \frac{DE}{OF} + \frac{PF}{OF}$ since $DE = RP$

Line 4 $= \frac{DE}{\dots} \times \frac{\dots}{OF} + \frac{PF}{EF} \times \frac{EF}{OF}$

Line 5 $= \dots + \cos A \sin B$

14 (a)

Explain why $\frac{PF}{EF} \times \frac{EF}{OF}$ in Line 4 leads to $\cos A \sin B$ in Line 5

[2 marks]



14 (b) Complete Line 4 and Line 5 to prove the identity

$$\text{Line 4} \quad = \frac{DE}{\dots} \times \frac{\dots}{OF} + \frac{PF}{EF} \times \frac{EF}{OF}$$

$$\text{Line 5} \quad = \dots + \cos A \sin B$$

[1 mark]

14 (c) Explain why the argument used in part (a) only proves the identity when A and B are acute angles.

[1 mark]

14 (d) Another student claims that by replacing B with $-B$ in the identity for $\sin(A + B)$ it is possible to find an identity for $\sin(A - B)$.

Assuming the identity for $\sin(A + B)$ is correct for all values of A and B , prove a similar result for $\sin(A - B)$.

[3 marks]

Turn over ►



[4 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

[2 marks]

There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

