

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

StemUp Model Solution

Forename(s)

Candidate signature

AS CHEMISTRY

Paper 2: Organic and Physical Chemistry

Friday 9 June 2017

Afternoon

Time allowed: 1 hour 30 minutes

Materials

For this paper you must have:

- the Periodic Table/Data Sheet, provided as an insert (enclosed)
- a ruler with millimetre measurements
- a calculator, which you are expected to use where appropriate.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided.
Do not write outside the box around each page or on blank pages.
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

Advice

- You are advised to spend about 65 minutes on **Section A** and 25 minutes on **Section B**.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
Section B	
TOTAL	




J U N 1 7 7 4 0 4 2 0 2

Section A

Answer **all** questions in this section.

0 1

Figure 1 shows the Maxwell–Boltzmann distribution of molecular energies in a sample of gas at a fixed temperature.



0 1 . 1

Label the horizontal axis in **Figure 1**.

[1 mark]

0 1 . 2

On **Figure 1**, sketch a distribution of molecular energies for this sample of gas at a higher temperature.

[2 marks]

0 1 . 3

This gas decomposes on heating.

Explain why an increase in temperature increases the rate at which this gas decomposes.

[2 marks]

At a higher temperature, more molecules have energy greater than or equal to the activation energy ($E \geq E_a$).

So there are more successful collisions in a given time (a higher frequency of successful collisions), so the rate of decomposition increases.



0 2

An experiment was carried out to determine the relative molecular mass (M_r) of a volatile hydrocarbon **X** that is a liquid at room temperature.

A known mass of **X** was vaporised at a known temperature and pressure and the volume of the gas produced was measured in a gas syringe.

Data from this experiment are shown in **Table 1**.

Table 1

Mass of X	194 mg
Temperature	373 K
Pressure	102 kPa
Volume	72 cm ³

0 2 . 1

Calculate the relative molecular mass of **X**.

Show your working.

Give your answer to the appropriate number of significant figures.

The gas constant, $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$

[5 marks]

$$n = pV \div RT$$

$$p = 102 \times 10^3 \text{ Pa}, \quad V = 72 \times 10^{-6} \text{ m}^3, \quad T = 373 \text{ K}$$

$$n = (102 \times 10^3 \times 72 \times 10^{-6}) \div (8.31 \times 373)$$

$$n = 7.344 \div 3099.63 = 0.002369 \text{ mol}$$

$$\text{mass} = 194 \text{ mg} = 0.194 \text{ g}$$

$$M_r = \text{mass} \div \text{moles} = 0.194 \div 0.002369$$

$$M_r = 81.9 = 82 \text{ (2 s.f.)}$$

Relative molecular mass 82



0 2 . 2

Analysis of a different hydrocarbon **Y** shows that it contains 83.7% by mass of carbon.

Calculate the empirical formula of **Y**.

Use this empirical formula and the relative molecular mass of **Y** ($M_r = 86.0$) to calculate the molecular formula of **Y**.

[4 marks]

$$\%H = 100 - 83.7 = 16.3\%$$

$$C : H = 83.7 \div 12.0 : 16.3 \div 1.0$$

$$= 6.975 : 16.3$$

$$\div 6.975 \rightarrow 1 : 2.337$$

$$\times 3 \rightarrow 3 : 7.01 = 3 : 7$$

$$M_r \text{ of } C_3H_7 = 3(12.0) + 7(1.0) = 43.0$$

$$86.0 \div 43.0 = 2, \text{ so molecular formula} = 2 \times C_3H_7$$

Empirical formula C_3H_7

Molecular formula C_6H_{14}



0 3 . 1

Compounds **A**, **B** and **C** all have the molecular formula C_5H_{10}

A and **B** decolourise bromine water but **C** does not.

B exists as two stereoisomers but **A** does **not** show stereoisomerism.

Use this information to deduce a possible structure for each of compounds **A**, **B** and **C** and explain your deductions.

State the meaning of the term stereoisomers and explain how they arise in compound **B**.

[6 marks]

A and B decolourise bromine water, so they are unsaturated: they contain a $C=C$ double bond (they are alkenes).

C does not decolourise bromine water, so it is saturated and has no $C=C$ bond. As C_5H_{10} is not C_nH_{2n+2} , C must be a ring (cycloalkane): C is cyclopentane.

A is pent-1-ene, $CH_2=CHCH_2CH_2CH_3$. It has no stereoisomers because one carbon of its $C=C$ has two identical groups (two H atoms) attached.

B is pent-2-ene, $CH_3CH=CHCH_2CH_3$.

Stereoisomers are molecules with the same structural formula but a different arrangement of the atoms (bonds) in space.

In B there is restricted rotation about the $C=C$ double bond (it cannot rotate), and each carbon of the $C=C$ has two different groups attached (H and CH_3 on one, H and CH_2CH_3 on the other).

So B exists as two stereoisomers: Z-pent-2-ene, with the CH_3 and CH_2CH_3 groups on the same side of the $C=C$, and E-pent-2-ene, with them on opposite sides.



0	4
---	---

When alkanes are burned in an excess of oxygen they produce carbon dioxide and water.

0	4	.	1
---	---	---	---

Write an equation for the complete combustion of propane in oxygen.

[1 mark]



0	4	.	2
---	---	---	---

An expression can be derived using bond enthalpy data to estimate the enthalpy of combustion ($\Delta_c H$) of an alkane.

For an alkane with n carbon atoms: $\Delta_c H = -(496n + 202) \text{ kJ mol}^{-1}$

The enthalpy of combustion of an alkane was calculated to be $-6650 \text{ kJ mol}^{-1}$ using this expression.

Deduce the molecular formula of this alkane.
Show your working.

[2 marks]

$$-6650 = -(496n + 202)$$

$$496n = 6650 - 202 = 6448$$

$$n = 6448 \div 496 = 13$$

$$\text{alkane} = \text{C}_n\text{H}_{2n+2} = \text{C}_{13}\text{H}_{28}$$

Molecular formula of alkane $\text{C}_{13}\text{H}_{28}$

0	4	.	3
---	---	---	---

Suggest **one** reason, other than the use of mean bond enthalpies, why a value for the enthalpy of combustion of a liquid alkane is different from the value obtained using the expression in Question 4.2

[1 mark]

The expression assumes the alkane is a gas, but this alkane is a liquid, so energy is needed to convert it into a gas.



0 4 . 4

Values of the enthalpy change for combustion of 1 g of some alkanes are shown in **Table 2**.


Table 2

	methane	ethane	propane	butane	pentane
Enthalpy change in kJ for combustion of 1 g	-55.6	-52.0	-50.5	-49.6	-48.7

Plot the enthalpy change for the combustion of 1 g against the number of carbon atoms in the alkanes in **Table 2**.

Draw a best fit line and use this to estimate the enthalpy change for combustion of 1 g of propane.

Write your answer in **Table 2**.

[3 marks]

0 4 . 5

Isooctane (2,2,4-trimethylpentane) is an important component of petrol used in cars.

When isooctane is burned, the enthalpy change is -47.8 kJ g^{-1}

Isooctane is a liquid at room temperature with a density of 0.692 g cm^{-3}

Calculate the heat energy released, in kJ, when 1.00 dm^3 of isooctane burns in excess oxygen.

Give your answer to the appropriate number of significant figures.

[2 marks]

$$1.00 \text{ dm}^3 = 1000 \text{ cm}^3$$

$$\text{mass} = 1000 \times 0.692 = 692 \text{ g}$$

$$\text{heat energy} = 692 \times 47.8 = 33\,077.6 \text{ kJ}$$

$$= 3.31 \times 10^4 \text{ kJ (3 s.f.)}$$


Heat energy released 3.31×10^4 kJ



0 5

Ethanedioic acid ($\text{H}_2\text{C}_2\text{O}_4$) is a diprotic acid. Beekeepers use a solution of this acid as a pesticide.

A student carried out a titration with sodium hydroxide solution to determine the mass of the acid in the solution. The student repeated the titration until concordant titres were obtained.



0 5 . 1

The student found that 25.0 cm^3 of the ethanedioic acid solution reacted completely with 25.30 cm^3 of $0.500 \text{ mol dm}^{-3}$ sodium hydroxide solution.

Calculate the mass, in mg, of the acid in 25.0 cm^3 of this solution.

[4 marks]

$$n(\text{NaOH}) = 25.30 \times 10^{-3} \times 0.500 = 0.01265 \text{ mol}$$

$$n(\text{H}_2\text{C}_2\text{O}_4) = 0.01265 \div 2 = 0.006325 \text{ mol}$$

$$M_r(\text{H}_2\text{C}_2\text{O}_4) = 2(1.0) + 2(12.0) + 4(16.0) = 90.0$$

$$\text{mass} = 0.006325 \times 90.0 = 0.569 \text{ g}$$

$$= 569 \text{ mg}$$

Mass of acid 569 mg

0 5 . 2

The student used a wash bottle containing deionised water when approaching the end-point to rinse the inside of the conical flask.

Explain why this improved the accuracy of the titration.

[1 mark]

It washes any ethanedioic acid or sodium hydroxide on the sides of the flask down into the solution, so all of the reactants are in the reaction mixture and can react.

0 5 . 3

Give the meaning of the term concordant titres.

[1 mark]

Titres that are within 0.10 cm^3 of each other.



0 6

2-Methylpropan-1-ol can be prepared by reacting 1-bromo-2-methylpropane with dilute aqueous sodium hydroxide.


0 6 . 1

Name and outline the mechanism for this reaction.

[3 marks]

Name of mechanism Nucleophilic substitution

Mechanism



0 6 . 2

When 2.0 cm^3 of 1-bromo-2-methylpropane ($M_r = 136.9$) were reacted with an excess of sodium hydroxide, 895 mg of 2-methylpropan-1-ol ($M_r = 74.0$) were obtained.

The density of 1-bromo-2-methylpropane is 1.26 g cm^{-3}

Calculate the percentage yield for this reaction.

[3 marks]

$$\text{mass of 1-bromo-2-methylpropane} = 2.0 \times 1.26 = 2.52 \text{ g}$$

$$\text{moles} = 2.52 \div 136.9 = 0.0184 \text{ mol}$$

$$\text{moles of alcohol expected} = 0.0184 \text{ mol}$$

$$\text{theoretical mass} = 0.0184 \times 74.0 = 1.36 \text{ g}$$

$$\% \text{ yield} = (0.895 \div 1.36) \times 100 = 65.7\%$$

Percentage yield 65.7%



0 6 . 3

When 1-bromo-2-methylpropane reacts with hot, concentrated ethanolic potassium hydroxide rather than dilute aqueous sodium hydroxide, a different product is formed.

Name this organic product and name the mechanism for this reaction.

[2 marks]

Name of organic product methylpropene

Name of mechanism elimination

Turn over for the next question



0 7


Alcohols undergo dehydration in the presence of concentrated phosphoric acid, via a carbocation intermediate, to form alkenes.

0 7 . 1

Complete the mechanism for the conversion of 2-methylcyclohexanol into 1-methylcyclohexene via carbocation **D** by drawing

- the structure of the missing intermediate
- all necessary curly arrows.


[4 marks]



0 7 . 2

Draw the structure of a different cyclic alkene formed from carbocation **D**.


[1 mark]



0 7 . 3

Carbocation **D** can undergo a type of reaction called a rearrangement to form carbocation **E**. In this reaction, a hydrogen atom and its bonding pair of electrons move from carbon **a** to carbon **b** as shown in **Figure 2**.

Figure 2



Use your knowledge of carbocations to explain why this rearrangement takes place.

[2 marks]

The rearrangement forms a more stable carbocation.


Carbocation D is secondary but carbocation E is tertiary: a tertiary carbocation has more alkyl groups releasing electrons (inductive effect) towards the positive carbon.

0 7 . 4

As a result of the rearrangement in Question 7.3, a third alkene is formed in this reaction.

Draw the structure of this third alkene.

[1 mark]




Turn over for the next question



0 7 . 5

Cyclohexene is prepared by the dehydration of cyclohexanol using concentrated phosphoric acid as a catalyst. The structure of concentrated phosphoric acid is shown.



Identify the factors that influence the boiling points of each of the compounds in this reaction mixture. State how and explain why cyclohexene can be separated from the reaction mixture.

[6 marks]

Cyclohexene is a non-polar hydrocarbon, so there are only van der Waals' forces between its molecules.

Cyclohexanol has an O-H group, so there are hydrogen bonds between cyclohexanol molecules.

Phosphoric acid has three O-H groups, so there are hydrogen bonds between phosphoric acid molecules.

Hydrogen bonds are much stronger than van der Waals' forces, so cyclohexene has the weakest intermolecular forces and the least energy is needed to overcome them.

Therefore cyclohexene has the lowest boiling point of the three compounds in the mixture.

Cyclohexene is separated by (fractional) distillation: the mixture is heated and cyclohexene boils off first, so it is condensed and collected, while cyclohexanol and phosphoric acid stay in the flask.




0 8

This question is about the structures of some organic molecules.

0 8 . 1

Draw the **skeletal** formula of 3-methylbutanal.


[1 mark]



0 8 . 2

Draw the **displayed** formula of $C_5H_{11}Br$ that is the major product of the reaction of 2-methylbut-2-ene with hydrogen bromide.

[1 mark]




0 8 . 3

Thermal cracking of hydrocarbons produces molecules that are attacked by electrophiles because they have a region of high electron density.

Draw the structure of one of these molecules that contains four carbon atoms.

[1 mark]




Turn over for the next question



0 9

Chloroethene can be polymerised to form poly(chloroethene), commonly known as PVC. This polymer can be used to make pipes, window frames and electrical insulation. Plasticisers can be added to change the properties of PVC

A section of poly(chloroethene) is shown.



0 9 . 1

Chloroethene has a melting point of $-154\text{ }^{\circ}\text{C}$

All types of PVC melt at temperatures over $100\text{ }^{\circ}\text{C}$

Explain why PVC melts at a higher temperature than chloroethene.


[2 marks]

PVC molecules are much bigger and longer than chloroethene molecules, so they have many more electrons and a larger surface area (a greater M_r).

So PVC has stronger van der Waals' forces between its molecules (intermolecular forces), which need more energy to overcome, so PVC melts at a higher temperature.

0 9 . 2

This structure shows a molecule that has been used as a plasticiser in PVC.



Deduce the number of hydrogen atoms in this molecule.

[1 mark]

38



0 9 . 3


Use your understanding of the properties of PVC to explain whether you would expect to find a plasticiser in the PVC used to insulate electrical cables.

[1 mark]

Yes, a plasticiser would be present, because the PVC used to insulate electrical cables needs to be flexible (bendy).


0 9 . 4

A section of the polymer poly(chloroprene), a synthetic rubber, is shown.



Draw the **displayed** formula for the repeating unit of poly(chloroprene).

[1 mark]



Turn over for the next question



Section B

Answer **all** questions in this section.Only **one** answer per question is allowed.

For each answer completely fill in the circle alongside the appropriate answer.

CORRECT METHOD



WRONG METHODS

If you want to change your answer you must cross out your original answer as shown. If you wish to return to an answer previously crossed out, ring the answer you now wish to select as shown. You may do your working in the blank space around each question but this will not be marked.
Do **not** use additional sheets for this working.

1 0

What is the burette reading for this transparent liquid?

[1 mark]

A 24.10 cm³
☐
B 24.30 cm³
☒
C 25.70 cm³
☐
D 25.90 cm³
☐


1 1

A volumetric flask was used to prepare 250 cm³ of a solution.

The solute was added from a plastic weighing container.

	Mass / g
Weighing container with solute	10.13
Weighing container after solute added to volumetric flask	4.48

Each reading from the balance has an uncertainty of ± 0.005 g

What is the percentage uncertainty in the mass of the solute used?

[1 mark]

A 0.09%

☐

B 0.11%

☐

C 0.18%


☒

D 0.22%

☐

1 2

The infrared spectrum of an organic compound is shown.



Which compound produces this spectrum?

[1 mark]

A butanone

☒

B ethanol

☐

C pent-2-ene

☐

D propanoic acid

☐


1	3
---	---

Which is the most likely bond angle around the oxygen atom in ethanol?

[1 mark]**A** 104.5°☒**B** 109.5°☐**C** 120°☐**D** 180°☐

1	4
---	---

Which compound is a structural isomer of Z-but-2-ene?

[1 mark]**A** butane☐**B** E-but-2-ene☐**C** cyclobutane☒**D** methylbut-2-ene☐

1	5
---	---

Which equation is a propagation step in the conversion of trichloromethane into tetrachloromethane by reaction with chlorine in the presence of ultraviolet light?

[1 mark]**A** $\text{CHCl}_3 + \text{Cl}_2 \rightarrow \text{CCl}_4 + \text{HCl}$ ☐**B** $\bullet\text{CCl}_3 + \bullet\text{Cl} \rightarrow \text{CCl}_4$ ☐**C** $\text{CHCl}_3 + \bullet\text{Cl} \rightarrow \text{CCl}_4 + \bullet\text{H}$ ☐**D** $\bullet\text{CCl}_3 + \text{Cl}_2 \rightarrow \text{CCl}_4 + \bullet\text{Cl}$ ☒

1 6

Which compound has the fastest rate of reaction with potassium cyanide to form pentanenitrile?

[1 mark]

A 1-bromobutane

☐

B 1-chlorobutane

☐

C 1-fluorobutane

☐

D 1-iodobutane

☒

1 7

Which alcohol can be oxidised by acidified potassium dichromate(VI) but cannot be dehydrated by heating with concentrated sulfuric acid?

[1 mark]

A 2,3-dimethylbutan-2-ol

☐

B 2,2-dimethylpropan-1-ol

☒

C 2-methylpropan-2-ol

☐

D pentan-3-ol

☐

1 8

How many structural isomers are there with the molecular formula C_3H_6BrCl ?

[1 mark]

A 4

☐

B 5

☒

C 6

☐

D 7

☐

1	9
---	---

Which sample contains the most molecules?

The Avogadro constant, $L = 6.022 \times 10^{23} \text{ mol}^{-1}$

[1 mark]

A 2.10×10^{22} molecules of methane, CH_4

☒

B 1.00 g of oxygen, O_2

☐

C 65.0 mg of hydrogen, H_2

☐

D 0.0300 mol of ethane, C_2H_6

☐

2	0
---	---

Which compound forms a molecular ion with a different precise molecular mass from the other three?

[1 mark]

A butanone

☐

B cyclobutanol

☐

C dimethylpropane


☒

D methylpropanal

☐

2	1
---	---

Hydrogen can be produced by this reaction.



In an experiment 4.20 mol of carbon monoxide were mixed with 2.00 mol of steam. When the reaction reached equilibrium, 1.60 mol of hydrogen had been formed.

What is the value of the equilibrium constant, K_c , for this reaction?

[1 mark]

A 0.30

☐

B 0.41

☐

C 1.54

☐

D 2.46

☒

2	2
---	---

A sample of 2.0 mol dm^{-3} acid has a volume of 100 cm^3

What volume of water, in cm^3 , should be added to this acid to dilute the sample to a concentration of 1.5 mol dm^{-3} ?

[1 mark]

A 25

☐

B 33.3

☒

C 50

☐

D 66.7

☐

Turn over for the next question



2	3
---	---

Two sealed flasks with the same volume are left side by side.

Flask **A** contains 4.0×10^{-3} mol of methane.

Flask **B** contains 340 mg of a different gas.

Both gases are at the same temperature and pressure.

Which gas could be in Flask **B**?

[1 mark]

A CH_2Cl_2

☒

B HBr

☐

C Kr

☐

D PF_3

☐

2	4
---	---

Analysis of a sample of a chemical with formula $\text{C}_{22}\text{H}_{30}\text{N}_6\text{O}_4\text{S}$, showed that it contained 0.0195 mol of carbon.

What mass of nitrogen was present in the sample?

[1 mark]

A 0.041 g

☐

B 0.057 g

☐

C 0.074 g

☒

D 0.420 g

☐

END OF QUESTIONS

Copyright Information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2017 AQA and its licensors. All rights reserved.

