

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

StemUp Model Solution

Forename(s)

Candidate signature

AS CHEMISTRY

Paper 2 Organic and Physical Chemistry

Friday 25 May 2018

Morning

Time allowed: 1 hour 30 minutes

Materials

For this paper you must have:

- the Periodic Table/Data Sheet, provided as an insert (enclosed)
- a ruler with millimetre measurements
- a scientific calculator, which you are expected to use where appropriate.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do **not** write outside the box around each page or on blank pages.
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

Advice

- You are advised to spend about 65 minutes on **Section A** and 25 minutes on **Section B**.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
Section B	
TOTAL	




J U N 1 8 7 4 0 4 2 0 1

Section A

Answer **all** questions in this section.

0 1

Hydrogen peroxide solution decomposes slowly to form water and oxygen.
The reaction is much faster in the presence of a manganese(IV) oxide catalyst.




Three experiments, shown in **Table 1**, were carried out to investigate how the volume of oxygen produced varied over time under different conditions. The same mass of catalyst was used in each experiment.

Table 1

Experiment	Concentration of $\text{H}_2\text{O}_2(\text{aq})$ / mol dm^{-3}	Volume of $\text{H}_2\text{O}_2(\text{aq})$ / cm^3	Temperature / $^{\circ}\text{C}$	Catalyst
1	1.0	50	20	lumps
2	1.0	50	20	powder
3	0.5	50	20	lumps

Figure 1 shows how the volume of oxygen collected varied with time in Experiment 1.

Figure 1



0 1 . 1

Explain, in general terms, how a catalyst increases the rate of a reaction.

[2 marks]

A catalyst provides an alternative reaction route (pathway) with a lower activation energy.

So more of the particles have energy greater than or equal to the lower activation energy, so there are more successful collisions per second.

0 1 . 2

Draw **two** lines on **Figure 1** to show how the volume of oxygen collected varied with time in Experiments **2** and **3**.

Label each line with the experiment number.

[2 marks]

0 1 . 3

Explain, in terms of collision theory, the effect of increasing the concentration of hydrogen peroxide on the rate of reaction.

[2 marks]

Increasing the concentration means there are more H_2O_2 particles in the same volume, so they are closer together.

So successful collisions (with energy at least equal to the activation energy) are more frequent, so the rate of reaction increases.

Turn over ►




0 2

Citric acid, $\text{C}_3\text{H}_5\text{O}(\text{COOH})_3$, occurs naturally in many fruits and can also be synthesised in the laboratory for use as a food flavouring. A student analysed a sample of citric acid to determine its percentage purity.

The student dissolved 784 mg of impure citric acid in water to prepare 250 cm^3 of solution in a volumetric flask.

The student titrated 25.0 cm^3 samples of this solution with $0.0500 \text{ mol dm}^{-3}$ sodium hydroxide solution using phenolphthalein as the indicator.



0 2 . 1

The student rinsed the burette before filling it with the sodium hydroxide solution.

State why the student should use sodium hydroxide solution rather than water for the final rinse of the burette.

[1 mark]

Water would dilute the sodium hydroxide, which would change (lower) its concentration and make the titre larger.

0 2 . 2

The student carried out several titrations. The results are shown in **Table 2**.

Complete **Table 2** to show the titre in each titration.

[1 mark]

Table 2

Titration	Rough	1	2	3
Final reading / cm^3	25.2	23.95	47.65	24.10
Start reading / cm^3	0.0	0.05	23.95	0.10
Titre / cm^3	25.2	23.90	23.70	24.00

0 2 . 3

Calculate the mean titre using the concordant results.
Give your answer to the appropriate number of significant figures.

[2 marks]

Concordant results: titrations 1 and 3 (within 0.10 cm^3)

Mean titre = $(23.90 + 24.00) \div 2 = 23.95 \text{ cm}^3$

Mean titre 23.95 cm^3



0 2 . 4

The total uncertainty when using the burette is $\pm 0.15 \text{ cm}^3$. This is the combination of uncertainties in the start reading, final reading and the determination of the end point.

Use your answer to Question **02.3** to calculate the percentage uncertainty for the use of the burette in this experiment.

[1 mark]

$$\text{Percentage uncertainty} = (0.15 \div 23.95) \times 100 = 0.626\%$$

Percentage uncertainty 0.63 %

0 2 . 5

Use your answer to Question **02.3** to find the mass, in mg, of citric acid dissolved in 250 cm^3 of the solution.

The relative molecular mass (M_r) of citric acid is 192.0

[3 marks]

$$\text{Moles of NaOH} = (23.95 \div 1000) \times 0.0500 = 1.198 \times 10^{-3} \text{ mol}$$

$$\text{Ratio citric acid : NaOH} = 1 : 3$$

$$\text{Moles of citric acid in } 25.0 \text{ cm}^3 = 1.198 \times 10^{-3} \div 3 = 3.992 \times 10^{-4} \text{ mol}$$

$$\text{Moles in } 250 \text{ cm}^3 = 3.992 \times 10^{-4} \times 10 = 3.992 \times 10^{-3} \text{ mol}$$

$$\text{Mass} = 3.992 \times 10^{-3} \times 192.0 = 0.766 \text{ g} = 766 \text{ mg}$$

Mass 766 mg

0 2 . 6

Calculate the percentage purity of this sample of citric acid.

[1 mark]

$$\text{Percentage purity} = (766 \div 784) \times 100 = 97.7\%$$


Percentage purity 97.7 %

Turn over ►



0 3

This question is about enthalpy changes.

0 3 . 1When ethanoic acid reacts with sodium hydroxide, the enthalpy change, ΔH , is $-56.1 \text{ kJ mol}^{-1}$ Calculate the temperature rise when 25 cm^3 of 2.0 mol dm^{-3} aqueous ethanoic acid react with 25 cm^3 of 2.0 mol dm^{-3} aqueous sodium hydroxide.Assume that both solutions have the same initial temperature, have a density of 1.0 g cm^{-3} and a specific heat capacity of $4.18 \text{ J K}^{-1} \text{ g}^{-1}$ **[4 marks]**

$$\text{Moles of CH}_3\text{COOH} = (25 \div 1000) \times 2.0 = 0.050 \text{ mol}$$

$$(\text{NaOH} = 0.050 \text{ mol, so neither is in excess})$$

$$\text{Heat released, } q = 0.050 \times 56.1 = 2.805 \text{ kJ} = 2805 \text{ J}$$

$$\text{Mass of solution} = 25 + 25 = 50 \text{ cm}^3 \times 1.0 \text{ g cm}^{-3} = 50 \text{ g}$$

$$q = mc\Delta T, \text{ so } \Delta T = q \div mc$$

$$\Delta T = 2805 \div (50 \times 4.18) = 13.4 \text{ }^\circ\text{C}$$

Temperature rise 13.4 $^\circ\text{C}$ 

0 3 . 2

A student recorded the temperature of aqueous ethanoic acid in a polystyrene cup for three minutes.

At the fourth minute, the student added sodium hydrogencarbonate.

The student stirred the mixture and carried on recording the temperature every minute for several minutes.


The student's measurements are shown in **Figure 2**.

A best-fit line showing the temperature before mixing has been drawn.

Draw an appropriate best-fit line on **Figure 2** and use it to find the temperature change at the time of mixing.

[2 marks]

Figure 2



Temperature change at time of mixing -5.8 °C


Turn over ►



0 4


The alkanes nonane and 2,4-dimethylheptane are structural isomers with the molecular formula C_9H_{20} . They are found in crude oil and can be separated by fractional distillation. Both can be used in fuels or cracked to form other products.

nonane



boiling point 151 °C

2,4-dimethylheptane



boiling point 134 °C

0 4 . 1

State the general formula of an alkane containing n carbon atoms.
Deduce an expression for the relative molecular mass (M_r) of an alkane in terms of n .
[2 marks]

General formula C_nH_{2n+2}

$$M_r = 12.0n + 1.0(2n + 2)$$

Expression $14.0n + 2.0$

0 4 . 2

Explain why nonane has a higher boiling point than 2,4-dimethylheptane.
[2 marks]

Nonane is unbranched, so its molecules can pack closer together and
have a greater surface area of contact.

So nonane has stronger van der Waals' forces between molecules, which
need more energy to overcome.



0 4 . 3

Give an equation for the complete combustion of nonane.

[1 mark]



0 4 . 4

Nonane is often found in fuel for jet engines. Combustion in jet engines produces pollutants including nitrogen monoxide (NO).

Explain how this nitrogen monoxide is formed.

[2 marks]

Nitrogen and oxygen from the air (drawn into the engine) react together.

This happens at the very high temperature inside the jet engine: $N_2 + O_2 \rightarrow 2NO$

0 4 . 5

Nonane can be cracked to form large quantities of propene.

Name the type of cracking used.

[1 mark]


Thermal cracking

0 4 . 6

The main use of propene, formed from cracking, is to make poly(propene).

Draw the repeating unit of poly(propene).

[1 mark]



Turn over for the next question

Turn over ►



0 5 . 1

A hydrocarbon contains 87.8% by mass of carbon and has a relative molecular mass (M_r) of 82.0

The hydrocarbon decolourises bromine water.

Determine the empirical and molecular formulae of the hydrocarbon.

Suggest **two** possible structures for the hydrocarbon.

Name the type of reaction taking place when bromine water reacts with the hydrocarbon.

[6 marks]

$$C : H = 87.8/12.0 : 12.2/1.0 = 7.32 : 12.2$$


Divide by the smallest: 1 : 1.67, then multiply by 3 to give 3 : 5

So the empirical formula is C_3H_5


$$M_r \text{ of } C_3H_5 = 3(12.0) + 5(1.0) = 41.0 \text{ and } 82.0 \div 41.0 = 2$$

So the molecular formula is C_6H_{10}

It decolourises bromine water, so it contains a $C=C$ double bond. Two possible structures are:



cyclohexene



hexa-1,3-diene

The reaction of the hydrocarbon with bromine water is electrophilic addition: bromine adds across the $C=C$ double bond.




[illegible]

There are no questions printed on this page

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



0 6

Compound **A** is a halogenoalkane.**Compound A**

0 6 . 1

Name Compound **A**.


[1 mark]

3-chloro-2,4-dimethylpentane

0 6 . 2

Compound **A** has a relative molecular mass (M_r) of 134.5The main isotope of hydrogen is ^1H The main isotope of carbon is ^{12}C Chlorine consists of two common isotopes, ^{35}Cl and ^{37}Cl , of which 75% is ^{35}Cl The mass spectrum of **A** was recorded when **A** was ionised by electron impact to form A^+ ions.Draw, on **Figure 3**, the peaks for the main molecular ions in the mass spectrum of **A**.

[2 marks]

Figure 3

Question 6 continues on the next page

Turn over ►



0 6 . 3 Reaction of **A** with warm, dilute aqueous sodium hydroxide forms alcohol **B**.



Name the mechanism for this reaction.

Outline the mechanism using the structure of **A** shown.
Include the structure of the product, alcohol **B**.

[4 marks]

Mechanism Nucleophilic substitution

Outline of mechanism



alcohol B (2,4-dimethylpentan-3-ol)



0 6 . 4 Reaction of **A** with hot, ethanolic potassium hydroxide gives alkene **C**.

Name the mechanism for this reaction.
State the role of the hydroxide ions.


Outline the mechanism using the structure of **A** shown.
Include the structure of the product, alkene **C**.

[6 marks]

Mechanism Elimination

Role of hydroxide ions Base (proton acceptor)

Outline of mechanism



alkene **C** (2,4-dimethylpent-2-ene)


Question 6 continues on the next page

Turn over ►



0 6 . 5 The infrared spectrum in **Figure 4** is that of either alcohol **B** or alkene **C**.

Figure 4



Tick the box that shows the correct compound.

Explain your answer with reference to a bond and the wavenumber of its absorption.

[1 mark]

Alcohol **B**

☐

Alkene **C**



☒

Explanation Alkene C: there is no broad O-H absorption at 3230 to 3550 cm⁻¹, and there is a C=C absorption at about 1650 cm⁻¹ (1620 to 1680 cm⁻¹).



0 6 . 6

Compound **D** reacts with dilute aqueous sodium hydroxide in a similar way to **A** to form alcohol **B**.

Compound **A**Compound **D**

Explain why **D** reacts more quickly than **A** with dilute aqueous sodium hydroxide at the same temperature.

[1 mark]

The C-Br bond is weaker than the C-Cl bond (C-Br has a lower bond enthalpy), so the C-Br bond breaks more easily.

15

Turn over for the next question

Turn over ►



0 7 . 1 Four compounds, all colourless liquids, are

- butan-2-ol
- butanal
- butanone
- 2-methylpropan-2-ol

Two of these compounds can be identified using different test-tube reactions.

Describe these **two** test-tube reactions by giving reagents and observations in each case.

Suggest how the results of a spectroscopic technique could be used to distinguish between the **other** two compounds.

[6 marks]

Test 1: add Tollens' reagent (ammoniacal silver nitrate) to a sample of each of the four compounds and warm in a water bath.

Only butanal (an aldehyde) gives a silver mirror. The other three show no change, so butanal is identified.

Test 2: add acidified potassium dichromate(VI) ($K_2Cr_2O_7$ with dilute H_2SO_4) to a sample of each compound and warm.

Butanal and butan-2-ol (a secondary alcohol) are oxidised, so the solution turns from orange to green. Butanone and 2-methylpropan-2-ol (a tertiary alcohol) do not react, so the solution stays orange.

Butanal is already identified by Test 1, so the other compound that turns green is butan-2-ol.



This leaves butanone and 2-methylpropan-2-ol. These can be distinguished using infrared spectroscopy.

Butanone has a strong $C=O$ absorption at 1680 to 1750 cm^{-1} and no broad $O-H$ absorption.


2-methylpropan-2-ol has a broad $O-H$ (alcohol) absorption at 3230 to 3550 cm^{-1} and no $C=O$ absorption at 1680 to 1750 cm^{-1} .

(Alternatively, mass spectrometry: the molecular ion peak for butanone is at $m/z = 72.0$, but for 2-methylpropan-2-ol it is at $m/z = 74.0$.)



0	8
---	---

Methanol can be manufactured in a reversible reaction as shown by the equation.



0	8	.	1
---	---	---	---

State and explain the effect of using a catalyst on the yield of methanol in this equilibrium.

[2 marks]

There is no effect on the yield of methanol.

A catalyst increases the rate of both the forward and reverse reactions equally (by the same amount), so the position of equilibrium does not change; equilibrium is just reached faster.

0	8	.	2
---	---	---	---

Give an expression for the equilibrium constant (K_c) for this reaction.

[1 mark]

$$K_c = \frac{[\text{CH}_3\text{OH}]}{[\text{CO}][\text{H}_2]^2}$$



08.3

A mixture of carbon monoxide and hydrogen was allowed to reach equilibrium in a container of volume 250 cm^3 at temperature T .
At equilibrium, the mixture contained 0.340 mol of carbon monoxide, 0.190 mol of hydrogen and 0.0610 mol of methanol.

Calculate the value of the equilibrium constant (K_c) for this reaction at temperature T .

[3 marks]

$$[\text{CO}] = 0.340 \div 0.250 = 1.36 \text{ mol dm}^{-3}$$

$$[\text{H}_2] = 0.190 \div 0.250 = 0.760 \text{ mol dm}^{-3}$$

$$[\text{CH}_3\text{OH}] = 0.0610 \div 0.250 = 0.244 \text{ mol dm}^{-3}$$

$$K_c = 0.244 \div (1.36 \times 0.760^2) = 0.3106$$

$$K_c \underline{\hspace{2cm} 0.311 \hspace{2cm}} \text{ mol}^{-2} \text{ dm}^6$$

08.4

Methanol decomposes on heating in a reaction that is the reverse of that used in its manufacture.



Use your answer from Question **08.3** to determine the value of K_c for this equilibrium at temperature T .

State the units for this value of K_c .

(If you were unable to complete the calculation in Question **08.3**, assume a value of $K_c = 0.825 \text{ mol}^{-2} \text{ dm}^6$. This is **not** the correct value.)

[2 marks]

$$K_c (\text{reverse}) = 1 \div K_c (\text{forward}) = 1 \div 0.311 = 3.22$$

$$\text{Value of } K_c \underline{\hspace{2cm} 3.22 \hspace{2cm}}$$

$$\text{Units of } K_c \underline{\hspace{2cm} \text{mol}^2 \text{ dm}^{-6} \hspace{2cm}}$$

Turn over ►



Section B

Answer **all** questions in this section.Only **one** answer per question is allowed.

For each answer completely fill in the circle alongside the appropriate answer.

CORRECT METHOD



WRONG METHODS



If you want to change your answer you must cross out your original answer as shown.

If you wish to return to an answer previously crossed out, ring the answer you now wish to select as shown.





You may do your working in the blank space around each question but this will not be marked.
Do **not** use additional sheets for this working.**0 9**A student has a 10 cm^3 sample of $1.00 \times 10^{-2} \text{ mol dm}^{-3}$ methanoic acid solution. The student is asked to dilute the methanoic acid solution to a concentration of $2.00 \times 10^{-4} \text{ mol dm}^{-3}$ by adding distilled water.

Which volume of water should be added?


[1 mark]**A** 200 cm^3 **B** 490 cm^3 **C** 500 cm^3 **D** 510 cm^3 **1 0**Which molecule does **not** have a permanent dipole?**[1 mark]****A** CH_3Br **B** CH_2Br_2 **C** CHBr_3 **D** CBr_4 

1 1

Which is the major product of the reaction between 2-methylbut-2-ene and iodine monochloride (ICl)?

[1 mark]☐☐☒☐**1 2**

Which statement is **not** correct about the industrial preparation of ethanol by the hydration of ethene at 300 °C?

**[1 mark]**

A The reaction is catalysed by an acid.

☐

B The higher the pressure, the higher the equilibrium yield of ethanol.

☐

C The higher the temperature, the higher the equilibrium yield of ethanol.

☒

D A low equilibrium yield of ethanol is acceptable because unreacted ethene is recycled.

☐**Turn over ►**

1 3 Which compound has the highest boiling point?

[1 mark]

A butanal

☐

B butan-2-ol

☒

C but-2-ene

☐

D 1-fluorobutane

☐

1 4 Which statement is correct about the fractional distillation of crude oil?

[1 mark]

A A zeolite catalyst is used.

☐

B Each fraction contains a mixture of hydrocarbons.

☒

C Gaseous fractions are formed by breaking covalent bonds.

☐

D The fractionating column is hottest at the top.

☐

1 5 How many structural isomers with an unbranched carbon chain have the molecular formula $C_4H_8Br_2$?

[1 mark]

A 4

☐

B 5

☐

C 6

☒


D 7

☐

1 6

The Maxwell–Boltzmann distribution of molecular energies in a sample of gas at a fixed temperature is shown.

Which letter represents the mean energy of the molecules?

[1 mark]**A**☐**B**☐**C**☐**D**☒**1 7**

Ethanol can be made from glucose by fermentation.



In an experiment, 268 g of ethanol ($M_r = 46.0$) were made from 1.44 kg of glucose ($M_r = 180.0$).

What is the percentage yield?

[1 mark]**A** 18.6%☐**B** 36.4%☒**C** 51.1%☐**D** 72.8%☐**Turn over ►**

1 8 Which species could act as a nucleophile?

[1 mark]

A BH_3

☐

B NH_4^+

☐

C PH_3

☒

D SiH_4

☐

1 9 Which statement is correct about poly(chloroethene)?

[1 mark]

A It has the empirical formula CHCl

☐

B It decolourises bromine water.

☐

C Its brittleness is reduced by plasticisers.

☒

D Its polymer chain contains alternate single and double bonds.

☐

2 0 What is the enthalpy of formation of buta-1,3-diene, $\text{C}_4\text{H}_6(\text{g})$?

Substance	Enthalpy of combustion / kJ mol^{-1}
$\text{C}_4\text{H}_6(\text{g})$	-2546
$\text{C}(\text{s})$	-394
$\text{H}_2(\text{g})$	-286

[1 mark]

A $+112 \text{ kJ mol}^{-1}$

☒

B -112 kJ mol^{-1}

☐

C $+746 \text{ kJ mol}^{-1}$

☐

D -746 kJ mol^{-1}

☐


2 1 A gas cylinder contains 5.0 kg of propane.

How many propane molecules are in the cylinder?

The Avogadro constant, $L = 6.022 \times 10^{23} \text{ mol}^{-1}$

[1 mark]

A 6.8×10^{22}

☐

B 7.2×10^{22}

☐

C 6.8×10^{25}

☒

D 7.2×10^{25}

☐

2 2 Which sample of liquid has the greatest volume?

[1 mark]

A 500 mg of pentane (density = 0.63 g cm^{-3})

☐

B 650 mg of propan-1-ol (density = 0.80 g cm^{-3})

☐

C 1.20 g of dichloromethane (density = 1.33 g cm^{-3})

☒

D 1.30 g of trichloromethane (density = 1.48 g cm^{-3})

☐

2 3 Which equation represents an initiation step?

[1 mark]

A $\text{CH}_3\text{CH}_2\dot{\text{C}}\text{HBr} + \text{Br}_2 \longrightarrow \text{CH}_3\text{CH}_2\text{CHBr}_2 + \dot{\text{Br}}$

☐

B $\text{O}_3 + \dot{\text{Cl}} \longrightarrow \text{O}_2 + \dot{\text{ClO}}$

☐

C $\text{RCH}_2\dot{\text{C}}\text{H}_2 + \text{H}_2\text{C}=\text{CH}_2 \longrightarrow \text{RCH}_2\text{CH}_2\text{CH}_2\dot{\text{C}}\text{H}_2$

☐

D $\text{CH}_3\text{CFCl}_2 \longrightarrow \text{CH}_3\dot{\text{C}}\text{FCl} + \dot{\text{Cl}}$

☒

END OF QUESTIONS



There are no questions printed on this page

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2018 AQA and its licensors. All rights reserved.

