

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

StemUp Model Solution

Forename(s)

Candidate signature

AS CHEMISTRY

Paper 1 Inorganic and Physical Chemistry

Monday 20 May 2019

Morning

Time allowed: 1 hour 30 minutes

Materials

For this paper you must have:

- the Periodic Table/Data Sheet, provided as an insert (enclosed)
- a ruler with millimetre measurements
- a scientific calculator, which you are expected to use where appropriate.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do **not** write outside the box around each page or on blank pages.
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

Advice

You are advised to spend about 65 minutes on **Section A** and 25 minutes on **Section B**.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
Section B	
TOTAL	



J U N 1 9 7 4 0 4 1 0 1

There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



Section A

Answer **all** questions in this section.

0 1

This question is about compounds that contain fluorine.

0 1 . 1

Sodium fluoride contains sodium ions (Na^+) and fluoride ions (F^-).
 Na^+ and F^- have the same electron configuration.

Explain why a fluoride ion is larger than a sodium ion.

[2 marks]

*A fluoride ion has fewer protons (9) than a sodium ion (11), so it has
a lower nuclear charge.*

*So there is a weaker attraction between the nucleus and the outer
electrons in the fluoride ion, so the outer electrons are held less closely
and the ion is larger.*

0 1 . 2

Explain, in terms of structure and bonding, why the melting point of sodium fluoride is high.

[2 marks]

*Sodium fluoride has a giant ionic lattice with strong electrostatic forces
of attraction between the oppositely charged ions, Na^+ and F^- .*


*A lot of energy is needed to overcome these strong forces of attraction
(ionic bonds), so the melting point is high.*

Question 1 continues on the next page

Turn over ►

0 1 . 3

The ion H_2F^+ is formed when hydrogen fluoride gains a proton as shown in the equation



Name the type of bond formed when HF reacts with H^+
Explain how this bond is formed.

[2 marks]

Type of bond Coordinate (dative covalent) bond

Explanation A lone pair of electrons on the fluorine atom in HF is donated to the H^+ ion, which has no electrons, so both electrons in the shared pair come from F.



0 1 . 4

Fluoroantimonic acid contains two ions, SbF_6^- and H_2F^+

Draw the shape of the SbF_6^- ion and the shape of the H_2F^+ ion. Include any lone pairs that influence the shape.

Name the shape of each ion.

[4 marks]

	SbF_6^-	H_2F^+
Shape		 (two lone pairs on F)
Name of shape	<u>Octahedral</u>	<u>Bent (V-shaped)</u>



0 1 . 5

Hydrogen fluoride reacts with ethyne (C_2H_2) as shown in the equation. All compounds are in the gaseous state.




Table 1 shows some mean bond enthalpy data.

Table 1

Bond	C-H	C $\equiv$ C	H-F	C-C
Mean bond enthalpy / kJ mol^{-1}	412	837	562	348

Use the data in **Table 1** to calculate a value for the bond enthalpy of a C-F bond in the product.

[3 marks]

$$\Delta H = \Sigma \Delta H(\text{bonds broken}) - \Sigma \Delta H(\text{bonds formed})$$

$$\text{Bonds broken: } 2(\text{C-H}) + \text{C}\equiv\text{C} + 2(\text{H-F})$$

$$= 2(412) + 837 + 2(562) = 2785 \text{ kJ mol}^{-1}$$

$$\text{Bonds formed: } \text{C-C} + 4(\text{C-H}) + 2(\text{C-F})$$

$$= 348 + 4(412) + 2(\text{C-F}) = 1996 + 2(\text{C-F})$$

$$-179 = 2785 - (1996 + 2(\text{C-F}))$$

$$2(\text{C-F}) = 2785 - 1996 + 179 = 968$$

$$\text{C-F} = 968 \div 2 = 484 \text{ kJ mol}^{-1}$$

C-F bond enthalpy 484 kJ mol^{-1}

13

Turn over for the next question

Turn over ►






0 2

Time of flight (TOF) mass spectrometry is an important analytical technique.

A mixture of three compounds is analysed using a TOF mass spectrometer.
The mixture is ionised using electrospray ionisation.

The three compounds are known to have the molecular formulas:



0 2 . 1

Describe how the molecules are ionised using electrospray ionisation.

[3 marks]

The sample is dissolved in a volatile solvent (such as water or methanol).

The solution is injected through a fine hypodermic needle (capillary) held at a high voltage, so it is positively charged.

Each molecule gains a proton (H^+), so it becomes a positive ion, MH^+ , as the solvent evaporates.

0 2 . 2

Give the formula of the ion that reaches the detector first in the TOF mass spectrometer.

[1 mark]



0 2 . 3

A sample of germanium is analysed in a TOF mass spectrometer using electron impact ionisation.

Give an equation, including state symbols, for the process that occurs during the ionisation of a germanium atom.

[1 mark]



0 2 . 4

In the TOF mass spectrometer, a germanium ion reaches the detector in 4.654×10^{-6} s
The kinetic energy of this ion is 2.438×10^{-15} J
The length of the flight tube is 96.00 cm

The kinetic energy of an ion is given by the equation $KE = \frac{1}{2}mv^2$

where

m = mass / kg

v = speed / ms^{-1}

The Avogadro constant $L = 6.022 \times 10^{23} \text{ mol}^{-1}$

Use this information to calculate the mass, in g, of one mole of these germanium ions.
Use your answer to state the mass number of this germanium ion.

[5 marks]

$$v = \text{length} \div \text{time} = 0.9600 \div (4.654 \times 10^{-6})$$

$$v = 2.0627 \times 10^5 \text{ m s}^{-1}$$

$$KE = \frac{1}{2}mv^2, \text{ so } m = 2KE \div v^2$$

$$m = (2 \times 2.438 \times 10^{-15}) \div (2.0627 \times 10^5)^2$$

$$\text{mass of one ion} = 1.146 \times 10^{-25} \text{ kg}$$

$$\text{mass of 1 mol of ions} = 1.146 \times 10^{-25} \times 6.022 \times 10^{23}$$

$$= 0.06901 \text{ kg}$$

$$= 0.06901 \times 1000 = 69.01 \text{ g}$$

Mass of one mole of germanium ions 69.01 g

Mass number of ion 69

10

Turn over for the next question

Turn over ►



0 3

This question is about chromium and its compounds.

0 3 . 1

Complete the full electron configuration of a chromium atom.

[1 mark]



0 3 . 2

An atom has 2 more protons and 3 more neutrons than an atom of ^{52}Cr .

Deduce the symbol, including the mass number and the atomic number, for this atom.

[1 mark]



0 3 . 3

A sample of chromium contains four isotopes and has a relative atomic mass of 52.09

Table 2 shows the mass number and the percentage abundance of three of these isotopes.

Table 2

Mass number	52	53	54
Abundance (%)	82.8	10.9	2.7

Determine the percentage abundance of the fourth isotope.
Show by calculation that the mass number of this isotope is 50

[3 marks]

Percentage abundance 3.6 %

Calculation

$$\% \text{ of fourth isotope} = 100 - (82.8 + 10.9 + 2.7) = 3.6 \%$$

$$52.09 = [(52 \times 82.8) + (53 \times 10.9) + (54 \times 2.7) + (3.6x)] \div 100$$

$$5209 = 4305.6 + 577.7 + 145.8 + 3.6x = 5029.1 + 3.6x$$

$$3.6x = 5209 - 5029.1 = 179.9$$

$$x = 179.9 \div 3.6 = 49.97, \text{ so the mass number is } 50$$



03.4

Deduce the oxidation state of chromium in the $\text{Cr}_2\text{O}_7^{2-}$ ion.

[1 mark]

 $+6$

03.5

Iodide ions can be oxidised to iodine using $\text{Cr}_2\text{O}_7^{2-}$ ions.

Deduce a half-equation to show the oxidation of iodide ions to iodine.

State symbols are **not** required.

[1 mark]

 $2\text{I}^- \rightarrow \text{I}_2 + 2\text{e}^-$

03.6

Deduce a half-equation for the conversion in acidic solution of $\text{Cr}_2\text{O}_7^{2-}$ ions to Cr^{3+} ions.State symbols are **not** required.

[1 mark]

 $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^- \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$

03.7

Use your answers from questions **03.5** and **03.6** to deduce the overall redox equation for the reaction between iodide ions and acidified $\text{Cr}_2\text{O}_7^{2-}$ ions.State symbols are **not** required.

[1 mark]

 $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{I}^- \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O} + 3\text{I}_2$

9

Turn over for the next question

Turn over ►



0 4

The first ionisation energies of the elements in Period 2 change as the atomic number increases.

Explain the pattern in the first ionisation energies of the elements from lithium to neon.

[6 marks]

General trend: the first ionisation energy generally increases across Period 2 from lithium to neon.

This is because the number of protons increases, so the nuclear charge increases. The outer electron is removed from the same energy level (the second shell) each time, so there is similar shielding by the inner 1s electrons across the period.

So there is a stronger attraction between the nucleus and the outer electrons, which are also pulled closer to the nucleus, and more energy is needed to remove an outer electron.

Deviation 1: boron has a lower first ionisation energy than beryllium.

In beryllium ($1s^2 2s^2$) the electron removed is from the 2s sub-level, but in boron ($1s^2 2s^2 2p^1$) the outer electron is in the 2p sub-level.

The 2p sub-level is higher in energy than the 2s sub-level (and slightly further from the nucleus), so less energy is needed to remove the 2p electron from boron.

Deviation 2: oxygen has a lower first ionisation energy than nitrogen.

In nitrogen ($2p^3$) the three 2p electrons are each in a separate 2p orbital, unpaired. In oxygen ($2p^4$) two of the 2p electrons must pair up in the same 2p orbital. These paired electrons repel each other, so less energy is needed to remove one of them, and so the first ionisation energy of oxygen is lower than that of nitrogen.

After oxygen the general increase continues through fluorine to neon, which has the highest first ionisation energy in Period 2, while lithium has the lowest.




[illegible]

6

0 5

Nitrogen monoxide reacts with chlorine to form nitrosyl chloride (NOCl).



0 5 . 1

1.50 mol of NO are mixed with 1.00 mol of Cl₂ and the mixture is left to reach equilibrium at a given temperature.

The equilibrium mixture contains 0.350 mol of NOCl

Calculate the amount, in moles, of NO and of Cl₂ in the equilibrium mixture.

[2 marks]

Moles of NOCl formed = 0.350, so NO used = 0.350 mol

and Cl₂ used = 0.350 ÷ 2 = 0.175 mol

NO = 1.50 - 0.350 = 1.15 mol; Cl₂ = 1.00 - 0.175 = 0.825 mol

Amount of NO 1.15 mol

Amount of Cl₂ 0.825 mol

0 5 . 2

Give the expression for the equilibrium constant, K_c , for the reaction between nitrogen monoxide and chlorine to form nitrosyl chloride.

[1 mark]

$$K_c = \frac{[\text{NOCl}]^2}{[\text{NO}]^2 [\text{Cl}_2]}$$



0 5 . 3

A different equilibrium mixture is prepared in a flask of volume 800 cm^3 at a different temperature.

At equilibrium this mixture contains 0.850 mol of NO and 0.458 mol of Cl_2

For the reaction at this temperature $K_c = 1.32 \times 10^{-2} \text{ mol}^{-1} \text{ dm}^3$

Determine the amount, in moles, of NOCl in this equilibrium mixture.

[4 marks]

$$[\text{NO}] = 0.850 \div 0.800 = 1.0625 \text{ mol dm}^{-3}$$

$$[\text{Cl}_2] = 0.458 \div 0.800 = 0.5725 \text{ mol dm}^{-3}$$

$$K_c = [\text{NOCl}]^2 \div ([\text{NO}]^2 [\text{Cl}_2])$$

$$[\text{NOCl}]^2 = K_c \times [\text{NO}]^2 \times [\text{Cl}_2]$$

$$= 1.32 \times 10^{-2} \times (1.0625)^2 \times 0.5725$$

$$[\text{NOCl}]^2 = 8.53 \times 10^{-3} \text{ mol}^2 \text{ dm}^{-6}$$

$$[\text{NOCl}] = \sqrt{(8.53 \times 10^{-3})} = 0.0924 \text{ mol dm}^{-3}$$

$$n(\text{NOCl}) = 0.0924 \times 0.800 = 0.0739 \text{ mol}$$

Amount of NOCl 0.0739 mol

7


Turn over for the next question

Turn over ►



0 6

A student does an investigation to determine the relative formula mass, M_r , of a solid unknown diprotic acid, H_2A



- 250 cm³ of aqueous solution are prepared using 1300 mg of H_2A
- A pipette is used to add 25.0 cm³ of 0.112 mol dm⁻³ aqueous sodium hydroxide to a conical flask.
- This aqueous sodium hydroxide is titrated with the acid solution.

The titration results are shown in **Table 3**.

Table 3

	Rough	1	2	3
Final volume / cm ³	27.35	26.75	38.90	35.70
Initial volume / cm ³	0.00	0.35	12.15	9.20
Titre / cm ³	27.35	26.40	26.75	26.50

0 6

. 1

Use the results to calculate the M_r of H_2A

[5 marks]

Concordant titres are 26.40 and 26.50, so mean titre = 26.45 cm³

$$n(NaOH) = 25.0 \times 0.112 \div 1000 = 2.80 \times 10^{-3} \text{ mol}$$

$$n(H_2A) \text{ in mean titre} = 2.80 \times 10^{-3} \div 2 = 1.40 \times 10^{-3} \text{ mol}$$

$$n(H_2A) \text{ in } 250 \text{ cm}^3 = 1.40 \times 10^{-3} \times 250 \div 26.45 = 0.01323 \text{ mol}$$

$$\text{Mass of } H_2A = 1300 \text{ mg} = 1.300 \text{ g}$$

$$M_r = \text{mass} \div \text{moles} = 1.300 \div 0.01323 = 98.2$$

M_r of H_2A 98.2



0 6 . 2

The uncertainty in using the pipette in this experiment is $\pm 0.06 \text{ cm}^3$

Calculate the percentage uncertainty in using the pipette.

[1 mark]

$$\% \text{ uncertainty} = (0.06 \div 25.0) \times 100 = 0.24 \%$$

% uncertainty 0.24 %

0 6 . 3

Before adding the solution from the burette in the rough titration, there was an air bubble below the tap.

At the end of this titration the air bubble was not there.

Explain why this air bubble increases the final burette reading of the rough titration.

[1 mark]

The air bubble takes up volume below the tap. During the titration acid solution fills this space (replaces the bubble), so this volume is read as acid added although it never reached the flask.

0 6 . 4

During the titration the student washed the inside of the conical flask with some distilled water.

Suggest why this washing does **not** give an incorrect result.

[1 mark]

Water does not react with the alkali, so it does not change the number of moles of NaOH in the flask.

8

Turn over for the next question

Turn over ►



0 7

This question is about the reactions of magnesium and its compounds.

0 7 . 1

Magnesium is used in one of the stages in the extraction of titanium.

Give an equation for the reaction between titanium(IV) chloride and magnesium.
State the role of magnesium in this reaction.

[2 marks]

Equation





Role of magnesium Reducing agent (it donates electrons)

0 7 . 2

A mixture of magnesium oxide and magnesium hydroxide has a mass of 3200 mg

This mixture is reacted with carbon dioxide to form magnesium carbonate and water.
The mass of water produced is 210 mg



Calculate the percentage by mass of magnesium oxide in this mixture.

[4 marks]

$$n(\text{H}_2\text{O}) = 0.210 \text{ g} \div 18.0 = 0.0117 \text{ mol}$$

$$n(\text{Mg(OH)}_2) = n(\text{H}_2\text{O}) = 0.0117 \text{ mol}$$

$$M_r \text{ of } \text{Mg(OH)}_2 = 24.3 + 2(16.0 + 1.0) = 58.3$$

$$\text{Mass of } \text{Mg(OH)}_2 = 0.0117 \times 58.3 = 0.680 \text{ g}$$

$$\text{Mass of } \text{MgO} = 3.200 - 0.680 = 2.52 \text{ g}$$

$$\% \text{ MgO} = (2.52 \div 3.200) \times 100 = 78.7 \%$$

% of magnesium oxide 78.7



0 8

The following pairs of compounds, each in aqueous solution, can be distinguished by simple test-tube reactions.

Give a reagent, or combination of reagents, that can be added to the solutions in each pair to distinguish between them in a single reaction.

State what is observed in each case.

0 8

. 1

NaCl(aq) and BaCl₂(aq)

[3 marks]

Reagent Dilute sulfuric acid, H₂SO₄(aq)

Observation with NaCl No visible change

Observation with BaCl₂ White precipitate forms

0 8

. 2

NaCl(aq) and Na₂CO₃(aq)

[3 marks]

Reagent Dilute hydrochloric acid, HCl(aq)

Observation with NaCl No visible change

Observation with Na₂CO₃ Effervescence (fizzing, CO₂ gas given off)

6

Turn over for Section B

Turn over ►



Section B

Answer **all** questions in this section.Only **one** answer per question is allowed.

For each answer completely fill in the circle alongside the appropriate answer.

CORRECT METHOD



WRONG METHODS



If you want to change your answer you must cross out your original answer as shown.

If you wish to return to an answer previously crossed out, ring the answer you now wish to select as shown.

You may do your working in the blank space around each question but this will not be marked.
Do **not** use additional sheets for this working.

0 9

Which sample, measured at room temperature and pressure, contains the greatest number of the stated particles?

[1 mark]

A 1 g of hydrogen molecules



B 1 g of helium atoms

C 1 dm³ of hydrogen moleculesD 1 dm³ of helium atoms

1 0

5.0 g of an oxide of molybdenum contain 4.0 g of molybdenum.

What is the empirical formula of this oxide?

[1 mark]

A MoO₂B Mo₄O₅C Mo₂O₃D Mo₃O₂

1	1
---	---

Which substance has delocalised electrons?

[1 mark]

A graphite

☒

B iodine

☐

C sodium chloride

☐

D tetrachloromethane

☐

1	2
---	---

Which species is **not** pyramidal in shape?

[1 mark]

A PF_3 ☐B H_3O^+ ☐C CH_3^- ☐D BF_3 ☒

1	3
---	---

Which change occurs when water is vaporised?

[1 mark]

A An exothermic change occurs.

☐

B Covalent bonds are broken.

☐

C Intermolecular forces are overcome.

☒

D The total energy of the molecules decreases.





☐

Turn over ►







1 4

Which equation represents the reaction that has a standard enthalpy change equal to the standard enthalpy of formation for barium chloride?

[1 mark]☐☐☒☐**1 5**

Which equation does **not** represent a redox reaction?

[1 mark]☐☐☐☒**1 6**

Which property would you expect the element radium, Ra, to possess?

[1 mark]

A It forms a soluble sulfate.

☐

B It does not react with water.

☐

C It is a good conductor of electricity.

☒

D It forms a covalent fluoride.

☐

1 7Which statement is **not** correct?**[1 mark]**

- A** Strontium has a lower first ionisation energy than calcium.
- B** Strontium has a larger ionic radius than calcium.
- C** Strontium reacts less vigorously with water than calcium.
- D** Strontium hydroxide is more soluble in water than calcium hydroxide.

☐☐☒☐**1 8**

Which property of the Group 2 elements, Ca to Ba, increases with increasing atomic number?

[1 mark]

- A** Atomic Radius
- B** Electronegativity
- C** First ionisation energy
- D** Melting Point

☒☐☐☐**1 9**

What is the best oxidising agent?

[1 mark]

- A** F_2
- B** F^-
- C** I_2
- D** I^-

☒☐☐☐**Turn over ►**

2 0

Some fuel in a spirit burner is burned, and the heat produced is used to heat a container of water.

In this experiment:

The mass of water heated = m g

The temperature rise = y °C

The specific heat capacity of water = c J K⁻¹ g⁻¹

What is the amount of heat energy absorbed by the water?

[1 mark]

A mcy

☒

B $mc(y + 273)$

☐


C y / mc

☐

D $(y + 273) / mc$

☐
2 1

The equation below represents the complete combustion of butane.



20 cm³ of butane are completely burned in 0.20 dm³ of oxygen.
Which statement is correct?

All volumes are measured at the same temperature and pressure.

[1 mark]

A 40 cm³ of carbon dioxide are formed

☐

B 0.065 dm³ of oxygen react

☐

C 70 cm³ of oxygen remain

☒

D 0.50 dm³ of steam are formed

☐


2 2


Which statement is correct about reactions involving halide ions?

[1 mark]

- A** Sodium chloride forms chlorine when added to concentrated sulfuric acid.
- B** Sodium chloride forms chlorine when added to bromine.
- C** Sodium bromide forms bromine when added to concentrated sulfuric acid.
- D** Sodium bromide forms bromine when added to iodine.

☐☐☒☐**2 3**

What is the percentage yield when 20 g of aluminium are produced from 50 g of aluminium oxide?

**[1 mark]**

- A** 76%
- B** 40%
- C** 33%
- D** 19%

☒☐☐☐

15**END OF QUESTIONS**

There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third-party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2019 AQA and its licensors. All rights reserved.



2 4



1 9 6 A 7 4 0 4 / 1