AQA

Please write clearly in block capitals.

Centre number | | | | | | Candidate number | | | | |

Surname StemUp Model Solution

Forename(s) _____

Candidate signature _____

AS
FURTHER MATHEMATICS

Paper 1

Monday 14 May 2018 Afternoon Time allowed: 1 hour 30 minutes

Materials
- You must have the AQA formulae and statistical tables booklet for A-level Mathematics and A-level Further Mathematics.
- You should have a scientific calculator that meets the requirements of the specification. (You may use a graphical calculator.)

Instructions
- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer each question in the space provided for that question. If you require extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do **not** write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information
- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

Advice
- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	
TOTAL	

Do not write outside the box

Answer **all** questions in the spaces provided.

1 $z = 3 - \mathrm{i}$

Determine the value of zz^*

Circle your answer.

[1 mark]

(10) $\sqrt{10}$ $10 - 2\mathrm{i}$ $10 + 2\mathrm{i}$

2 Three matrices **A**, **B** and **C** are given by

$$\mathbf{A} = \begin{bmatrix} 5 & 2 & -3 \\ 0 & 7 & 6 \\ 4 & 1 & 0 \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} 1 & 0 \\ 3 & -5 \\ -2 & 6 \end{bmatrix} \quad \text{and } \mathbf{C} = \begin{bmatrix} 6 & 4 & 3 \\ 1 & 2 & 0 \end{bmatrix}$$

Which of the following **cannot** be calculated?

Circle your answer.

[1 mark]

AB (**AC**) **BC** $\mathbf{A}^2$

3 Which of the following functions has the fourth term $-\frac{1}{720}x^6$ in its Maclaurin series expansion?

Circle your answer.

[1 mark]

$\sin x$ ($\cos x$) e^x $\ln(1 + x)$


Do not write outside the box

4 Sketch the graph given by the polar equation

$$r = \frac{a}{\cos\theta}$$

where a is a positive constant.

[2 marks]



5 Describe fully the transformation given by the matrix $\begin{bmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} & 0 \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix}$

[3 marks]

$\cos\theta = -\frac{1}{2}$, $\sin\theta = \frac{\sqrt{3}}{2}$ (top-left 2 × 2 part)

$\Rightarrow \theta = 120°$ (second quadrant)

Third row and column are 0, 0, 1 so the z-coordinate is unchanged.

Rotation through 120° anticlockwise about the z-axis

Do not write outside the box

6 (a) Matthew is finding a formula for the inverse function $\operatorname{arsinh} x$.
He writes his steps as follows:

$$\text{Let } y = \sinh x$$

$$y = \frac{1}{2}(e^x - e^{-x})$$

$$2y = e^x - e^{-x}$$

$$0 = e^x - 2y - e^{-x}$$

$$0 = (e^x)^2 - 2ye^x - 1$$

$$0 = (e^x - y)^2 - y^2 - 1$$

$$y^2 + 1 = (e^x - y)^2$$

$$\pm\sqrt{y^2 + 1} = e^x - y$$

$$y \pm \sqrt{y^2 + 1} = e^x$$

To find the inverse function, swap x and y: $x \pm \sqrt{x^2 + 1} = e^y$

$$\ln\left(x \pm \sqrt{x^2 + 1}\right) = y$$

$$\operatorname{arsinh} x = \ln\left(x \pm \sqrt{x^2 + 1}\right)$$

Identify, and explain, the error in Matthew's proof.

[2 marks]

The error is the ± sign: only the + sign is valid.

$e^x > 0$ for all x, but $\sqrt{(y^2 + 1)} > |y|$ so $y - \sqrt{(y^2 + 1)} < 0$

so $e^x = y - \sqrt{(y^2 + 1)}$ is impossible (a contradiction).

Hence $e^x = y + \sqrt{(y^2 + 1)}$ and $\operatorname{arsinh} x = \ln(x + \sqrt{(x^2 + 1)})$

Do not write outside the box

6 (b) Solve $\ln\left(x + \sqrt{x^2 + 1}\right) = 3$

[1 mark]

$x + \sqrt{(x^2 + 1)} = e^3 \Rightarrow \operatorname{arsinh} x = 3 \Rightarrow x = \sinh 3$

$x = \frac{1}{2}(e^3 - e^{-3}) = 10.0178...$

$x = 10.0$

7 Find two invariant points under the transformation given by $\begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}$

[2 marks]

$\begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ y \end{bmatrix}$

$2x + 3y = x \Rightarrow x + 3y = 0$

$x + 4y = y \Rightarrow x + 3y = 0$

Invariant points lie on the line $x = -3y$

Two invariant points: $(0, 0)$ and $(-3, 1)$

Do not write outside the box

8 $2 - 3\mathrm{i}$ is one root of the equation

$$z^3 + mz + 52 = 0$$

where m is real.

8 (a) Find the other roots.

[3 marks]

Coefficients are real, so the conjugate $2 + 3i$ is also a root.

No z^2 term, so the sum of the roots = 0

$\alpha + (2 - 3i) + (2 + 3i) = 0$

$\alpha + 4 = 0 \Rightarrow \alpha = -4$

(Check: product of roots $= \alpha(2 - 3i)(2 + 3i) = 13\alpha = -52 \Rightarrow \alpha = -4$)

Other roots: $2 + 3i$ and -4

Do not write outside the box

8 (b) Determine the value of m.

[2 marks]

$z = -4$ is a root: $(-4)^3 + m(-4) + 52 = 0$

$-64 - 4m + 52 = 0$


$-12 = 4m$

$m = -3$

Do not write outside the box

9 (a) Sketch the graph of $y^2 = 4x$

[1 mark]



9 (b) Ben is using a 3D printer to make a plastic bowl which holds exactly 1000 cm^3 of water.

Ben models the bowl as a region which is rotated through 2π radians about the x-axis.

He uses the finite region enclosed by the lines $x = d$ and $y = 0$ and the curve with equation $y^2 = 4x$ for $y \geq 0$

9 (b) (i) Find the depth of the bowl to the nearest millimetre.

[4 marks]

$V = \pi\int_0^d y^2\,dx = \pi\int_0^d 4x\,dx = \pi[\,2x^2\,]_0^d = 2\pi d^2$

$2\pi d^2 = 1000$

$d^2 = \frac{500}{\pi}$

$d = \sqrt{(500/\pi)} = 12.6157\ldots$ cm

Depth of bowl = 12.6 cm (126 mm)

Do not write outside the box

9 (b) (ii) What assumption has Ben made about the bowl?

[1 mark]

The thickness of the plastic is negligible (the bowl has zero thickness).

Turn over ▶

Do not write outside the box

10 (a) Prove by induction that, for all integers $n \geq 1$,

$$\sum_{r=1}^{n} r^3 = \frac{1}{4}n^2(n+1)^2$$

[4 marks]

$n = 1$: LHS $= 1^3 = 1$, RHS $= \frac{1}{4} \times 1^2 \times 2^2 = 1$, so true for $n = 1$

Assume true for $n = k$: $\sum_{r=1}^{k} r^3 = \frac{1}{4}k^2(k+1)^2$

Then $\sum_{r=1}^{k+1} r^3 = \frac{1}{4}k^2(k+1)^2 + (k+1)^3$

$= \frac{1}{4}(k+1)^2[k^2 + 4(k+1)]$

$= \frac{1}{4}(k+1)^2(k^2 + 4k + 4)$

$= \frac{1}{4}(k+1)^2(k+2)^2 = \frac{1}{4}(k+1)^2((k+1)+1)^2$

which is the formula with $n = k + 1$, so it is true for $n = k + 1$.

The result is true for $n = 1$, and if true for $n = k$ then it is true for $n = k + 1$, so by induction it is true for all integers $n \geq 1$.

Do not write outside the box

10 (b) **Hence** show that

$$\sum_{r=1}^{2n} r(r-1)(r+1) = n(n+1)(2n-1)(2n+1)$$

[4 marks]

$\Sigma_{r=1}^{2n}\ r(r-1)(r+1) = \Sigma_{r=1}^{2n}\ (r^3 - r) = \Sigma_{r=1}^{2n}\ r^3 - \Sigma_{r=1}^{2n}\ r$

$= \frac{1}{4}(2n)^2(2n+1)^2 - \frac{1}{2}(2n)(2n+1)$

$= n^2(2n+1)^2 - n(2n+1)$

$= n(2n+1)[\,n(2n+1) - 1\,]$

$= n(2n+1)(2n^2 + n - 1)$

$= n(2n+1)(2n-1)(n+1)$

$= n(n+1)(2n-1)(2n+1)$ as required


Turn over ▶

Do not write outside the box

11 Four finite regions *A*, *B*, *C* and *D* are enclosed by the curve with equation

$$y = x^3 - 7x^2 + 11x + 6$$

and the lines $y = k$, $x = 1$ and $x = 4$, as shown in the diagram below.



The areas of *B* and *C* are equal.

Find the value of k.

[3 marks]

B is above y = k and C is below it for $1 \leq x \leq 4$, so area B = area C gives

$\int_1^4 (x^3 - 7x^2 + 11x + 6 - k)\,dx = 0$

$\left[\frac{x^4}{4} - \frac{7x^3}{3} + \frac{11x^2}{2} + 6x - kx\right]_1^4 = 0$

$\left(64 - \frac{448}{3} + 88 + 24 - 4k\right) - \left(\frac{1}{4} - \frac{7}{3} + \frac{11}{2} + 6 - k\right) = 0$

$\left(\frac{80}{3} - 4k\right) - \left(\frac{113}{12} - k\right) = 0$

$\frac{69}{4} - 3k = 0$

$k = \frac{23}{4} = 5.75$

Do not write outside the box

12 (a) Show that the matrix $\begin{bmatrix} 5-k & 2 \\ k^3+1 & k \end{bmatrix}$ is singular when $k = 1$.

[1 mark]

$k = 1$: $\det = (5 - 1)(1) - 2(1^3 + 1) = 4 - 4 = 0$

$\det = 0$, so the matrix is singular when $k = 1$

12 (b) Find the values of k for which the matrix $\begin{bmatrix} 5-k & 2 \\ k^3+1 & k \end{bmatrix}$ has a negative determinant.

Fully justify your answer.

[5 marks]

$\det = k(5 - k) - 2(k^3 + 1) = 5k - k^2 - 2k^3 - 2$

Negative determinant: $5k - k^2 - 2k^3 - 2 < 0$

$\Rightarrow 2k^3 + k^2 - 5k + 2 > 0$

$k = 1$ is a root (part (a)): $(k - 1)(2k^2 + 3k - 2) > 0$

$(k - 1)(2k - 1)(k + 2) > 0$

Critical values: $k = -2, \frac{1}{2}, 1$

$k < -2$: $-$, $-2 < k < \frac{1}{2}$: $+$, $\frac{1}{2} < k < 1$: $-$, $k > 1$: $+$

$\therefore -2 < k < \frac{1}{2}$ or $k > 1$

Do not write outside the box

13 The graph of the rational function $y = f(x)$ intersects the x-axis exactly once at $(-3, 0)$

The graph has exactly two asymptotes, $y = 2$ and $x = -1$

13 (a) Find $f(x)$

[2 marks]


Asymptotes $x = -1$ and $y = 2 \Rightarrow f(x) = \frac{2x + c}{x + 1}$

$f(-3) = 0$: $\frac{-6 + c}{-2} = 0 \Rightarrow c = 6$

$f(x) = \frac{2x + 6}{x + 1}$ $\left(= 2 + \frac{4}{x + 1}\right)$

13 (b) Sketch the graph of the function.

[3 marks]



Do not write outside the box

13 (c) Find the range of values of x for which $\mathrm{f}(x) \leq 5$

[4 marks]

$\frac{2x + 6}{x + 1} \leq 5 \qquad (x \neq -1)$

Multiply by $(x + 1)^2 > 0$: $(2x + 6)(x + 1) \leq 5(x + 1)^2$

$0 \leq (x + 1)[\, 5(x + 1) - (2x + 6)\,]$

$0 \leq (x + 1)(3x - 1)$

Critical values: $x = -1$ and $x = \frac{1}{3}$ $\quad (f(x) = 5 \Rightarrow 5x + 5 = 2x + 6 \Rightarrow x = \frac{1}{3})$

$(x + 1)(3x - 1) \geq 0$ for $x \leq -1$ or $x \geq \frac{1}{3}$, but $x \neq -1$


$\therefore \quad x < -1 \quad$ or $\quad x \geq \frac{1}{3}$

Do not write outside the box

14 (a) Sketch, on the Argand diagram below, the locus of points satisfying the equation

$$|z - 3| = 2$$

[1 mark]



Do not write outside the box

14 (b) There is a unique complex number w that satisfies both

$$|w - 3| = 2 \quad \text{and} \quad \arg(w + 1) = \alpha$$

where α is a constant such that $0 < \alpha < \pi$

14 (b) (i) Find the value of α.

[2 marks]

A unique w means the half-line from −1 is a tangent to the circle.

Distance from −1 to the centre 3 is 4, and the radius is 2

$\sin \alpha = \frac{2}{4} = \frac{1}{2} \Rightarrow \alpha = \frac{\pi}{6}$

14 (b) (ii) Express w in the form $r(\cos\theta + \mathrm{i}\sin\theta)$.

Give each of r and θ to two significant figures.

[4 marks]

Length of tangent from −1 = $\sqrt{(4^2 - 2^2)} = \sqrt{12} = 2\sqrt{3}$

$w = -1 + 2\sqrt{3}(\cos\frac{\pi}{6} + i\sin\frac{\pi}{6})$

$= -1 + 2\sqrt{3}(\frac{\sqrt{3}}{2} + \frac{1}{2}i) = -1 + 3 + \sqrt{3}\,i = 2 + \sqrt{3}\,i$

$r = |w| = \sqrt{(2^2 + (\sqrt{3})^2)} = \sqrt{7} = 2.6457\ldots = 2.6$

$\theta = \arg w = \tan^{-1}(\frac{\sqrt{3}}{2}) = 0.71372\ldots = 0.71$

$w = 2.6(\cos 0.71 + i \sin 0.71)$

Turn over ▶

Do not write outside the box

15 (a) Show that

$$\frac{1}{r+2}-\frac{1}{r+3}=\frac{1}{(r+2)(r+3)}$$

[1 mark]

$\frac{1}{r+2}-\frac{1}{r+3}=\frac{(r+3)-(r+2)}{(r+2)(r+3)}=\frac{1}{(r+2)(r+3)}$

15 (b) Use the method of differences to show that

$$\sum_{r=1}^{n}\frac{1}{(r+2)(r+3)}=\frac{n}{3(n+3)}$$

[3 marks]

$\sum_{r=1}^{n}\frac{1}{(r+2)(r+3)}=\sum_{r=1}^{n}\left(\frac{1}{r+2}-\frac{1}{r+3}\right)$

$r=1$: $\frac{1}{3}-\frac{1}{4}$

$r=2$: $\frac{1}{4}-\frac{1}{5}$

$r=3$: $\frac{1}{5}-\frac{1}{6}$

⋮

$r=n-1$: $\frac{1}{n+1}-\frac{1}{n+2}$

$r=n$: $\frac{1}{n+2}-\frac{1}{n+3}$

Adding, all the middle terms cancel: sum $=\frac{1}{3}-\frac{1}{n+3}$

$=\frac{n+3-3}{3(n+3)}=\frac{n}{3(n+3)}$ as required

Do not write outside the box

16 Two matrices **A** and **B** satisfy the equation

$$\mathbf{AB} = \boldsymbol{I} + 2\mathbf{A}$$

where $\boldsymbol{I}$ is the identity matrix and $\mathbf{B} = \begin{bmatrix} 3 & -2 \\ -4 & 8 \end{bmatrix}$

Find **A**.

[3 marks]

$AB - 2A = I \Rightarrow A(B - 2I) = I \Rightarrow A = (B - 2I)^{-1}$

$B - 2I = \begin{bmatrix} 3 & -2 \\ -4 & 8 \end{bmatrix} - \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} = \begin{bmatrix} 1 & -2 \\ -4 & 6 \end{bmatrix}$

$\det(B - 2I) = 1 \times 6 - (-2)(-4) = 6 - 8 = -2$

$A = -\frac{1}{2}\begin{bmatrix} 6 & 2 \\ 4 & 1 \end{bmatrix} = \begin{bmatrix} -3 & -1 \\ -2 & -0.5 \end{bmatrix}$

Do not write outside the box

17 Find the exact solution to the equation

$$\sinh\theta(\sinh\theta + \cosh\theta) = 1$$

[4 marks]

$\sinh\theta = \frac{1}{2}(e^{\theta} - e^{-\theta})$, $\cosh\theta = \frac{1}{2}(e^{\theta} + e^{-\theta})$

so $\sinh\theta + \cosh\theta = e^{\theta}$

$\frac{1}{2}(e^{\theta} - e^{-\theta}) \times e^{\theta} = 1$

$e^{2\theta} - 1 = 2$

$e^{2\theta} = 3$

$2\theta = \ln 3$

$\theta = \frac{1}{2}\ln 3$

Do not write outside the box

18 α, β and γ are the real roots of the cubic equation

$$x^3 + mx^2 + nx + 2 = 0$$

By considering $(\alpha - \beta)^2 + (\gamma - \alpha)^2 + (\beta - \gamma)^2$, prove that

$$m^2 \geq 3n$$

[4 marks]

$(\alpha - \beta)^2 + (\gamma - \alpha)^2 + (\beta - \gamma)^2$

$= \alpha^2 - 2\alpha\beta + \beta^2 + \gamma^2 - 2\gamma\alpha + \alpha^2 + \beta^2 - 2\beta\gamma + \gamma^2$

$= 2(\alpha^2 + \beta^2 + \gamma^2) - 2(\alpha\beta + \beta\gamma + \gamma\alpha) = 2\Sigma\alpha^2 - 2\Sigma\alpha\beta$

$\Sigma\alpha = -m$, $\Sigma\alpha\beta = n$

$\Sigma\alpha^2 = (\Sigma\alpha)^2 - 2\Sigma\alpha\beta = m^2 - 2n$

$\therefore (\alpha - \beta)^2 + (\gamma - \alpha)^2 + (\beta - \gamma)^2 = 2(m^2 - 2n) - 2n = 2m^2 - 6n$

α, β, γ are real, so each of $(\alpha - \beta)^2$, $(\gamma - \alpha)^2$, $(\beta - \gamma)^2$ is ≥ 0

$\therefore (\alpha - \beta)^2 + (\gamma - \alpha)^2 + (\beta - \gamma)^2 \geq 0$

$\Rightarrow \quad 2m^2 - 6n \geq 0 \quad \Rightarrow \quad m^2 \geq 3n$

Do not write outside the box

19 A theme park has two zip wires.

Sarah models the two zip wires as straight lines using coordinates in metres.

The ends of one wire are located at (0, 0, 0) and (0, 100, −20)

The ends of the other wire are located at (10, 0, 20) and (−10, 100, −5)

19 (a) Use Sarah's model to find the shortest distance between the zip wires.

[7 marks]

Wire 1 direction: $d_1 = \begin{bmatrix} 0 \\ 100 \\ -20 \end{bmatrix} - \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 100 \\ -20 \end{bmatrix}$

Wire 2 direction: $d_2 = \begin{bmatrix} 10 \\ 0 \\ 20 \end{bmatrix} - \begin{bmatrix} -10 \\ 100 \\ -5 \end{bmatrix} = \begin{bmatrix} 20 \\ -100 \\ 25 \end{bmatrix}$

Let $n = (x, y, z)$ be perpendicular to both wires:

$n \cdot d_1 = 0$: $100y - 20z = 0 \Rightarrow z = 5y$

$n \cdot d_2 = 0$: $20x - 100y + 25z = 0 \Rightarrow 20x + 25y = 0 \Rightarrow x = -1.25y$

Take $y = -4$: $n = \begin{bmatrix} 5 \\ -4 \\ -20 \end{bmatrix}$, $|n| = \sqrt{(5^2 + 4^2 + 20^2)} = \sqrt{441} = 21$

Vector joining the wires: $\begin{bmatrix} 10 \\ 0 \\ 20 \end{bmatrix} - \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 10 \\ 0 \\ 20 \end{bmatrix}$

Shortest distance $= \frac{1}{21} \left| \begin{bmatrix} 10 \\ 0 \\ 20 \end{bmatrix} \cdot \begin{bmatrix} 5 \\ -4 \\ -20 \end{bmatrix} \right|$

$= \frac{|50 + 0 - 400|}{21} = \frac{350}{21}$

$= 16.666...$

Shortest distance = 16.7 m

Do not write outside the box

19 (b) State one way in which Sarah's model could be refined.

[1 mark]

Model the wires as curves rather than straight lines, since real zip wires sag under their own weight and the weight of the rider.

END OF QUESTIONS

Do not write outside the box

There are no questions printed on this page

DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2018 AQA and its licensors. All rights reserved.