Please write clearly in block capitals.

Centre number | | | | | | Candidate number | | | | |

Surname _____

Forename(s) _____

Candidate signature _____

I declare this is my own work.

AS
FURTHER MATHEMATICS

Paper 2 Mechanics

Friday 16 May 2025 Afternoon Time allowed: 1 hour 30 minutes

Materials
- You must have the AQA Formulae and statistical tables booklet for A-level Mathematics and A-level Further Mathematics.
- You should have a graphical or scientific calculator that meets the requirements of the specification.
- You must ensure you have the other optional Question Paper/Answer Book for which you are entered (**either** Discrete **or** Statistics). You will have 1 hour 30 minutes to complete **both** papers.

Instructions
- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer each question in the space provided for that question. If you require extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do **not** write outside the box around each page or on blank pages.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

Information
- The marks for questions are shown in brackets.
- The maximum mark for this paper is 40.

Advice
- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
TOTAL	

Do not write outside the box

Answer **all** questions in the spaces provided.

1 A particle travels around a horizontal circle of radius r metres with a constant angular speed of ω radians per second.

State an expression for the magnitude of the acceleration, in m s^{-2}, of the particle.

Circle your answer.

[1 mark]

$r^2\omega$ $\frac{\omega}{r^2}$ $r\omega^2$ $\frac{\omega^2}{r}$

2 State the dimensions of power.

Circle your answer.

[1 mark]

ML^2T ML^2T^{-1} ML^2T^{-2} ML^2T^{-3}

Do not write outside the box

3 A light elastic string has a modulus of elasticity of 20 newtons and a natural length of 0.5 metres.

One end of the string is attached to a fixed point.

A particle of weight 25 newtons is attached to the other end of the string.

The particle hangs in equilibrium with the string vertical.

Find the extension of the string.

[2 marks]

Turn over for the next question

Do not write outside the box

4 Two particles, P and Q, are moving on a smooth horizontal surface.

Particle P has mass 4 kg and is moving with velocity $\begin{bmatrix} -1 \\ 2 \end{bmatrix}$ m s^{-1}

Particle Q has mass 6 kg and is moving with velocity $\begin{bmatrix} 4 \\ -3 \end{bmatrix}$ m s^{-1}

The particles collide and coalesce to form a single particle.

Find the velocity of the single particle immediately after the collision.

[2 marks]

Do not write outside the box

5 The path of a person on a theme park ride can be modelled as a circle of radius 15 metres.

The person has a constant angular speed about the centre of the circle.

The person completes one revolution every 40 seconds.

Find the speed of the person on the ride in metres per second.

Give your answer to two significant figures.

[3 marks]

Turn over for the next question

Do not write outside the box

6 A ball of mass m is thrown vertically upwards with initial speed u

The ball travels a vertical distance h before instantaneously coming to rest.

It is thought that

$$h = km^a u^b g^c$$

where

- $[k] = 1$
- g is the acceleration due to gravity
- a, b and c are constants.

6 (a) Explain what is meant by $[k] = 1$

[1 mark]

6 (b) Use dimensional analysis to find the value of a, the value of b and the value of c

[3 marks]

Do not write outside the box

7 A car of mass 900 kg is travelling along a straight horizontal road.

The car experiences a constant resistance force of 500 newtons.

7 (a) At a particular time, the car has speed 10 m s^{-1} and is accelerating at 0.6 m s^{-2}

7 (a) (i) Show that the driving force has magnitude 1040 newtons at this time.

[2 marks]

7 (a) (ii) Find the power of the car at this time.

Give your answer in kW

[2 marks]

7 (b) It is given that the maximum speed of the car is 30 m s^{-1} on this horizontal road.

Find the maximum power of the engine.

[1 mark]

Do not write outside the box

8 A horizontal force, F newtons, is acting on a particle moving in a straight line on a smooth horizontal surface.

At time t seconds, $F = 20t + 2$ where $0 \leq t \leq T$

The magnitude of the impulse exerted on the particle by the horizontal force over the interval $0 \leq t \leq T$ is 168 N s

8 (a) (i) Show that

$$5T^2 + T - 84 = 0$$

[3 marks]

8 (a) (ii) Find the value of T

[1 mark]

Do not write outside the box

8 (b) It is given that the velocity of the particle increases from 10 m s^{-1} to 30 m s^{-1} over the interval $0 \leq t \leq T$

Find the mass of the particle.

[2 marks]

Turn over for the next question

Do not write outside the box


9 A block of mass 4 kg rests on a horizontal surface.

The block is attached to one end of a light elastic string.

The other end of the string is attached to a vertical wall.

The string has natural length 1.5 metres and a stiffness of 120 N m^{-1}

The block is released from rest at a distance of 2 metres from the wall as shown in the diagram.



Assume that no air resistance acts on the block.

9 (a) Find the elastic potential energy of the string when the block is 2 metres from the wall.

[2 marks]

9 (b) Assuming that the horizontal surface is modelled as smooth, use an energy method to find the speed of the block when it hits the wall.

Give your answer to three significant figures.

[2 marks]

Do not write outside the box

9 (c) In an improved model the horizontal surface is modelled as rough, with a constant friction force of magnitude F newtons.

Given that the block just reaches the wall, find the value of F

[2 marks]

9 (d) State how your answer to part **(c)** would change if air resistance was considered.

[1 mark]


Turn over for the next question

Do not write outside the box

10 Two smooth spheres, *A* and *B*, of equal size, initially lie at rest on a smooth horizontal surface.

Sphere *A* has mass $2m$ and sphere *B* has mass $3m$

Sphere *A* is projected with speed u directly towards sphere *B* as shown in the diagram.



Subsequently *A* collides directly with *B*

The coefficient of restitution between *A* and *B* is $\frac{3}{4}$

10 (a) Find, in terms of u, the speed of *A* and the speed of *B* immediately after the collision.

[5 marks]

Do not write outside the box

10 (b) Subsequently B hits a vertical wall and rebounds, travelling directly towards A

The coefficient of restitution between B and the wall is e

It is given that B **does not** collide again with A

Find the range of possible values of e

[4 marks]

END OF QUESTIONS

Do not write outside the box

There are no questions printed on this page

DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Copyright information

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.aqa.org.uk

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2025 AQA and its licensors. All rights reserved.