

AQA GCSE PHYSICS (HIGHER) • 8463

PHYSICS



EXAM QUESTION PACK

4.5.1.1 SCALAR AND VECTOR QUANTITIES (PACK 1)

SCORE

/66

PERCENTAGE

%

STEMUP

These questions belong to AQA, not to us.

The examination questions, mark schemes and examiner comments in this pack were written and published by AQA. They are past assessment materials, reproduced here so that students can revise from them.

What we do not own

STEMUP is not affiliated with AQA and does not own, and makes no claim to own, any of the question, mark scheme or examiner report material in this pack. AQA has not endorsed, approved or been involved in the production of it. AQA and the qualification titles used here belong to their respective owners and appear for identification only.

Where to find the originals

AQA publishes its past papers, mark schemes and examiner reports. Always check the current specification at aqa.org.uk before you revise, because content and wording can change between series.

If you hold the rights to any material in this pack and would like it amended or removed, contact admin@stemup.co.uk and we will act on it.

Q1.

- (a) Which of these is a scalar quantity?

Tick **one** box.

displacement

☐

distance

☐

force

☐


velocity

☐

(1)

- (b) A woman cycled along a straight flat road.

The graph below shows how the woman's velocity changed with time.



Which part of the graph shows the woman moving at constant velocity?

Tick **one** box.

BC

☐

CD

☐

DE

☐

(1)

- (c) Which part of the graph shows the woman stationary?

BC ☐ CD ☐ DE ☐

(1)

Between points **A** and **B** the woman was accelerating.

- (d) Use the graph above to determine the total time for which she was accelerating.

Time = _____ s

(1)

- (e) Use the graph above to determine her increase in velocity between points **A** and **B**.

Increase in velocity = _____ m/s

(1)

- (f) Calculate her acceleration between points **A** and **B**.

Use the equation:

$$\text{acceleration} = \frac{\text{change in velocity}}{\text{time taken}}$$

Acceleration = _____ m/s²

(2)

- (g) Estimate how a typical cycling speed of 6 m/s compares with a typical walking speed.

Tick **one** box.

about twice as fast ☐

about four times faster ☐

about eight times faster ☐

(1)

(Total 8 marks)

Q2.


A student investigated the frictional force between an object and a surface.

The student used a string to pull a small wooden block across different surfaces. The block was pulled at a constant speed in a straight line.

Pulling the block causes a tension force in the string.

The student kept the angle of the string the same each time.

The diagram below represents the block being pulled across a piece of carpet.



- (a) Measure angle **A** on the diagram above.

Angle **A** = _____ degrees

(1)

- (b) Complete the sentences.

Choose answers from the box.

controlled	dependent	scalar	valid	vector
-------------------	------------------	---------------	--------------	---------------

Force has both magnitude and direction, so is a _____ quantity.

A quantity with magnitude only is a _____ quantity.

(2)

- (c) Two forces acting on the block are tension and friction.

Name **one** other force acting on the block.

(1)

- (d) When the student pulled the block with a constant force, the velocity of the block did not change.

What is the best explanation for this?

Tick **one** box.

Force is directly proportional to velocity

☐

No work is done by the pulling force

☐

The block is moving in a straight line

☐

The resultant force on the block is zero

☐

(1)

The student pulled the block along four different surfaces:

- cardboard
- carpet
- glass
- sandpaper.

(e) Give **two** control variables for this investigation.

1. _____

2. _____

(2)

The table below shows the results.

Surface	Force to pull the block in newtons			Mean force in newtons
	Trial 1	Trial 2	Trial 3	
cardboard	1.4	1.6	1.5	1.5
carpet	2.5	3.0	3.9	3.2
glass	0.7	0.8	0.6	0.7
sandpaper	5.2	5.6	5.4	X

(f) Calculate value **X** in the table above.

X = _____ N

(1)

(g) Which surface produced the lowest friction force?

(1)

(Total 9 marks)

Q3.

Forces are vector quantities.

- (a) What is the difference between a vector quantity and a scalar quantity?


(2)

Figure 1 represents a wooden block being pulled across a surface at a constant speed in a straight line.

The block is in contact with the surface.

The arrow in **Figure 1** represents the tension force in the string pulling the block.

Figure 1




- (b) Complete **Figure 1** to show the other three forces acting on the block.

(3)

Figure 2 is a copy of **Figure 1** to help you answer the following question.

Figure 2



- (c) **Figure 2** is drawn to scale. The scale is 1 cm : 0.5 N

Determine the horizontal and vertical components of the tension in the string.

Show these components on **Figure 2**.

Horizontal component = _____ N

Vertical component = _____ N

(3)

A student collects data on the size of the force required to pull the block across different surfaces at a constant speed.

The table below shows the results.

Type of surface	Force in N			Mean force in N
	Trial 1	Trial 2	Trial 3	
Cardboard	1.4	1.6	1.5	1.5
Carpet	2.6	3.1	3.9	3.2
Glass	0.7	0.8	0.6	0.7
Sandpaper	5.2	X	5.3	5.4

(d) Calculate value **X** in the table above.

X = _____ N

(2)

(e) Give **three** control variables for this investigation.

1. _____

2. _____

3. _____

(3)


(Total 13 marks)

Q4.

Figure 1 shows a rollercoaster train as it is pulled up a slope on the track.

The arrows, **A**, **B**, **C** and **D**, represent the forces acting on the rollercoaster train.

Figure 1



(a) Give **two** ways that the force arrows show that forces are vector quantities.

1. _____
2. _____

(2)

(b) Which arrow shows the weight of the rollercoaster train?

Tick **one** box.

A	☐	B	☐	C	☐	D	☐
----------	--------------------------	----------	--------------------------	----------	--------------------------	----------	--------------------------

(1)

(c) Which arrow shows the normal contact force?


Tick **one** box.

A	☐	B	☐	C	☐	D	☐
----------	--------------------------	----------	--------------------------	----------	--------------------------	----------	--------------------------

(1)

Figure 2 shows the magnitude of the acceleration of the rollercoaster train during the ride.

Figure 2



(d) Why has a line graph been drawn instead of a bar chart?

Tick **one** box.

Acceleration is a control variable.

☐

Both variables are continuous.

☐

Line graphs are easier to read.

☐

Time is a categoric variable.

☐

(1)

(e) What conclusion can be made from **Figure 2** about the motion of the rollercoaster train between 10 and 15 seconds?

Tick **one** box.

It is moving at a constant velocity.

☐

Its velocity is decreasing.

☐

Its velocity is increasing.

☐

(1)

(f) What is the maximum acceleration of the rollercoaster train?

Use **Figure 2**.

Acceleration = _____ m/s²

(1)

(g) The maximum safe acceleration for most people is 5 times the acceleration due to gravity.

Acceleration due to gravity = 9.8 m/s^2

Explain whether the acceleration of this rollercoaster train is safe for most people

(3)

- (h) One of the passengers on the rollercoaster train has a mass of 58 kg

Calculate the maximum force experienced by the passenger during the ride.

Use the equation:

$$\text{force} = \text{mass} \times \text{acceleration}$$

Give the unit.

Maximum force = _____ Unit _____

(3)


(Total 13 marks)

Q5.

Figure 1 shows a skydiver training in an indoor wind tunnel.


Large fans below the skydiver blow air upwards.

Figure 1



- (a) The skydiver is in a stationary position.

Complete the free body diagram for the skydiver.



(2)

- (b) The skydiver now straightens his legs to increase his surface area.

This causes the skydiver to accelerate upwards.

Explain why straightening his legs cause the skydiver to accelerate upwards.

(2)

- (c) A small aeroplane used for skydiving moves along a runway.

The aeroplane accelerates at 2 m / s^2 from a velocity of 8 m / s .

After a distance of 209 m it reaches its take-off velocity.

Calculate the take-off velocity of the aeroplane.

Take-off velocity = _____ m / s

(3)


(d) A skydiver jumps from an aeroplane.

There is a resultant vertical force of 300 N on the skydiver.

There is a horizontal force from the wind of 60 N.

Draw a vector diagram on **Figure 2** to determine the magnitude and direction of the resultant force on the skydiver.

Figure 2



Magnitude of resultant force = _____ N

(5)

Q6.

This question is about forces, quantities and vectors.

- (a) Write down the equation that links gravitational field strength, mass and weight.

_____ (1)

- (b) A small ball weighs 1.4 N.

gravitational field strength, $g = 9.8 \text{ N / kg}$

Calculate the mass of the ball.

Mass = _____ kg (3)

- (c) A white ball with mass 143 g is moving at a velocity of 7.9 m / s.

It collides with a red ball with mass of 150 g.

The red ball is stationary before the collision. The white ball stops after the collision.

Calculate the velocity of the red ball after the collision.

Give your answer to two significant figures.

Velocity of red ball = _____ m / s (4)

- (d) The white ball is thrown high into the air.

After it is released the ball moves up and then back down in a vertical line.

The free body force diagram in the figure below shows the forces on the ball at one point in its flight.

The force arrows are drawn to scale.



Explain what is happening to the ball at this point in its flight.

(4)

(Total 12 marks)

Mark schemes

Q1.

- | | | | |
|-----|---------------------------|---|------------|
| (a) | distance | | 1 |
| (b) | BC | | 1 |
| (c) | DE | | 1 |
| (d) | 5.0 (s) | <i>allow 5 (s)</i> | 1 |
| (e) | 6.0 (m/s) | <i>allow 6 (m/s)</i> | 1 |
| (f) | 6.0 / 5.0 | <i>allow ecf from parts (d) and (e)</i> | 1 |
| | = 1.2 (m/s ²) | <i>an answer of 1.2 scores 2 marks</i> | 1 |
| (g) | about four times faster | | 1 |
| | | | [8] |

Q2.

- | | | | |
|-----|--------------------------|--|---|
| (a) | 20 (degrees) | <i>allow answers in the range 19–21 (degrees)</i> | 1 |
| (b) | vector | | 1 |
| | scalar | | 1 |
| | | <i>in this order only</i> | |
| (c) | weight | <i>allow gravity</i>
<i>do not accept gravitational potential energy</i> | |
| | or | | |
| | (normal) contact (force) | <i>allow (normal) reaction (force)</i>
<i>allow air resistance</i>
<i>allow drag</i> | |

(d) the resultant force on the block is zero

1

(e) any **two** from:

do not accept force

- the (same) block

allow named feature of block e.g. mass / weight

- speed (at which block is pulled)
- angle (of string)

2

(f)
$$\left(\frac{5.2 + 5.6 + 5.4}{3} \right) = 5.4 \text{ N}$$

1

(g) glass

allow ecf from part (f) if glass not given

1

[9]

Q3.

(a) scalars have magnitude only

allow size for magnitude

1

vectors have magnitude and direction

1

(b) any **three** from:

- arrow pointing vertically downwards originating in the block, labelled weight

allow gravity

- arrow pointing left along the surface labelled friction; part of the arrow must be between the block and the carpet

- arrow pointing to the left touching block labelled air resistance or drag

- arrow pointing vertically upwards from bottom surface of block, labelled (normal) contact force

allow reaction (force) for contact (force)

judge horizontal and vertical by eye

*allow 1 mark for three correctly labelled arrows,
in correct directions but in incorrect positions*

3

(c) clear attempt to draw horizontal **and** vertical components

*horizontal line should extend beyond dashed line
on diagram*

1

horizontal component: 2.6 (N)
allow a range from 2.5 to 2.7

1

vertical component: 1.7 (N)
allow a range from 1.6 to 1.8

1

*if 2nd and 3rd marking points not awarded, allow
1 mark for clear measurements of **both** 5.2 ± 0.2
 cm and 3.4 ± 0.2 cm*

(d)
$$\frac{5.2 + X + 5.3}{3} = 5.4$$

1

(X =) 5.7 (N)

1

*allow 5.6 (N) or 5.8 (N) for **2** marks
 an answer of 5.7 (N) scores **2** marks*

- (e) any **three** from:
- angle (of string)
 - speed (at which block is pulled)
allow velocity (at which block is pulled)
 - area of block in contact with surface
 - mass / weight of block
ignore same block

3

[13]

Q4.

- (a) the arrows have different lengths
allow forces can have different sizes

1

the arrows point in different directions
allow forces can have different directions

1

- (b) **D**

1

- (c) **B**

1

- (d) both variables are continuous

1

- (e) it is moving at a constant velocity

1

- (f) $24.5 \text{ (m/s}^2\text{)}$

1

- (g) $5 \text{ g} = 49 \text{ (m/s}^2\text{)}$

$$49 \text{ m/s}^2 > 24.5 \text{ m/s}^2$$

so the ride is safe

allow ecf from (f) (ie if their answer to (f) was greater than 49, then the ride is unsafe)

(h) force = 58×24.5

allow ecf from (f)

$$\text{force} = 1421$$

Newtons

allow N

[13]

Q5.

(a) arrow of equal size pointing vertically downwards

judged by eye

labelled 'weight'

(b) the upwards force is greater than the downwards force

because air resistance increases

(c) $v^2 = (2 \times 2 \times 209) + 8^2$

$$v = \sqrt{900}$$

$$v = 30 \text{ (m / s)}$$

allow 30 (m / s) without working shown for 3 calculation marks

(d) vertical force (300 N) drawn with a suitable scale

horizontal force (60 N) drawn to the same scale

resultant force drawn in correct direction

value of resultant in the range 304 N – 308 N

Q6.

(a) weight = mass × gravitational field strength

1

(b) mass = weight ÷ g

1

$$= 1.4 \div 9.8$$

1

$$= 0.143 \text{ (kg)}$$

allow 0.143 with no working shown for 3 marks

1

(c) momentum = mass × velocity

momentum before = momentum after

1

$$143 \times 7.9 = 150 \times v$$

1

$$v = \frac{143 \times 7.9}{150}$$

1

$$= 7.5 \text{ (m / s)}$$

allow 7.5 (m / s) with no working shown for 4 marks

1

incorrect number of sig. figs max. 3 marks

(d) ball is falling / moving down

1

at terminal velocity

1

air resistance and weight have the same magnitude / size

1

so no acceleration / constant speed

1

[12]

Examiner reports

Q1.

- (a) Nearly 34% scored this mark. The most common incorrect response was velocity.
- (b) Almost 74% of students were able to identify the part of the graph that showed the woman moving at a constant velocity.
- (c) 68% of students selected the correct response for the part of the graph where the woman was stationary. The most common incorrect response was BC.
- (d) The most common response was to misread the scale with incorrect values of 2.5 or 4 often given. Nearly 45% gave the correct answer. A significant minority tried to use various equations of motion, producing incorrect answers.
- (e) This required a more straightforward graph reading and almost 72% gave the correct answer.
- (f) 66% gained two marks, with students either using the correct values read from the graph or correctly calculating using their previously incorrect values. A small number multiplied instead of dividing, despite the equation being shown.
- (g) Almost 53% correctly stated that typical cycling speed would be about four times faster than the typical walking speed. The most common wrong answer was about twice as fast.

Q2.

- (a) Around two thirds of students were able to measure the angle (20°) correctly.

A large proportion of the incorrect answers were either 70° ($90 - 20$) or 160° ($180 - 20$), indicating that students were measuring the correct angle but were not too familiar with reading a protractor. Around 3% of students did not attempt this question.
- (b) Less than one third of students achieved both marks for this question. A fair number realised that vector and scalar went together but got the words in the wrong order.

All of the given words were used in a variety of combinations.
- (c) Around 12% of students did not attempt this question. Of those who did, the most popular correct answers given were gravity and weight.

Incorrect answers included quantities which were not forces but were linked to the investigation in some way, such as kinetic energy, movement and mass. Also seen were words which appeared to be totally unrelated to the situation, such as electricity, light and fractional distillation.
- (d) Twice as many students selected the incorrect response of 'force is directly proportional to velocity' than chose the correct response of 'the resultant force on the block is zero'. The other two incorrect options had about the same number of responses as the correct one.
- (e) Students who understood the term 'control variable' were usually able to give at least one correct answer. Around 9% of students gained both marks, with a further

40% scoring one mark.

A large number of students thought that the nature of the surface and the force would be control variables, showing some confusion with independent and dependent variables. A small number of students gave random answers such as 'oxygen' and 'valid'.

- (f) Around 70% of students gained the mark for the correct calculation of the mean. The most frequent incorrect response was 16.2 N, the sum of the three numbers.
- (g) Around 85% of students were able to identify glass as the surface with the lowest frictional force. The most common incorrect answer was 'cardboard'. Nearly 10% did not answer this question.

Q3.

- (a) Around one quarter knew the correct definitions of vector and scalar quantities, thereby scoring two marks.

Some students knew that vectors had both magnitude and direction, and realised that scalars don't, but opted for them having direction rather than magnitude.

A number of students defined a scalar as speed and a vector as velocity, rather than these being examples of the different types of quantity.

- (b) Fewer than 2% of students scored two or three marks, with a further 15% picking up one mark.

Many did not appear to be familiar with representing forces on diagrams, so did not realise that they were expected to draw labelled arrows to show the forces. A large number did not know what forces were acting in this situation, many indicating different types of energy, motion, speed or momentum.

Of those who were aware of forces, the most common ones shown were weight / gravity, air resistance and friction. Many did not realise that friction and air resistance act in the opposite direction to motion. In addition, a few seemed to think that gravity acts upwards or horizontally.

The main error amongst those who knew the forces and their directions, was to show them acting in random positions around the block rather than on the block itself; in such cases one mark was awarded.

- (c) It appeared that many students did not know what components of a force are.

Of those who made some attempt, very few showed the components on the diagram. Some lines on the diagram were measured, usually the dotted horizontal line and the line representing the tension force, and the forces were calculated using the scaling factor.

Around 9 % gained all three marks.

- (d) More than three quarters of students were able to calculate the missing value correctly, thereby gaining the two marks for this question. The method used by some students was to multiply the mean force by 3, then subtract the other two values from this figure. An alternative method, done by many more students, was a trial-and-error approach, substituting various values in turn as the missing 'X' and working out the mean, until they found a value of 5.4 N.

Some students who did not know how to do this calculation ended up with very odd values, indicating that they either did not understand the concept of a mean value or they had not thought about whether their answer was sensible.

- (e) Approximately 45 % of students achieved 1 mark, with a further 13 % gaining 2 marks, but very few gained all three marks. The most common answer was the weight of the block, with few students opting for the angle between the block and the string.

When asked for control variables, answers which include the same equipment or the same person doing the experiment are unlikely to gain credit. A large number of students included the independent variable, i.e. the type of surface, and / or the dependent variable, i.e. the force, as control variables.