Rewarding Learning

ADVANCED SUBSIDIARY (AS)

General Certificate of Education

2022

Further Mathematics

Assessment Unit AS 1

assessing

Pure Mathematics

SFM11

[SFM11]

MONDAY 16 MAY, AFTERNOON

TIME

1 hour 30 minutes.

INSTRUCTIONS TO CANDIDATES

Write your Centre Number and Candidate Number on the Answer Booklet provided.
Answer **all eight** questions.
Show clearly the full development of your answers.
Answers should be given to three significant figures unless otherwise stated.
You are permitted to use a graphic or scientific calculator in this paper.

INFORMATION FOR CANDIDATES

The total mark for this paper is 100.
Figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question.
A copy of the **Mathematical Formulae and Tables booklet** is provided.
Throughout the paper the logarithmic notation used is $\ln z$ where it is noted that $\ln z \equiv \log_e z$

Answer all eight questions.

Show clearly the full development of your answers.

Answers should be given to three significant figures unless otherwise stated.

1 Matrices **A** and **B** are given by

$$\mathbf{A} = \begin{pmatrix} 4 & 1 & 0 \\ 3 & -2 & p \end{pmatrix} \quad \text{and} \quad \mathbf{B} = \begin{pmatrix} 1 & 0 \\ -2 & 2 \\ 1 & p+1 \end{pmatrix}$$

(i) Calculate the matrix product **AB**, leaving your answer in terms of p. [2]

(ii) Given that **AB** is singular, find the possible values of p. [4]

2 The complex numbers z_1 and z_2 are given by

$$z_1 = 2 + 3\mathrm{i} \quad \text{and} \quad z_2 = 1 + q\mathrm{i}$$

where q is a real number.

(i) Find the modulus and argument of z_1 [3]

(ii) Given that $z_1 + 3z_2$ is a real number, find the value of q. [2]

(iii) Hence simplify

$$\frac{z_1}{\overline{z_2}}$$

giving your answer in the form $a + b\mathrm{i}$, where a and b are real numbers. [4]

3 The roots of the quadratic equation

$$2x^2 - 4x + 5 = 0$$

are α and β.

(i) Write down the values of $(\alpha + \beta)$ and $\alpha\beta$. [3]

(ii) By considering $(\alpha + \beta)^3$, find the value of $\alpha^3 + \beta^3$ [4]

(iii) Find a quadratic equation with integer coefficients whose roots are

$$\frac{1}{\alpha + 1} \quad \text{and} \quad \frac{1}{\beta + 1}$$ [6]

4 **(i)** The matrix $\mathbf{S} = \begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix}$

Describe fully the transformation represented by **S** [2]

(ii) The set of points which form the curve

$$x^2 + 10y^2 + 6xy - 25 = 0$$

is transformed by the matrix **S**

Show that the curve formed by the image points is a circle and state its centre and radius. [6]

(iii) The matrix $\mathbf{N} = \begin{pmatrix} 3 & 2 \\ 1 & 4 \end{pmatrix}$

Find the single matrix that represents the combined effect of the transformation represented by **S** followed by that represented by **N** [2]

5 Two lines are given by the equations

$$\frac{x+2}{4} = \frac{y-1}{2} = \frac{z+4}{3} \quad \text{and} \quad x-3 = \frac{y-2}{-1} = \frac{z+3}{-2}$$

(i) Show that the two lines intersect and find their point of intersection. [9]

(ii) Find the Cartesian equation of the plane which contains both these lines. [6]

(iii) Find the coordinates of the point at which this plane cuts the y-axis. [1]

6 A matrix **M** is given by

$$\mathbf{M} = \begin{pmatrix} 2 & 1 & p \\ 1 & -2 & -2 \\ 0 & p+1 & 2 \end{pmatrix}$$

(i) Show that the determinant of **M** is given by

$$p^2 + 5p - 6$$ [3]

A system of linear equations is given by

$$\begin{aligned} 2x + y + pz &= 3 \\ x - 2y - 2z &= 6 \\ (p+1)y + 2z &= -7 \end{aligned}$$

(ii) Find the possible values of p for which this system has a unique solution. [3]

(iii) If $p = 2$ find, without the use of a calculator, the inverse of **M** [5]

(iv) Hence for $p = 2$, find the unique solution of the system of equations. [4]

7 **(a)** **(i)** Find the real values of a and b such that

$$(a + b\mathrm{i})^2 = 12 - 16\mathrm{i}$$

[6]

(ii) Hence write down the solutions to the equation

$$z^2 = 12 - 16\mathrm{i}$$ [2]

(b) Sketch, on a carefully labelled Argand diagram, the region consisting of points which represent a complex number, z, satisfying both

$$|z - 2\mathrm{i}| \leqslant 4 \quad \text{and} \quad |z + 4 - \mathrm{i}| \leqslant |z + \mathrm{i}|$$ [7]

8 ABCDE is a rectangular-based pyramid with apex E, as shown in **Fig. 1** below.




Fig. 1

The vertices A, B, D and E have coordinates (−2, −2, 1), (1, 2, 6), (−3, 0, 0) and (2, −2, 2) respectively.

(i) Find the coordinates of vertex C. [3]

(ii) Find the volume of the pyramid. [7]

The line l has the vector equation

$$\mathbf{r} = \begin{pmatrix} -4 \\ 2 \\ -1 \end{pmatrix} + \lambda \begin{pmatrix} 3 \\ -1 \\ 2 \end{pmatrix}$$

(iii) Find the shortest distance between the point E and the line l. [6]

THIS IS THE END OF THE QUESTION PAPER

Permission to reproduce all copyright material has been applied for.
In some cases, efforts to contact copyright holders may have been unsuccessful and CCEA
will be happy to rectify any omissions of acknowledgement in future if notified.