Rewarding Learning

ADVANCED SUBSIDIARY (AS)
General Certificate of Education
2022

Further Mathematics *StemUp Model Solution*

SFM21

Assessment Unit AS 2
assessing
Applied Mathematics

[SFM21]
THURSDAY 26 MAY, AFTERNOON

TIME

1 hour 30 minutes.

INSTRUCTIONS TO CANDIDATES

Write your Centre Number and Candidate Number on the Answer Booklet provided and on the Section D Supplementary Answer Booklet (if applicable).
You must answer **all** questions from sections A and B **or** A and C **or** A and D **or** C and D.
You should spend equal time on each of the two sections.
Candidates taking Section D should use the insert provided and attach to your Answer Book using the treasury tag provided.
Show clearly the full development of your answers.
Answers should be given to three significant figures unless otherwise stated.
You are permitted to use a graphic or scientific calculator in this paper.

INFORMATION FOR CANDIDATES

The total mark for this paper is 100.
The total mark for each section of this paper is 50.
Figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question.
Answers should include diagrams where appropriate and marks may be awarded for them.
Take $g = 9.8\,\text{m}\,\text{s}^{-2}$, unless specified otherwise.
A copy of the **Mathematical Formulae and Tables booklet** is provided.
Throughout the paper the logarithmic notation used is $\ln z$ where it is noted that $\ln z \equiv \log_e z$

stemup.co.uk

BLANK PAGE

stemup.co.uk

SECTION A Mechanics 1

Answer all five questions in this section.

1 A bead B of mass 0.5 kg is attached to one end of a light inextensible string of length L metres.

The other end of the string is attached to a fixed point O.

The bead is made to describe horizontal circles with constant angular velocity 3 rad s^{-1} with centre vertically below O as shown in **Fig. 1** below.




Fig. 1

The string makes an angle θ with the downward vertical.

The tension in the string is 12 N.

(i) Draw a diagram showing all the external forces acting on the bead. [2]

(ii) Find the values of θ and L. [7]

stemup.co.uk

2 A car of mass 1200 kg is travelling along a straight horizontal road.

The car experiences a constant resistance to motion of 300 N.

At a certain instant, the car is accelerating at $0.4\,\text{m s}^{-2}$ and the engine of the car is working at a rate of 19.5 kW.

(i) Find the speed of the car at this instant. [4]

The car then moves up a hill inclined at 30° to the horizontal at a constant speed of $15\,\text{m s}^{-1}$

The resistance to motion remains 300 N.

(ii) Find the power of the car's engine as it moves up the line of greatest slope of the hill. [4]

3 A particle of mass 6 kg moves x metres from a fixed point O, in a positive direction along the x-axis.

It is acted upon by a variable force, F newtons, given by

$$F = px + 1$$

where p is a positive constant.

The particle is at rest at O.

When $x = 3$ the work done by F is $3p^2$ joules.

Find the speed of the particle at this point. [7]

stemup.co.uk

4 A climber of mass 80 kg is securely attached by a light elastic rope to a fixed point O on a vertical rock face.

The rope has natural length 20 metres and modulus of elasticity 18 500 newtons.

He climbs to a point which is d metres vertically above O where $d < 20$

He slips and falls from rest, but is brought to instantaneous rest by the force in the rope when the rope has extended by 8 metres.

(i) By taking the Gravitational Potential Energy to equal zero at O, use the Principle of Conservation of Mechanical Energy to find the value of d. [8]

(ii) Find the speed of the climber when he is 25 metres below O. [6]

(iii) Explain briefly why, in reality, his speed at this point is less than the value found. [1]

stemup.co.uk

5 A light elastic string is attached to a fixed point A on a plane inclined at an angle θ to the horizontal where $\sin\theta = \frac{3}{5}$

A particle of mass 10 kg is attached to the other end of the string.

The string has natural length 5 metres and modulus of elasticity 147 newtons.

The particle is first held at a point B on the plane, where AB = 5 metres and B is below A as shown in **Fig. 2** below.




Fig. 2

The string is parallel to a line of greatest slope of the plane.

The particle is released from rest.

(i) Given that the plane is smooth, find the distance down the slope from B that the particle moves before next coming to instantaneous rest. [5]

If instead the plane is rough, the particle next comes to instantaneous rest 2 metres down the slope from B.

(ii) Find the coefficient of friction between the particle and the plane. [6]

stemup.co.uk

SECTION B Mechanics 2

Answer all five questions in this section.

1 A violin has four strings, each of which has a pitch determined by its fundamental frequency.

The frequency f depends on the tension S in the string, the mass per unit length ρ of the string and the length l of the string.

It is known that the frequency is related to the other string properties as follows:

$$f = kS^x \rho^y l^z$$

where k is a dimensionless constant.

The dimensions of frequency are T^{-1}

(i) Use the method of dimensions to find x, y and z. [6]

The violinist holds one of the strings which effectively makes its length 15% shorter.

(ii) Find the percentage change in its frequency, given that the tension and mass per unit length are unchanged. [2]

2 A satellite of mass m kg is moving in a circular orbit of radius r metres around a planet of mass M kg.

(i) Show that the angular speed of the satellite around the planet can be expressed as

$$\sqrt{\frac{GM}{r^3}} \text{ rad s}^{-1}$$

where G is the universal gravitational constant. [6]

A satellite orbits Saturn in a circular path of radius 133 600 km.

The mass of Saturn is 5.683×10^{26} kg and the value of G is $6.674 \times 10^{-11}\,\text{m}^3\,\text{s}^{-2}\,\text{kg}^{-1}$

(ii) Find the period of orbit of the satellite in hours. [4]

stemup.co.uk

3 A pendulum consists of a bob P of mass 4 kg, attached to one end of a light rod of length 4 m.

The other end of the rod is freely hinged at the fixed point O.

The bob is at rest at its lowest point and is then given a horizontal impulse so that it starts to move with speed $u\,\mathrm{m\,s^{-1}}$

Find the possible values of u if the pendulum is to perform complete circles. [8]

4 A teacher on lunchtime duty notices Pauline standing 5 m North East of Roisin.

At the same instant, each girl starts to run.

Pauline has a velocity of $4\,\mathrm{m\,s^{-1}}$ due South.

Roisin has a velocity of $3\sqrt{2}\,\mathrm{m\,s^{-1}}$ South East.

If they maintain these velocities, find the shortest distance between Pauline and Roisin in the subsequent motion. [9]

stemup.co.uk

5 Two small identical rings, A and B, each of mass 0.6 kg are threaded on to a fixed rough horizontal wire.

The rings are attached to opposite ends of a light inextensible string of length $2L$ metres, as shown in **Fig. 1** below.




Fig. 1

A particle P of mass 0.2 kg is attached to the midpoint of the string.

When the rings are in limiting equilibrium P hangs vertically below the wire, with AB = 0.4 m.

The coefficient of friction between each ring and the wire is 0.5

(i) Draw a diagram showing all the forces acting on each of the three masses. [3]

(ii) Show that the normal reaction between each ring and the wire is $0.7g$ N. [4]

(iii) Find the value of L. [8]

stemup.co.uk

SECTION C Statistics

Answer all five questions in this section.

1 The owners of a football stadium would like to find out the views of the supporters who attend matches with regard to changing some of the stadium seating areas to standing sections.

By looking at previous attendance statistics, the owners have identified that the stadium crowd typically comprises 80% home supporters and 20% away supporters.

They intend to place a number of interviewers around the stadium at the next game.

Each interviewer has been instructed to interview 10 supporters, of which 8 must be home supporters and 2 must be away supporters.

(i) State the non-random sampling technique that is being used. [1]

(ii) Describe one advantage and one disadvantage of using this particular sampling technique. [4]

2 A researcher is investigating how the mass of a chimpanzee is related to its age.

She took a sample of 10 chimpanzees from a nature reserve with ages ranging from birth to 12 weeks old.

For each chimpanzee she recorded the age x, number of weeks since birth, and the mass y kg.

The largest mass recorded was 4 kg.

Some of the summary statistics are shown below.

$$\Sigma x = 60 \qquad \Sigma y = 24 \qquad \Sigma x^2 = 480 \qquad \Sigma y^2 = 62.4$$

From the data obtained the researcher calculated the product–moment correlation coefficient to be 0.9

(i) Find the equation of the least squares regression line of chimpanzee mass y on age x, in the form $y = a + bx$ where a and b are constants. [9]

(ii) Suggest a reason why it would not be appropriate to use the regression line calculated in part **(i)** to:

(a) estimate the mass of a chimpanzee at 14 weeks old; [1]

(b) estimate the age of a chimpanzee with a mass of 3 kg. [1]

stemup.co.uk

3 A fairground game involves the contestant rolling two fair tetrahedral dice at the same time.

Each tetrahedral die has four faces numbered 1, 2, 3 and 4

The positive difference between the scores shown on the two dice determines the prize won by the contestant.

The possible prizes that the contestant can win are given below.

- £5 if the difference is 3
- £1 if the difference is 1
- £0 if the difference is 0 or 2

Let the amount, in pounds, won by a contestant on a single attempt be modelled by the random variable X.

(i) Using a sample space, or otherwise, copy and complete **Table 1** below showing the probability distribution of X. [5]

Table 1

x	0	1	5
$P(X = x)$			

The game organiser would like to make an average profit of 50p per contestant attempt.

(ii) Find the price that the organiser should charge a contestant for a single attempt in order to achieve this average profit. [3]

(iii) Calculate Var(X). [3]

The random variable Y is related to X by the formula $Y = 3X - 4$

(iv) Find the mean and variance of Y. [3]

stemup.co.uk

4 An experiment involves a biased coin being tossed a number of times.

Let p be the probability that the coin lands on tails on any given throw.

The probability that the coin first lands on tails on the second throw is 0.24

(i) Find the two possible values of p. [4]

Using the smaller value of p, find the probability that:

(ii) the coin lands on tails for the first time on the eighth throw; [3]

(iii) the coin does not land on tails before the fifth throw. [2]

5 Each letter of the word DESTINY is written on a separate card.

The word DESTINY contains two vowels (E and I) and five consonants.

Four cards out of the seven are to be chosen at random.

(i) Find the number of different combinations possible. [1]

(ii) Find the probability that exactly three consonants are on the four chosen cards. [3]

The seven cards are to be arranged in a random order in a line.

(iii) Find the number of arrangements possible. [1]

(iv) Find the probability that the two cards containing vowels are next to each other in either order. [3]

Another word has n different letters, where $n \geqslant 3$

Two of the letters are vowels and the remainder are consonants.

Each letter is written on a separate card and the cards are arranged in a random order in a line.

(v) Find, in terms of n, the probability that the two cards containing the vowels are next to each other. [3]


SECTION D Discrete and Decision Mathematics

Answer all five questions in this section.

Note: There is a Supplementary Answer Booklet for questions **3(i)** and **(iv)**
Please attach this to your answer booklet using the treasury tag provided.

1 **(a)** The graph G_1 on 5 vertices is shown in **Fig. 1** below.

Draw a new version of G_1 to demonstrate that it is planar.



[2]

Fig. 1

(b) The graph G_2 on 6 vertices is shown in **Fig. 2** below.




Fig. 2

(i) How many edges must be removed from G_2 to leave a spanning tree? [1]

(ii) Draw a spanning tree that is a subgraph of G_2 [2]

(iii) How many edges must be added to G_2 to form K_6, the complete graph on 6 vertices? [2]

stemup.co.uk

(c) Write down an Eulerian trail of the graph $K_{3,2}$ shown in **Fig. 3** below. [2]




Fig. 3

2 The sequence u_n is defined by the relationship

$$u_{n+2} = \frac{5}{3}u_{n+1} + \frac{2}{3}u_n$$

where $u_0 = 2$ and $u_1 = \frac{5}{3}$ and n is an integer.

(i) Show that

$$u_n = \alpha^n + \beta^n$$

where α and β are real numbers to be found. [7]

(ii) Given that $u_t = 4096.000002$ to 10 significant figures, find the value of t. [3]

stemup.co.uk

3 The travel times in minutes between seven towns are given in the road network in **Fig. 4** below.




Fig. 4

A race is planned from Aysbury to Groomsport.

(i) Apply Dijkstra's algorithm to this network to calculate the shortest time from Aysbury to Groomsport.

Use the version of the road network **Diagram 1** given in the Section D Supplementary Answer Booklet.

Be careful to fill the algorithm details into the box at each vertex.

Do not delete any of the temporary times. [5]

(ii) Write down the fastest route from Aysbury to Groomsport and the corresponding minimum time. [2]

(iii) Write down the minimum time from Aysbury to Evensby. [1]

stemup.co.uk

Due to road works, the road from Drilly to Beacon is temporarily one way, as shown in **Fig. 5** below.




Fig. 5

(iv) Apply Dijkstra's algorithm to **Fig. 5** to recalculate the new shortest time from Aysbury to Groomsport, taking into account the one-way road.

Use the version of the road network **Diagram 2** given in the Section D Supplementary Answer Booklet.

Be careful to fill the algorithm details into the box at each vertex.

Do not delete any of the temporary times. [3]

stemup.co.uk

4 Let p, q and r be propositional statements.

The Boolean implication operator

$$p \Rightarrow q$$

is defined by the truth table in **Fig. 6** below.

p	q	$p \Rightarrow q$
T	T	T
T	F	F
F	T	T
F	F	T

Fig. 6

(i) Use truth tables to show that $p \Rightarrow q$ and $\sim q \Rightarrow \sim p$ are equivalent. [3]

(ii) Use truth tables to prove that

$$[(p \Rightarrow q) \text{ and } (q \Rightarrow r)] \Rightarrow (p \Rightarrow r)$$

is a tautology, i.e. is always true. [7]

stemup.co.uk

5 The set of six permutations p, q, r, s, t and u forms a group G under the operation $\circ$, the composition of permutations.

The permutations are defined by

$$p = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix} \qquad q = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \qquad r = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix}$$

$$s = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix} \qquad t = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix} \qquad u = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix}$$

An example composition is

$$q \circ s = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \circ \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix} = t$$

so $q \circ s = t$.

(i) Write down the identity permutation. [1]

A partially complete Cayley table for this group is given in **Fig. 7** below.

$\circ$	p	q	r	s	t	u
p		u	t	r	s	
q	u	p	s	t		
r	s		u			
s	t	r	q			
t						
u						

Fig. 7

(ii) Copy and complete the Cayley table in **Fig. 7** [4]

stemup.co.uk

(iii) Copy and complete the table of elements and their periods in **Fig. 8** below.

Element	p	q	r	s	t	u
Period						

[2]

Fig. 8

(iv) List all the subgroups of G. [3]

THIS IS THE END OF THE QUESTION PAPER


stemup.co.uk

Surname	StemUp Model Solution
Answer booklet	Write the question number in the left-hand margin at the start of each answer.

Question number

Section A Mechanics 1

A1 (i)



(ii) Resolve vertically (no vertical acceleration):

$12 \cos \theta = 0.5g = 4.9$

$\cos \theta = \frac{4.9}{12}$

$\theta = 65.9°$ (3 s.f.)

Radius of the circle $r = L \sin \theta$

Resolve horizontally (towards centre): $12 \sin \theta = mr\omega^2$

$12 \sin \theta = 0.5 \times L \sin \theta \times 3^2$

$12 = 4.5L$

$L = \frac{12}{4.5} = \frac{8}{3} = 2.67$ m (3 s.f.)

A2 (i) Driving force $F = \frac{P}{v} = \frac{19\,500}{v}$

Newton II along the road: $F - 300 = 1200 \times 0.4$

Question number

$\frac{19\,500}{v} - 300 = 480$

$\frac{19\,500}{v} = 780$

$v = 25\ m\ s^{-1}$

(ii) Constant speed, so the forces along the slope balance:

$F = 1200g \sin 30° + 300$

$F = 5880 + 300 = 6180\ N$

$P = Fv = 6180 \times 15 = 92\,700\ W$

Power = 92.7 kW

A3 Work done by $F = \int_0^3 (px + 1)\,dx = 3p^2$

$\left[\frac{px^2}{2} + x\right]_0^3 = 3p^2$

$\frac{9p}{2} + 3 = 3p^2$

$6p^2 - 9p - 6 = 0$

$2p^2 - 3p - 2 = 0$

$(2p + 1)(p - 2) = 0$

$p = 2$ or $p = -\frac{1}{2}$ but $p > 0$, so $p = 2$

Work done $= 3p^2 = 3 \times 2^2 = 12\ J$

Work done = change in kinetic energy (starts from rest):

$12 = \frac{1}{2} \times 6 \times v^2$

$v^2 = 4$

Question number

Speed = 2 m s^{-1}

A4 (i) Take GPE = 0 at O.

At the highest point (at rest, rope slack):

KE = 0, GPE = 80gd = 784d, EPE = 0

Total mechanical energy = 784d J

At the lowest point the rope is 20 + 8 = 28 m long,

so the climber is 28 m below O and at rest:

KE = 0, GPE = 80g × (−28) = −21 952 J

$EPE = \frac{\lambda x^2}{2l} = \frac{18\,500 \times 8^2}{2 \times 20} = 29\,600$ J

Total mechanical energy = 29 600 − 21 952 = 7648 J

Conservation of mechanical energy:

784d = 7648

d = 9.76 m (3 s.f.)

(ii) 25 m below O the extension is 25 − 20 = 5 m.

$KE = \frac{1}{2} \times 80v^2 = 40v^2$

GPE = 80g × (−25) = −19 600 J

$EPE = \frac{18\,500 \times 5^2}{2 \times 20} = 11\,562.5$ J

Total = $40v^2$ − 19 600 + 11 562.5 = $40v^2$ − 8037.5

Conservation of mechanical energy:

$40v^2 - 8037.5 = 7648$

$40v^2 = 15\,685.5$

$v^2 = 392.1375$

v = 19.8 m s^{-1} (3 s.f.)

Question number

(iii) Air resistance has been ignored in the model; in reality it does work against the climber, so his speed is less.

AS (i) Let the particle move d m down the slope from B to C.

At B the string is at its natural length (AB = 5 m).

$\sin\theta = \frac{3}{5}$, take GPE = 0 at C.

Energy at B = Energy at C:

$0 + 10g \times d\sin\theta + 0 = 0 + 0 + \frac{147d^2}{2 \times 5}$

$10 \times 9.8 \times \frac{3}{5}d = 14.7d^2$

$58.8d = 14.7d^2$

$d \neq 0$, so $58.8 = 14.7d$

$d = 4$ m

(ii) $\cos\theta = \frac{4}{5}$

$R = 10g\cos\theta = 10g \times \frac{4}{5} = 8g$ N

Friction (limiting, particle sliding) $= \mu R = 8\mu g$ N

Now d = 2 m. Work-energy from B to C:

Loss in GPE = gain in EPE + work done against friction

$10g \times 2 \times \frac{3}{5} = \frac{147 \times 2^2}{2 \times 5} + 8\mu g \times 2$

$117.6 = 58.8 + 156.8\mu$

$156.8\mu = 58.8$

Question number

$\mu = \frac{3}{8} = 0.375$

Section B Mechanics 2

B1 (i)

$[f] = T^{-1}$ $[S] = MLT^{-2}$

$[\rho] = ML^{-1}$ $[l] = L$

$T^{-1} = (MLT^{-2})^x (ML^{-1})^y L^z$

Compare powers:

M: $0 = x + y$

L: $0 = x - y + z$

T: $-1 = -2x$

$x = \frac{1}{2}$

$y = -x = -\frac{1}{2}$

$z = y - x = -\frac{1}{2} - \frac{1}{2} = -1$

$x = \frac{1}{2}$, $y = -\frac{1}{2}$, $z = -1$

(ii)

$f = kS^{\frac{1}{2}} \rho^{-\frac{1}{2}} l^{-1}$, so with S and ρ fixed $f \propto \frac{1}{l}$

New length = 0.85l

$f_1 = \frac{f_0}{0.85} = 1.176 f_0$

Frequency increases by 17.6% (3 s.f.)

Question number

B2 (i) Gravitational force provides the centripetal force:

$\frac{GMm}{r^2} = mr\omega^2$

$GMm = mr^3\omega^2$

$\omega^2 = \frac{GM}{r^3}$

$\omega = \sqrt{\frac{GM}{r^3}}$ rad s^{-1} as required

(ii) $r = 133\,600$ km $= 1.336 \times 10^8$ m


$\omega = \sqrt{\frac{6.674 \times 10^{-11} \times 5.683 \times 10^{26}}{(1.336 \times 10^8)^3}}$

$\omega = 1.2612 \times 10^{-4}$ rad s^{-1}

$T = \frac{2\pi}{\omega} = \frac{2\pi}{1.2612 \times 10^{-4}} = 49\,820$ s

Period $= \frac{49\,820}{3600} = 13.8$ hours (3 s.f.)

B3



Take GPE = 0 at the lowest point A. B is the top, 8 m above A.

At A: KE $= \frac{1}{2} \times 4 \times u^2 = 2u^2$, GPE = 0

Question number

At B: $KE = \frac{1}{2} \times 4 \times v^2 = 2v^2$, $GPE = 4g \times 8 = 32g$

Conservation of energy: $2u^2 = 2v^2 + 32g$

The bob is on a rod, so it completes circles provided

it still has speed at the top: $v > 0$, i.e. $v^2 > 0$

$2u^2 - 32g > 0$

$u^2 > 16g = 156.8$

$u > 12.5 \text{ m s}^{-1}$ (3 s.f.)

B4

Let $\mathbf{i}$ be a unit vector due East and $\mathbf{j}$ a unit vector due North.

$\mathbf{v}_P = -4\mathbf{j}$

$\mathbf{v}_R = 3\sqrt{2}\cos 45° \mathbf{i} - 3\sqrt{2}\sin 45° \mathbf{j} = 3\mathbf{i} - 3\mathbf{j}$

Velocity of P relative to R:

${}_P\mathbf{v}_R = \mathbf{v}_P - \mathbf{v}_R = -4\mathbf{j} - (3\mathbf{i} - 3\mathbf{j}) = -3\mathbf{i} - \mathbf{j}$

Initial position of P relative to R (5 m North East):

${}_P\mathbf{r}_R = \frac{5}{\sqrt{2}}\mathbf{i} + \frac{5}{\sqrt{2}}\mathbf{j}$

After t seconds: ${}_P\mathbf{r}_R = (\frac{5}{\sqrt{2}} - 3t)\mathbf{i} + (\frac{5}{\sqrt{2}} - t)\mathbf{j}$

$D^2 = (\frac{5}{\sqrt{2}} - 3t)^2 + (\frac{5}{\sqrt{2}} - t)^2$

$\frac{d(D^2)}{dt} = -6(\frac{5}{\sqrt{2}} - 3t) - 2(\frac{5}{\sqrt{2}} - t) = 0$

$20t = \frac{40}{\sqrt{2}} \Rightarrow t = \sqrt{2}$ s


${}_P\mathbf{r}_R = (\frac{5}{\sqrt{2}} - 3\sqrt{2})\mathbf{i} + (\frac{5}{\sqrt{2}} - \sqrt{2})\mathbf{j} = -\frac{1}{\sqrt{2}}\mathbf{i} + \frac{3}{\sqrt{2}}\mathbf{j}$

Question number

$D^2 = \frac{1}{2} + \frac{9}{2} = 5$

Shortest distance $= \sqrt{5} = 2.24$ m (3 s.f.)

B5 (i)



θ is the angle each string makes with the vertical.

Friction 0.5R on each ring acts outwards (rings about to slip inwards).

(ii) Resolve vertically for P: $2T \cos\theta = 0.2g$

$T \cos\theta = 0.1g$

Resolve vertically for ring A: $R = 0.6g + T\cos\theta$

$R = 0.6g + 0.1g$

$R = 0.7g$ N as required

(iii) Limiting equilibrium, so friction $F = \mu R = 0.5 \times 0.7g = 0.35g$

Resolve horizontally for ring A: $T \sin\theta = 0.35g$

From (ii): $T \cos\theta = 0.1g$

Divide: $\tan\theta = \frac{0.35g}{0.1g} = 3.5$

$\theta = 74.05°$

P is below the midpoint of AB, so the horizontal

distance from A to P is 0.2 m:

Question number

$L \sin\theta = 0.2$

$L = \dfrac{0.2}{\sin 74.05°}$

$L = 0.208$ m (3 s.f.)

Section C Statistics

C1 (i) Quota sampling

(ii) Advantage: it is quick and cheap, and no sampling frame is needed; the interviewer simply fills the quota, and the sample still reflects the 80% : 20% split of supporters.

Disadvantage: the interviewer chooses who to interview, which introduces interviewer bias, so the sample is not random (and non-responses are simply replaced).

C2 (i) $S_{xx} = \Sigma x^2 - \dfrac{(\Sigma x)^2}{n} = 480 - \dfrac{60^2}{10} = 120$

$S_{yy} = \Sigma y^2 - \dfrac{(\Sigma y)^2}{n} = 62.4 - \dfrac{24^2}{10} = 4.8$

$r = \dfrac{S_{xy}}{\sqrt{S_{xx}S_{yy}}}$

$0.9 = \dfrac{S_{xy}}{\sqrt{120 \times 4.8}} = \dfrac{S_{xy}}{24}$

$S_{xy} = 21.6$

$\bar{x} = \dfrac{60}{10} = 6, \quad \bar{y} = \dfrac{24}{10} = 2.4$

Question number

$b = \frac{S_{xy}}{S_{xx}} = \frac{21.6}{120} = 0.18$

$a = \bar{y} - b\bar{x} = 2.4 - 0.18 \times 6 = 1.32$

$y = 1.32 + 0.18x$

(ii)(a) 14 weeks is outside the range of the data (0 to 12 weeks), so this would be extrapolation, which is unreliable.

(b) This is the regression line of y on x, so it should only be used to estimate y from x. To estimate age from mass the regression line of x on y would be needed.

C3 (i) Sample space of positive differences:

	1	2	3	4
1	0	1	2	3
2	1	0	1	2
3	2	1	0	1
4	3	2	1	0

$P(\text{diff } 0) = \frac{4}{16}$, $P(\text{diff } 1) = \frac{6}{16}$, $P(\text{diff } 2) = \frac{4}{16}$, $P(\text{diff } 3) = \frac{2}{16}$

$P(X = 0) = P(\text{diff } 0 \text{ or } 2) = \frac{8}{16} = \frac{1}{2}$

$P(X = 1) = \frac{6}{16} = \frac{3}{8}$, $P(X = 5) = \frac{2}{16} = \frac{1}{8}$

x	0	1	5
$P(X = x)$	$\frac{1}{2}$	$\frac{3}{8}$	$\frac{1}{8}$

Question number

(ii) $E(X) = 0 \times \frac{1}{2} + 1 \times \frac{3}{8} + 5 \times \frac{1}{8}$

$E(X) = \frac{3}{8} + \frac{5}{8} = £1$

Charge = 1 + 0.50 = £1.50

(iii) $E(X^2) = 0^2 \times \frac{1}{2} + 1^2 \times \frac{3}{8} + 5^2 \times \frac{1}{8} = \frac{28}{8} = 3.5$

$Var(X) = E(X^2) - [E(X)]^2 = 3.5 - 1^2 = 2.5$

(iv) $E(Y) = 3E(X) - 4 = 3 \times 1 - 4 = -1$

$Var(Y) = 3^2 \times Var(X) = 9 \times 2.5 = 22.5$

C4 (i) X = number of throws up to and including the first tail

$X \sim Geo(p)$

$P(X = 2) = (1 - p)p = 0.24$

$p - p^2 = 0.24$

$p^2 - p + 0.24 = 0$

$(p - 0.4)(p - 0.6) = 0$

$p = 0.4$ or $p = 0.6$

(ii) $p = 0.4$, $q = 0.6$

$P(X = 8) = q^7 p = 0.6^7 \times 0.4$

$= 0.0112$ (3 s.f.)

(iii) $P(X \geq 5) = P(\text{no tail in the first four throws}) = q^4$

$= 0.6^4 = 0.1296$

$= 0.130$ (3 s.f.)

Question number

C5 (i) ${}^7C_4 = 35$

(ii) 3 consonants from 5 and 1 vowel from 2:

${}^5C_3 \times {}^2C_1 = 10 \times 2 = 20$

$P = \frac{20}{35} = \frac{4}{7} = 0.571$ (3 s.f.)

(iii) $7! = 5040$

(iv) Treat the two vowels as one block: 6! arrangements,

and the vowels can be in 2 orders within the block.

$2 \times 6! = 1440$

$P = \frac{1440}{5040} = \frac{2}{7} = 0.286$ (3 s.f.)

(v) Total arrangements = $n!$

Vowels together: $2 \times (n-1)!$


$P = \frac{2 \times (n-1)!}{n!} = \frac{2 \times (n-1)!}{n \times (n-1)!}$

$P = \frac{2}{n}$

Section D Discrete and Decision Mathematics

Question number

D1 (a)




No edges cross, so G_1 is planar.

(b)(i) G_2 has 12 edges; a spanning tree on 6 vertices has 5 edges.

Edges to remove = 12 − 5 = 7

(ii)



(5 edges of G_2, connected, no cycles, all 6 vertices)

(iii) Number of edges in $K_6 = \frac{1}{2} \times 6 \times 5 = 15$

Edges to add = 15 − 12 = 3

(c) d and e have odd degree (3), so the trail starts at d, ends at e:

d a e b d c e

D2 (i) Try $u_n = kx^n$:

Question number

$kx^{n+2} = \frac{5}{3}kx^{n+1} + \frac{2}{3}kx^n$

Multiply by 3: $3kx^{n+2} = 5kx^{n+1} + 2kx^n$

$kx^n(3x^2 - 5x - 2) = 0$

$x \neq 0$, so $3x^2 - 5x - 2 = 0$

$(3x + 1)(x - 2) = 0$

$x = 2$ or $x = -\frac{1}{3}$

General solution: $u_n = A \times 2^n + B(-\frac{1}{3})^n$

$n = 0$: $2 = A + B$

$n = 1$: $\frac{5}{3} = 2A - \frac{1}{3}B \Rightarrow 5 = 6A - B$

Add: $7 = 7A \Rightarrow A = 1, \ B = 1$

$u_n = 2^n + (-\frac{1}{3})^n$

So $u_n = \alpha^n + \beta^n$ with $\alpha = 2, \ \beta = -\frac{1}{3}$

(ii) As n increases, $(-\frac{1}{3})^n \to 0$, so $u_t \approx 2^t$


$2^t \approx 4096 = 2^{12} \Rightarrow t = 12$

Check: $u_{12} = 4096 + (-\frac{1}{3})^{12} = 4096.0000019 = 4096.000002$ (10 s.f.)

$t = 12$

D3 (i) Box key: vertex | order of labelling | final time; temporary times below.

Question number



Shortest time from Aysbury to Groomsport = 34 minutes


(ii) Route: Aysbury → Beacon → Drilly → Firshaw → Groomsport

(A B D F G), minimum time 34 minutes

(iii) Aysbury to Evensby (A B D E): 25 minutes

(iv) Drilly to Beacon is one way, so B → D can no longer be used.

Question number



New shortest time from Aysbury to Groomsport = 35 minutes

(route A C F G)

D4 (i)

p	q	p ⇒ q	~p	~q	~q ⇒ ~p
T	T	T	F	F	T
T	F	F	F	T	F
F	T	T	T	F	T
F	F	T	T	T	T

The columns for p ⇒ q and ~q ⇒ ~p are identical (T F T T),

so p ⇒ q and ~q ⇒ ~p are equivalent.

(ii)

Let X = (p ⇒ q) and (q ⇒ r)

p	q	r	p ⇒ q	q ⇒ r	X	p ⇒ r	X ⇒ (p ⇒ r)
T	T	T	T	T	T	T	T
T	T	F	T	F	F	F	T
T	F	T	F	T	F	T	T
T	F	F	F	T	F	F	T
F	T	T	T	T	T	T	T

Question number

F	T	F	T	F	F	T	T
F	F	T	T	T	T	T	T
F	F	F	T	T	T	T	T

The final column is T in every row, so

$[(p \Rightarrow q) \text{ and } (q \Rightarrow r)] \Rightarrow (p \Rightarrow r)$ is a tautology.

DS (i) Identity: $u = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix}$

(ii)

$\circ$	p	q	r	s	t	u
p	q	u	t	r	s	p
q	u	p	s	t	r	q
r	s	t	u	p	q	r
s	t	r	q	u	p	s
t	r	s	p	q	u	t
u	p	q	r	s	t	u

(iii)

Element	p	q	r	s	t	u
Period	3	3	2	2	2	1

(iv) Subgroups of G:

$\{u\}$, $\{u, r\}$, $\{u, s\}$, $\{u, t\}$, $\{u, p, q\}$,

$\{p, q, r, s, t, u\}$ (G itself)

Permission to reproduce all copyright material has been applied for.
In some cases, efforts to contact copyright holders may have been unsuccessful and CCEA
will be happy to rectify any omissions of acknowledgement in future if notified.

12931/4