



Rewarding Learning

ADVANCED SUBSIDIARY (AS)
General Certification of Education
2023

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--	--

Physics

StemUp Model Solution

Assessment Unit AS 2

assessing

Module 2: Waves, Photons
and Astronomy



[SPH21]

SPH21

WEDNESDAY 24 MAY, AFTERNOON

TIME

1 hour 45 minutes.

INSTRUCTIONS TO CANDIDATES

Write your Centre Number and Candidate Number in the spaces provided at the top of this page.

You must answer the questions in the spaces provided.

Do not write outside the boxed area on each page or on blank pages.

Complete in black ink only. **Do not write with a gel pen.**

Answer **all ten** questions.

INFORMATION FOR CANDIDATES

The total mark for this paper is 100.

Figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question.

Your attention is drawn to the Data and Formulae Sheet which is inside this question paper.

You may use an electronic calculator.

13548



1 (a) Water waves are classified as transverse waves.

(i) State another example of a transverse wave.

Light (any electromagnetic wave)

[1]

(ii) State an example of a longitudinal wave.

Sound wave

[1]

(iii) With reference to the motion of the particles of the medium through which the wave passes, what distinguishes a transverse wave from a longitudinal wave?

In a transverse wave the particles oscillate perpendicular to the direction of energy transfer.

In a longitudinal wave they oscillate parallel to it.

[3]

(b) A graph of displacement against time for a water wave is shown in Fig. 1.1.




Fig. 1.1

(i) On Fig. 1.1, label the amplitude A of the wave.

[1]



- (ii) Use the graph in **Fig. 1.1** to determine an accurate value for the period of the wave.

3 complete waves take 2.0 s

$$T = 2.0 / 3$$

$$T = 0.67 \text{ s}$$

Period = 0.67 s [2]

- (iii) The wave is travelling at a speed of 3.50 m s^{-1} . Calculate the wavelength of the wave.

$$f = 1 / T = 1 / 0.667 = 1.5 \text{ Hz}$$

$$v = f\lambda, \text{ so } \lambda = v / f$$

$$\lambda = 3.50 / 1.5$$

$$\lambda = 2.33 \text{ m}$$

Wavelength = 2.33 m [3]

- (iv) On **Fig. 1.1**, draw a wave of the same amplitude and period but 90° out of phase with the wave shown. [2]

[Turn over]




2 Describe an investigation that will allow you to accurately determine the focal length of a converging lens using a graphical method.

Include in your answer:

- a labelled sketch of the apparatus required;
- the measurements which should be taken;
- the graph that you would draw using your results;
- how the graph is used to find an accurate value for the focal length of the lens.

(a) A labelled sketch of the apparatus required:



[2]

(b) The measurements which should be taken;

Move the screen until a sharp image of the object is seen. Measure the object distance u (object to lens) and the image distance v (lens to screen) with the metre rule.

Repeat for at least 5 different values of u (all greater than f).

[3]



(c) The graph you would draw using your results;

A graph of $1/v$ (y-axis) against $1/u$ (x-axis): a straight line of gradient

-1 .

[1]

(d) How the graph is used to find an accurate value for the focal length of the lens.

Both intercepts (on the $1/v$ and $1/u$ axes) are equal to $1/f$, so $f =$

$1/\text{intercept}$.

Find f from each intercept and take the average of the two values.

[2]

[Turn over]



- 3 (a) Explain the difference between Doppler red-shift and cosmological red-shift of light from distant galaxies. Refer to the source and the observer in each case.

Doppler red-shift is caused by the source (galaxy) moving away from the observer through space (relative motion of source and observer).

Cosmological red-shift is caused by the expansion of space itself between the source and the observer, stretching the wavelength.

[2]

- (b) The spectrum of light from a distant galaxy contains a spectral line of wavelength 398.6 nm. The same line, when measured in the laboratory, has a wavelength of 393.3 nm.

- (i) Calculate the recession speed of the galaxy. Give your answer to three significant figures.

$$\Delta\lambda = 398.6 - 393.3 = 5.3 \text{ nm}$$

$$z = \Delta\lambda / \lambda = 5.3 / 393.3 = 0.0135$$

$$v = zc = 3.00 \times 10^8 \times (5.3 / 393.3)$$

$$v = 4.04 \times 10^6 \text{ m s}^{-1}$$

Recession speed = 4.04×10^6 m s⁻¹

[3]



(ii) Use Hubble's law to estimate the distance to this galaxy.

$$v = H_0 d, \text{ so } d = v / H_0$$

$$d = 4.04 \times 10^6 / 2.4 \times 10^{-18}$$

$$d = 1.7 \times 10^{24} \text{ m}$$

Distance = 1.7×10^{24} m [2]

(iii) Why is this considered to be an estimate of the distance?

The Hubble constant H_0 is only an estimate (its value is
uncertain), so d is an estimate. [1]



- 4 The wavelength of monochromatic laser light can be calculated using Young's double-slit interference experiment. An arrangement for carrying out the experiment is shown in **Fig. 4.1**.




Fig. 4.1

- (a) The light emerging from a laser is coherent. Explain what is meant by coherent in this context.

The waves have a constant phase difference: they are in phase and have the same frequency.

[1]

In one such experiment, the interference pattern of bright and dark fringes, shown in **Fig. 4.2**, is observed on a screen placed 2.50 m from the double-slit. The separation of the slits is 0.27 mm.




Fig. 4.2



(b) Use the data to calculate the wavelength of the laser light.

6 dark fringes have 5 fringe spacings: $y = 30.5 / 5 = 6.1 \text{ mm}$

$$y = 6.1 \times 10^{-3} \text{ m}, a = 0.27 \times 10^{-3} \text{ m}, d = 2.50 \text{ m}$$

$$\lambda = ay / d = (0.27 \times 10^{-3} \times 6.1 \times 10^{-3}) / 2.50$$

$$\lambda = 6.6 \times 10^{-7} \text{ m}$$

Wavelength = 6.6×10^{-7} m

[4]



- 5 (a) A person uses 'reading glasses' in order to correct hypermetropia (long sight). Each lens has a power +2.0 dioptres. While wearing the glasses, the person's near point is 25 cm.

- (i) Show that the focal length of each lens in the glasses is +50 cm.

$$P = 1 / f, \text{ so } f = 1 / P$$

$$f = 1 / 2.0 = 0.50 \text{ m}$$

$$f = +50 \text{ cm}$$

[2]

- (ii) When an object is placed at a distance of 25 cm from the glasses, describe the image produced.

Virtual

Upright (erect)

Magnified (enlarged), formed 50 cm from the lens

[3]

- (iii) Complete **Table 5.1** to show the range of vision with and without the glasses. Show any calculations in the space below.

With glasses, an object at 25 cm gives a virtual image at v:

$$1/u + 1/v = 1/f: 1/25 + 1/v = 1/50, \text{ so } v = -50 \text{ cm}$$

so near point without glasses = 50 cm

with glasses, far point = f = 50 cm (image formed at infinity)

Table 5.1

	near point	far point
without the glasses	<u>50</u> cm	infinity
with the glasses	25 cm	<u>50</u> cm

[3]



(b) **Fig. 5.2** shows a diverging lens with an object OA placed perpendicular to the principal axis. The points F and F' are the principal foci of the lens.

On **Fig. 5.2**, draw suitable construction rays to locate the image of OA. Label the image IB. Show a suitable position of the eye for viewing the image.




Fig. 5.2

[5]

[Turn over]



- 6 (a) Explain what is meant by the photoelectric effect.

When photons of electromagnetic radiation are incident on a metal surface, electrons (photoelectrons) are ejected from the surface, provided the photon energy is greater than the work function (frequency above the threshold frequency).

[3]

- (b) A piece of zinc metal is represented in **Fig. 6.1**. It contains fixed ions and delocalised electrons as shown. The threshold frequency of zinc is 1.04×10^{15} Hz.




Fig. 6.1

- (i) Calculate the work function of zinc, giving your answer in eV.

$$\phi = hf_0$$

$$\phi = 6.63 \times 10^{-34} \times 1.04 \times 10^{15}$$

$$\phi = 6.90 \times 10^{-19} \text{ J}$$

$$\phi = 6.90 \times 10^{-19} / 1.60 \times 10^{-19} = 4.31 \text{ eV}$$

Work function = 4.31 eV

[4]



- (ii) When a piece of zinc is illuminated by radiation of a single frequency, the emitted particles have a range of kinetic energies. Explain why this is the case.

Electrons are at different positions (depths) within the metal.

Electrons below the surface must use up some energy to reach

the surface, so they are emitted with less than the maximum

kinetic energy.

[2]

[Turn over]



- 7 A simplified energy level diagram for an atom of hydrogen is shown in Fig. 7.1.




Fig. 7.1

- (a) Explain why the energy levels have negative values.

Energy must be supplied to remove an electron from the atom; a free electron (level E) has zero energy.

[2]

- (b) An electron can move between energy levels within the atom.

- (i) If a photon of wavelength 652 nm is emitted, between which energy levels has the electron moved?

$$\lambda = 652 \text{ nm} = 652 \times 10^{-9} \text{ m}$$

$$E = hc / \lambda = (6.63 \times 10^{-34} \times 3.00 \times 10^8) / 652 \times 10^{-9}$$

$$E = 3.05 \times 10^{-19} \text{ J}$$

$$C \text{ to } B: -2.40 \times 10^{-19} - (-5.45 \times 10^{-19}) = 3.05 \times 10^{-19} \text{ J}$$

Photon emitted, so the electron falls from C down to B.

Electron moved from C to B

[5]



- (ii) An electron in level A is struck by a photon of energy $1.75 \times 10^{-18} \text{ J}$. State which transition might occur, if any, and why?

No transition: $1.75 \times 10^{-18} \text{ J}$ is not exactly equal to the energy

difference between A and any other level. [2]

- (c) Electron transitions between energy levels are involved in the production of X-rays for medical imaging. This is a result of high energy electrons striking a metal anode.

- (i) What do the letters CT represent in CT scanning?

Computed Tomography [1]

- (ii) State three differences between CT scanning and conventional X-ray imaging.

X-ray images are 2-D, while CT scans give 3-D images (slices through the body).

CT needs an X-ray tube rotating around the patient; a conventional X-ray does not.

CT gives a higher radiation dose than a conventional X-ray. [3]

[Turn over]



- 8 (a) Calculate (i) the momentum and (ii) the velocity of an electron with a de Broglie wavelength $3.17 \times 10^{-11} \text{ m}$.

$$\lambda = h / p, \text{ so } p = h / \lambda$$

$$p = 6.63 \times 10^{-34} / 3.17 \times 10^{-11} = 2.09 \times 10^{-23} \text{ kg m s}^{-1}$$

(i) Momentum = 2.09×10^{-23} kg m s⁻¹ [2]

$$p = mv, \text{ so } v = p / m$$

$$v = 2.09 \times 10^{-23} / 9.11 \times 10^{-31} = 2.30 \times 10^7 \text{ m s}^{-1}$$

(ii) Velocity = 2.30×10^7 m s⁻¹ [2]

- (b) Diffraction is the name given to the spreading of waves after they pass through an opening. What is the condition under which maximum diffraction occurs?

When the wavelength of the wave is equal to the width of the opening (aperture). [1]



- (c) In an experiment to demonstrate electron diffraction, a narrow beam of electrons is directed at a thin metal foil.

- (i) The beam of electrons is produced by an 'electron gun', where a high voltage is applied between a cathode and an anode in order to accelerate the electrons.

Describe and explain the effect on the diffraction pattern when the accelerating voltage is increased.

The radius of the diffraction rings decreases (rings closer together).

The electrons are given a greater velocity (momentum), so their de Broglie wavelength ($\lambda = h/p$) is shorter and they diffract less.

[2]

- (ii) In this demonstration, the thin foil acts as a diffraction grating. What is the effective number of lines per mm if the second order diffraction occurs at an angle of 40° ?

The de Broglie wavelength of the electron is $3.17 \times 10^{-11} \text{ m}$.

$$d \sin \theta = n\lambda, \text{ with } n = 2$$

$$d = 2 \times 3.17 \times 10^{-11} / \sin 40^\circ$$

$$d = 9.86 \times 10^{-11} \text{ m}$$

$$\text{lines per metre} = 1 / d = 1.01 \times 10^{10} \text{ m}^{-1}$$

$$\text{lines per mm} = 1.01 \times 10^{10} / 1000 = 1.01 \times 10^7$$

Lines per mm 1.01×10^7

[4]

[Turn over]



- 9 (a) State Snell's law of refraction.

For light passing between a given pair of media,

the ratio $\sin i / \sin r$ is a constant (equal to the refractive index).

[2]

- (b) In an experiment to verify Snell's law, the path of a ray of light was traced as it passed from air through a rectangular glass block and out into the air. This is shown in Fig. 9.1.




Fig. 9.1

Use the angles given to show that the refractive index of the glass is 1.51.

$$n = \sin i / \sin r$$

$$n = \sin 60^\circ / \sin 35^\circ = 0.8660 / 0.5736$$

$$n = 1.510 = 1.51$$

[2]



- (c) A ray of monochromatic light passing through a section of an optical fibre is shown in **Fig. 9.2**. The ray travels from air into the core material with an incident angle A.

The core material is glass of refractive index 1.51 and the cladding material surrounding the core is of refractive index 1.45.

The ray is incident on the core–cladding boundary at the critical angle C for this pair of materials.




Fig. 9.2

- (i) What is the speed of light in the core material?

$$n = c / v, \text{ so } v = c / n$$

$$v = 3.00 \times 10^8 / 1.51$$

$$v = 1.99 \times 10^8 \text{ m s}^{-1}$$

Speed = 1.99 × 10⁸ m s⁻¹ [3]

- (ii) What is the refractive index for light travelling from the core to the cladding?

$$n \text{ (core to cladding)} = n_{\text{cladding}} / n_{\text{core}}$$

$$= 1.45 / 1.51$$

$$= 0.960$$

Refractive index = 0.96 [3]

[Turn over]



- 10 The speed of sound was measured using a resonance tube arrangement as shown in Fig. 10.1. A loudspeaker was placed above the open end of the tube. The length L of the tube is 75 cm. The signal frequency was raised from 0 Hz until an exceptionally loud sound was heard for the first time. This frequency is called the fundamental frequency f_0 .




Fig. 10.1

- (a) Resonance is the effect which results in an exceptionally loud sound being heard. Explain why resonance occurs.

The driving frequency of the signal equals the natural frequency of the air column in the tube, so the incident and reflected waves superpose to form a standing wave of maximum amplitude.

[2]



(b) The method described will identify the first position of resonance.

(i) On **Fig. 10.1** label the position of a node with the letter N and the position of an antinode with the letter A for the first position of resonance. [1]

(ii) Describe how the air particles in the resonance tube are behaving at N and A.

N *The air particles do not vibrate (zero amplitude).*

A *The air particles vibrate with maximum amplitude.* [2]

(iii) The fundamental frequency f_0 is 114 Hz when the first position of resonance is identified. Use this information to calculate the speed of sound in air.

$$\text{First resonance: } L = \lambda/4$$

$$\lambda = 4L = 4 \times 0.75 = 3.00 \text{ m}$$

$$v = f\lambda = 114 \times 3.00 = 342 \text{ m s}^{-1}$$

Speed = 342 m s⁻¹ [3]

(iv) The frequency is increased from 114 Hz until the next position of resonance is identified. Calculate the frequency at which this will occur.

$$\text{Next resonance: } L = 3\lambda/4, \text{ so } f = 3f_0$$

$$\lambda = 4 \times 0.75 / 3 = 1.00 \text{ m}$$

$$f = v / \lambda = 342 / 1.00 = 342 \text{ Hz}$$

Frequency = 342 Hz [2]

THIS IS THE END OF THE QUESTION PAPER



BLANK PAGE
DO NOT WRITE ON THIS PAGE

13548





BLANK PAGE

DO NOT WRITE ON THIS PAGE

13548



DO NOT WRITE ON THIS PAGE

For Examiner's use only	
Question Number	Marks
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	

Total Marks	
-------------	--

Examiner Number

Permission to reproduce all copyright material has been applied for.
In some cases, efforts to contact copyright holders may have been unsuccessful and CCEA
will be happy to rectify any omissions of acknowledgement in future if notified.

SPH21/5
276070





Rewarding Learning

ADVANCED SUBSIDIARY
General Certificate of Education

Physics

Assessment Units AS 1 and AS 2

[SPH11/SPH21]

DATA AND FORMULAE SHEET

Data and Formulae Sheet for AS 1 and AS 2

Values of constants

speed of light in a vacuum	$c = 3.00 \times 10^8 \text{ m s}^{-1}$
elementary charge	$e = 1.60 \times 10^{-19} \text{ C}$
the Planck constant	$h = 6.63 \times 10^{-34} \text{ J s}$
mass of electron	$m_e = 9.11 \times 10^{-31} \text{ kg}$
mass of proton	$m_p = 1.67 \times 10^{-27} \text{ kg}$
acceleration of free fall on the Earth's surface	$g = 9.81 \text{ m s}^{-2}$
electron volt	$1 \text{ eV} = 1.60 \times 10^{-19} \text{ J}$
the Hubble constant	$H_0 \approx 2.4 \times 10^{-18} \text{ s}^{-1}$

Useful formulae

The following equations may be useful in answering some of the questions in the examination:

Mechanics

conservation of energy	$\frac{1}{2} mv^2 - \frac{1}{2} mu^2 = Fs$
	for a constant force

Waves

two-source interference	$\lambda = \frac{ay}{d}$
diffraction grating	$d \sin \theta = n\lambda$

Light

lens equation

$$\frac{1}{u} + \frac{1}{v} = \frac{1}{f}$$

Electricity

terminal potential difference

$$V = E - Ir \text{ (e.m.f., } E; \text{ Internal Resistance, } r)$$

potential divider

$$V_{\text{out}} = \frac{R_1 V_{\text{in}}}{R_1 + R_2}$$

Particles and photons

Einstein's equation

$$\frac{1}{2} m v_{\text{max}}^2 = hf - hf_0$$

de Broglie equation

$$\lambda = \frac{h}{p}$$

Astronomy

red shift

$$z = \frac{\Delta\lambda}{\lambda}$$

recession speed

$$z = \frac{v}{c}$$

Hubble's law

$$v = H_0 d$$

