CAMBRIDGE INTERNATIONAL AS & A LEVEL BIOLOGY · 9700

BIOLOGY



MARK SCHEME

12 ENERGY AND RESPIRATION

SCORE	PERCENTAGE
/	%

STEMUP

These questions belong to CIE, not to us.

The examination questions, mark schemes and examiner comments in this pack were written and published by Cambridge International (CIE). They are past assessment materials, reproduced here so that students can revise from them.

What we do not own

STEMUP is not affiliated with CIE and does not own, and makes no claim to own, any of the question, mark scheme or examiner report material in this pack. CIE has not endorsed, approved or been involved in the production of it. CIE and the qualification titles used here belong to their respective owners and appear for identification only.

Where to find the originals

CIE publishes its past papers, mark schemes and examiner reports. Always check the current syllabus at cambridgeinternational.org before you revise.

If you hold the rights to any material in this pack and would like it amended or removed, contact admin@stemup.co.uk and we will act on it.

Paper 1 Multiple Choice Answers: Energy and Respiration

Question	Paper	Q. No.	Answer
1602	9700_s15_qp_12	40	C
1603	9700_s15_qp_13	38	C
1604	9700_s15_qp_13	40	A
1605	9700_w15_qp_12	40	B

Q1.

2 (a) provides energy;
suitable examples;
e.g. muscle contraction, protein synthesis, DNA replication, cell movement, active transport

3

(b)

substrate level phosphorylation	cytoplasm (in glycolysis);
	matrix of mitochondria (in Krebs cycle);
oxidative phosphorylation	inner membrane of mitochondria/cristae;

2 max

(c) oxidative phosphorylation more than substrate level phosphorylation;
ref. to quantity, e.g. 32/34 vs. 4/6 per glucose;

2

(d) requires proton gradient produced by ETC;
with no oxygen ETC does not occur/no electron flow;
NAD cannot be reformed/NADH cannot be oxidised;
oxygen combines with electron/proton/oxygen final acceptor in ETC;

3 max

Total: 10

Q2.

STEMUP.CO.UK

Question 2

(a) **(i)** correctly indicated on inner membrane ;
(ii) correctly indicated in matrix ;

2

(b) folded inner membrane / cristae ;
increases surface area available ;

intermembrane space ;
allows accumulation of H^+ ;

impermeability of inner membrane to H^+ ;
maintains H^+ gradient / H^+ only go through channels ;

stalked particles / ATPase ;
channel for H^+ / ATP synthesis ;

linear arrangement of ETC on inner membrane ;
greater efficiency ;

4 max

(c) (no) oxygen to combine with e^- / H^+ / H / 2H / proton ;
at the end of the ETC ;
no H^+ gradient produced ;
no ATP synthesized / oxidative phosphorylation does not occur ;
no NAD regenerated / NADH not oxidized ;
stops Krebs cycle ;

3 max

Total: 9

Q3.

STEMUP.CO.UK

Question			Marks
1	(a)	a nucleotide ;	
		with <u>three</u> phosphate groups ;	
		an organic / nitrogenous base / adenine ;	
		a pentose sugar / ribose ;	
		ref. ester linkages / covalent bonds ;	**3 max**
	(b)	synthesized from ADP and P_i ;	
		soluble molecule ;	
		diffuses rapidly / transported easily ;	
		on hydrolysis / removal of (third) phosphate ;	
		energy released / 30.5 kJ (mol^{-1}) ;	
		ref.(idea) intermediary (between energy yielding and energy requiring reactions) ;	**3 max**
	(c)	oxidative phosphorylation ;	
		$NADH_2$ to, cristae / inner membrane ;	
		oxidised to NAD ;	
		ref. transfer of electrons to electron carriers / ETC ;	
		H^+ pumped into intermembrane space ;	
		ref. to H^+ gradient ;	
		H^+ (diffuses) through ATP synthase / stalked particle ;	
		results in ADP and P_i to ATP ;	
		ref. chemiosmosis ;	
		ref. substrate level phosphorylation ;	**4 max**

Q4.

STEMUP.CO.UK

Question		Expected Answers	Marks
1	(a)	cytoplasm ;	[1]
	(b)	hexose bisphosphate / phosphorylated 6C sugar / fructose bisphosphate ;	[1]
	(c)	hexose energy rich ; does not react easily / AW ; phosphorylation activates hexose ; maintains concentration gradient of glucose inside and outside cell ;	[2 max]
	(d)	does not enter Krebs cycle ; decarboxylated / CO_2 released ; forms ethanal ; reduced / ref: reduced NAD ; to ethanol ; reduced NAD does not enter ETC ; ref: alcohol dehydrogenase ; irreversible ;	[4 max]
			[Total: 8]

Q5.

2

process	major products
glycolysis	ATP ; pyruvate ; reduced NAD ;
Krebs cycle	ATP ; reduced NAD / reduced FAD ; CO_2 ;
oxidative phosphorylation	ATP ; water ; NAD / FAD ;

[8 max]

R NADP throughout

[Total: 8]

Q6.

STEMUP.CO.UK

1 (a) (i) 18 ; [1]

(ii) 0.72 ;

allow ecf from (i) [1]

(b) 1 RQ value falls steeply, initially / 40–80 min ;

2 then, very little change / AW ;

3 sugar / carbohydrate, metabolised at start ; **A** named carbohydrate

4 then fat metabolised ;

5 (due to) fasting / carbohydrate running out ; [4 max]

(c) 1 increase in rate of respiration ;

2 kinetic energy increases / more enzyme-substrate complexes / enzyme activity increases ;

3 effects of too high a rise in temperature ; e.g. denaturation of enzymes

4 AVP ; e.g. $Q_{10} = 2$ [2 max]

[Total: 8]

Q7.

STEMUP.CO.UK

7 (a) (i) glycolysis ; [1]

(ii) cytoplasm/cytosol ; [1]

(iii) 4 ; **A** 4 – 2 = 2 [1]

(b) (i) inner membrane/cristae/stalked particles ; [1]

(ii) 1 reduced, NAD/FAD ;

2 dehydrogenase enzymes ;

3 release hydrogen ; **A** H **R** H_2/H^+

4 hydrogen splits into proton and electron ;

5 electrons flow down, ETC/AW ;

6 energy released ;

7 protons pumped (across inner membrane/from matrix) ;

8 into intermembrane space ;

9 proton gradient ;

10 protons pass through, ATP synthase/stalked particle ;

11 oxygen final, hydrogen/proton, acceptor ; [5 max]

(c) (i) nuclei **and** ribosomes ; [1]

(ii) 1 glycolysis, does not occur in mitochondrion/only occurs in cytosol or cytoplasm ;

2 pyruvate produced in glycolysis ;

3 pyruvate can enter mitochondrion/glucose cannot enter mitochondrion ;

4 carbon dioxide produced/decarboxylation, in, Krebs/link reaction ; [3 max]

(iii) 1 cyanide, inhibits cytochrome oxidase is a non-competitive inhibitor ;

2 reduced NAD not oxidised/AW ;

3 Krebs cycle stops ;

4 alternative H acceptor needed/pyruvate is H acceptor/pyruvate is reduced ; **R** H^+

5 lactate produced in cytoplasm ;

6 by anaerobic respiration ; [3 max]

[Total: 16]

Q8.

STEMUP.CO.UK

7 **(a)** **(i)** removal of, carbon dioxide/carboxyl group ;
removal of hydrogen ; [2]

(ii) **P** and **Q** ; [1]

(b) **(i)** 3 ; [1]

(ii) 1 inner mitochondrial membrane/cristae ;

2 dehydrogenase enzymes ;

3 release hydrogen ;

4 hydrogen splits into protons and electrons ;

5 electrons flow down, ETC/Electron Transfer Chain/AW ;

6 energy released ;

7 protons pumped across (inner membrane) ;

8 into intermembrane space ;

9 proton gradient ;

10 protons pass through, ATP synthase/stalked particles ;

11 ATP formed ; *linked to 10*

12 oxygen (final), hydrogen/proton and electron, acceptor ; *max 4* [5 max]

(c) 1 pyruvate converted to ethanal ;

2 ethanal reduced ;

3 by reduced NAD ;

4 NAD, oxidised/regenerated ;

5 allows glycolysis to continue ;

6 ethanal dehydrogenase ;

7 ethanol formed ;

8 prevents H^+ from lowering pH ; [4 max]

STEMUP.CO.UK

(d) 1 no, decarboxylation/carbon dioxide removed ; **A** ora

2 single step ;

3 lactate dehydrogenase ;

4 reversible ; [3 max]

[Total: 16]

Q9.

7 **(a)** active transport ;
ribose ;
water ;
hydrolysis ; **A** dephosphorylation
heat ; [5]

(b) **(i)** (converted to) glycogen / lipid ;
(used in) glycolysis / respiration ; [1 max]

(ii) *anaerobic*
1. less ATP / only 2 ATP ;
2. per mol glucose ;
3. lactate still contains energy / only glycolysis involved / stages other than glycolysis not involved ;
4. not sustainable / cannot go on indefinitely / AW ; [2 max]

(iii)

process	precise location
glycolysis	cytoplasm / cytosol ;
link reaction	mitochondrial matrix ;
Krebs cycle	mitochondrial matrix ;
oxidative phosphorylation	inner mitochondrial membrane / cristae ;

[4]

(iv) 1. cannot pass through phospholipid bilayer ;
2. too big to fit through (glucose's) protein channel ;
3. no specific transport protein ;
4. AVP ; e.g. used up as soon as it is made [2 max]

(v) oxygen debt ; [1]

[Total:15]

Q10.

STEMUP.CO.UK

6 (a) (i) decarboxylation ; [1]

(ii) dehydrogenation / oxidation ; [1]

(iii) substrate level phosphorylation ; [1]

(b) **K** – reduced NAD ; **A** NADH etc.
L – oxaloacetate ; [2]

(c) 1. hydrogens split into protons and electrons ;
2. electrons pass along ETC ;
3. energy released used to pump protons ;
4. (from matrix) to intermembrane space ;
5. inner membrane impermeable to protons ;
6. proton gradient forms ;
7. protons move down gradient ;
8. through ATP, synthase / ATP synthetase ; **R** ATPase
9. enzyme rotates ;
10. ATP produced ; [5 max]

[Total: 10]

Q11.

4 (a) (i) 1. ATP is made, in the electron transport chain/by oxidative phosphorylation;

2. oxygen is the final electron acceptor;

3. in the, inner membrane of the mitochondrion/cristae;

4. transfer of electron (between electron carriers) provides energy;

5. energy used to pump hydrogen ions (into intermembrane space);

6. creates proton gradient;

7. diffusion of hydrogen ions down their electrochemical gradient causes ATP to be synthesised;

8. ref. chemiosmosis/ATP synthase/stalked particles;

9. idea that if less oxygen (consumed/available) then fewer electrons transferred along the chain; [max 4]

(ii) 1. at high temperatures, reactions/enzyme activity/metabolism, faster;

2. because, molecules/enzymes/substrates, have more kinetic energy;

3. more frequent collisions;

4. therefore, respiration/Krebs cycle/electron transport chain/production of reduced NAD, take place at a faster rate;

5. idea of increase in rate of anabolic reactions (requiring more ATP); [max 3]

STEMUP.CO.UK

(b) (i) 1. oxygen consumed = oxygen inhaled – oxygen exhaled;

2. measure oxygen consumption at rest (x) **and** after exercise stops (y);

3. extra oxygen consumed/oxygen debt = y – x;

4. measure mass of lizard; [max 2]

(ii) 1. less (oxygen debt)(for *Varanus*); **ora**

2. difference is greater at higher temperatures;

3. any two comparative figures at one temperature including units; [3]
A 102.0 cm^3 O_2 kg^{-1} at 30°C and 40°C

(iii) 1. *Varanus* uses, less anaerobic/more aerobic, respiration (when running);

2. more ATP produced per glucose molecule;

3. able to run for long time;

4. good chance of catching prey; [max 3]

(iv) *assume Varanus throughout*
1. larger surface area, in lungs/for gas exchange;

2. more oxygen absorbed into blood (per unit time)/faster rate of gas exchange;

3. more oxygen supplied to muscles (so oxygen debt lower); [max 2]

[Total: 17]

Q12.

STEMUP.CO.UK

4 (a) (i) inner membrane / crista(e); [1]

(ii) 1. (electron comes from) hydrogen (atom); **R** H^+ / H_2

2. (from) reduced NAD / reduced FAD;

3. (from) dehydrogenation / oxidation, reactions;

4. (from substances in) Krebs cycle / link reaction / glycolysis;

5. in, matrix of mitochondrion / cytoplasm; [max 3]

(iii) 1. final electron acceptor / accepts electron from last carrier;

2. so carrier can be reduced again;

3. so electrons can keep flowing (along ETC) / so ETC can continue to work;

4. (oxygen) combines with H^+ to form water; [2 max]

(b) (i) 1. (when pump stops working), resting potential not maintained
or
pump usually maintains the resting potential;

2. (during resting potential) membrane polarised
or
positive charge outside (neurone) / negative charge inside (neurone) / -70mV inside neurone relative to outside / potential difference across membrane;

3. (when pump stops working), ions (only) move by diffusion;

4. Na^+ **into** the neurone;

5. outward diffusion of K^+ is limited / K^+ stay in neurone;

6. ref. non voltage-gated channels;

7. (eventually) inside of the neurone, becomes less negative / contains (relatively) more positive ions
or
there is a reduced potential difference across the membrane; [max 4]

(ii) 1. voltage gated (calcium) channels open;

2. (calcium ions move in) by diffusion / move down their concentration gradient; [2]

STEMUP.CO.UK

(c) (i) 1. Na^+ / K^+, cannot move through membrane;

2. so potential across membrane maintained even when pump stops / so membrane depolarisation does not happen;

3. calcium ions cannot enter cell;

4. so, (destructive) enzymes not activated; [max 2]

(ii) 1. gene (for protein channels), expressed less / switched off;

2. transcription, reduced / stopped;

3. AVP; e.g. reduced aerobic respiration / less ATP, for transcription [max 2]

[Total: 16]

Q13.

Question 2
(a)

	name of structure	stage of respiration
A	matrix	Krebs cycle ;
B	cristae / inner membrane **A** intermembrane space	oxidative phosphorylation/ETC ; **A** build up of protons

Penalise once if rows A and B are correct but swapped
If both structure names are correct (but stages incorrect) allow one mark

2

STEMUP.CO.UK

(b)
membranes separate from rest of cytoplasm ;
allows different pH ;
inner membrane attachment of stalked particles / ATPase ;
allows linear / ordered arrangement of carriers/ETC/respiratory chain ;
ref. to large internal surface area/AW ;
matrix contains enzymes;

3 max

(c)
carries / transfers protons/hydrogen(atoms) ;
and electrons ;
in/to ETC /FAD/respiratory chain;
ref. to dehydrogenation/oxidising ;
energy used to form ATP;
ref. to coenzyme ;
ref. alternative pathways (named);

3 max

(d)
light involved ;
occurs in chloroplasts/chlorophyll ;
on thylakoid membranes ;
ref. to cyclic and non-cyclic ;
photolysis of water/produces oxygen;

If oxidative phosphorylation stated
light not involved;
oxygen final hydrogen acceptor/oxygen not evolved;

3max

Total:11

Q14.

STEMUP.CO.UK

Question 2

(a) cytoplasm ;
matrix in mitochondria ; **2**

(b) coenzyme ;
carries electrons / protons / hydrogen ions / hydrogen / H / 2H / H^+ ; **R** H_2
to electron transfer chain / AW ;
from glycolysis / link reaction / Krebs cycle ;
role of NAD in conversion / oxidation of triose phosphate to pyruvate in glycolysis ;
role of NAD in anaerobic respiration ; **3 max**

(c) in absence of oxygen electron transfer chain does not work ;
oxygen final acceptor at end of electron transfer chain ;
reduced NAD cannot be oxidised ; **3**

(d) aerobic respiration produces more ATP / (ora);
to produce the same amount of ATP more glucose broken down in glycolysis ;
glycolysis is the only part of respiration used / no ETC or oxidative phosphorylation ; **2 max**

Total : 10

Q15.

3 **(a) 1** no increase below 40 au;

2 (most) rapid production above 60 au;

3 correct reference to Figs.; **2 max**

(b) 1 glucose to pyruvate/glycolysis;

2 pyruvate to lactate;

3 reference lactate dehydrogenase;

4 in absence/shortage of oxygen to muscles;

5 pyruvate acts as a hydrogen acceptor;

6 reduced NAD to NAD/NAD regenerated; **3 max**

(c) 1 lactate must be oxidised;

2 extra oxygen required;

3 this is the oxygen debt; *linked to point 2* **3**

(d) more anaerobic respiration/insufficient oxygen supply; **1**

Total 9

STEMUP.CO.UK

Q16.

Question 1

(a) matrix of mitochondria ; **[1]**

(b)(i) 3 sites labeled ;; *deduct one mark for each additional or missing label*

(ii) 5 sites labeled ;; **[3 max]**

(c) reduced NAD ;
ref. to ETC ;
oxidized / give up hydrogen ; **[3]**

(d) ref. to substrate level phosphorylation;
no proton gradient involved ;
no ATP synthase ;
no ETC ; **[1 max]**

Total [8]

Q17.

Question 3

(a) $RQ = \frac{\text{volume of carbon dioxide given off}}{\text{volume of oxygen taken up}}$; **[1]**

(b)(i) 18 H_2O ;
$18CO_2$; **[2]**

(ii) 18/26 ;
= 0.7 ; 2 marks for correct answer **[2]**

(iii) fatty acid A lipids / triglycerides / fat / oil ; **[1]**

(c) less C-C bonds ;
less C-H bonds ;
more oxygen ; **A** O **R** O_2 **[2 max]**

Total [8]

Q18.

STEMUP.CO.UK

Question			Expected Answers	Marks
1	(a)		*(carbohydrates)*	
		1	less reduced / less hydrogen / less C-H bonds ; **R** H_2	
		2	for, aerobic respiration / ETC / NAD / ATP ;	
		3	less energy ;	
		4	per, unit mass / mole ; *accept figs for 3 and 4*	
		5	carbohydrate has lower energy density ;; *accept as alternative to 3 & 4 for 2 marks*	**3 max**
	(b)		carbohydrate = 1.0 ;	
			lipid = 0.6 – 0.8 ;	**2**
	(c)		RQ remains stable between 3°C and 10°C / AW ;	
			rise between 10°C and, 20°C / 25°C ;	
			0.74 to, 0.76 / 0.8 ; *accept difference for figs marks*	
			sharp rise, between 25°C and 27°C / after 25°C ;	
			0.8 to 0.91 / peaks at 0.91 ; *3 max*	
			at low temperatures hamster uses lipids ;	
			reason ; e.g. more heat generated from lipid respiration	
			at higher temperatures more carbohydrates are used ;	**4 max**
	(d)		anaerobic respiration / conversion of carbohydrate to fats as animal hibernates;	**1**
				[Total: 10]

Q19.

7	(a)	1	provides, H^+ / protons / protons and electrons ; ***A*** *hydrogen* ***R*** H_2 ***R*** *produce* H^+	
		2	forms, reduced NAD / reduced FAD ; ***A*** *NAD / FAD, accepts* H^+	
		3	passed to ETC / cytochromes ;	
		4	oxidative phosphorylation ;	
		5	cytochrome oxidase ;	
		6	forms water (with oxygen) ;	[3 max]
	(b)	**(i)**	(initial) steep rise up to 40 (μmol) Al ;	
			2 paired figs ;	
			ref. plateau above 40 (μmol) Al ;	[2 max]
		(ii)	(initially) Al is, activator / cofactor / coenzyme ;	
			detail of shape change of enzyme ;	
			enzyme / substrate, limiting, after 40 (μmol) Al / high conc Al ; ***A*** *end product inhibition after 40 (μmol) Al*	[2 max]
				[Total: 7]

Q20.

STEMUP.CO.UK

6 (a) (i) adenine ;

(ii) ribose ; **R** pentose [2]

(b) 1 energy is released when it is hydrolysed ; **A** equation **A** joules for energy

2 easily hydrolysed ;

3 (energy) used in, processes / reactions ; **A** named process

4 rapid turnover ;

5 links catabolic and anabolic reactions / AW ;

6 found in, most cells / all organisms ;

7 soluble so easily moved (within cell) ;

8 ATP produced from variety of reactions ; **A** named reactions [4 max]

(c) 1 ETC / inner mitochondrial membrane / crista / stalked particles ;

2 grana / thylakoids / inner chloroplast membrane ;

3 cytoplasm / cytosol ;

4 mitochondrial matrix ; [2 max]

[Total: 8]

Q21.

STEMUP.CO.UK

6 (a) (i) phosphorylation ; [1]

(ii) lysis ; [1]

© University of Cambridge International Examinations 2011

(iii) dehydrogenation / oxidation ; *ignore reduction of NAD* [1]

(b) provides activation energy / AW ;
for it to split / AW ; [2]


(c) 1. decarboxylated / carbon dioxide given off ;
2. ethanal produced ;
3. ethanal reduced ;
4. by reduced NAD ;
5. to ethanol ;
6. dehydrogenase ; [4 max]

[Total: 9]

Q22.

STEMUP.CO.UK

6 (a)



[5]

(b) *in mammals*

1. lactate produced / no ethanol produced ;
2. no, decarboxylation / carbon dioxide released ;
3. single step ;
4. lactate dehydrogenase ;
5. reversible ;

[3 max]

(c) *in anaerobic respiration*

1. only glycolysis occurs / Krebs cycle stops / link reaction stops ;
2. glucose, not fully broken down / still contains energy ;
3. pyruvate does not enter mitochondrion ;
4. (no oxygen) so no final electron acceptor (in ETC) ;
5. ETC stops ;
6. no oxidative phosphorylation ;

[3 max]

[Total: 11]

Q23.

STEMUP.CO.UK

8 (a) **X** = crista(e) / inner membrane ;
Y = matrix ; [2]

(b) (i) raise chemical PE of glucose / provide activation energy / AW ; [1]

(ii) removes hydrogen / hydrogen carrier / coenzyme ; [1]

(iii) 4 ; **A** net 2 [1]

(iv) dehydrogenation ; **A** oxidation
decarboxylation ;
accept 'oxidative decarboxylation' for two marks [2]

(v) matrix ; [1]

(vi) 1. accepted by NAD ;
2. passed to ETC ;
3. for oxidative phosphorylation ;
4. ref. proton pump / chemiosmosis ; [2 max]

(c) 1. found in all organisms ;
2. loss of phosphate / hydrolysis, leads to, energy release / release of 30.5 kJ (per mole) ;
3. ADP + Pi ⟷ ATP / reversible reaction ;
4. small packets of energy ;
5. small / water soluble, so can move around cell ;
6. (used by cells as) immediate energy donor ;
7. link between energy yielding and energy requiring reactions / AW ;
8. high turnover ;
9. example of use ; e.g. active transport / muscle contraction / Calvin cycle / protein synthesis [5 max]

[Total:15]

Q24.

STEMUP.CO.UK

8 **(a) (i)** cytoplasm / cytosol ; [1]

(ii) 1 NAD regenerated ;
2 so glycolysis can continue ;
3 to produce ATP ; [2 max]

(iii) lactate dehydrogenase ; [1]

(iv) *reaction* - condensation / polymerisation ;
bond - glycosidic ; [2]

(b) *in yeast*
1 decarboxylation / CO_2 removed ;
2 ethanal (as intermediate step) ;
3 ethanol produced ;
4 two steps (from pyruvate) ;
5 ethanol dehydrogenase ;
6 not a reversible reaction / ethanol cannot be converted back to pyruvate ;
7 *idea of* process less energy efficient ;
allow ***ora*** *for mp1, mp4, mp5, mp6 and mp7* [4 max]

(c) (i) carbon dioxide produced divided by oxygen consumed ;
volume / number of moles (of both gases) ; [2]

(ii) carbohydrate = 1.0 ;
lipid = 0.7 ; [2]

(iii) increase / go above one / infinity ; [1]

[Total: 15]

Q25.

3 **(a)** adenine / nitrogen(ous) base / purine ; **R** adenosine

ribose / pentose ; [2]

(b) 1. (cell uses) ATP as source of energy ;

2. ATP broken down ;

3. (so) cell must regenerate ATP ;

4. from ADP and Pi ;

5. ref. ADP / AMP, must be synthesised in the cell ; [max 2]

STEMUP.CO.UK

(c) (i) 1. palmitic acid has **more**, hydrogens / C-H bonds ;

2. per mole ;

3. hydrogens needed for, ATP production / chemiosmosis / oxidative phosphorylation ;
[max 2]

(ii) *alanine* – starvation / lack of fat or carbohydrate ;

lactate – after anaerobic respiration ; [2]

[Total: 8]

Q26.

3 (a) 1. oxidative phosphorylation ;

2. oxygen is **final** electron acceptor ;

3. reduced to water / accepts hydrogen ion to form water ; **A** equation

4. so electron transport chain can continue ; ora

5. increases ATP production ; ora

6. in absence of oxygen only glycolysis continues ; [max 3]

© Cambridge International Examinations 2013

(b) (i) 1. lipid releases most energy ;

2. because it has more, hydrogens / C-H bonds ;

3. per unit mass ;

4. hydrogens needed for, ATP production / chemiosmosis ; [max 3]

(ii) many more hydrogens available to, reduce / convert, oxygen to water ; [1]

[Total: 7]

Q27.

STEMUP.CO.UK

2 (a) (i) substrate level ;
protein synthesis / DNA replication / glycogenesis / polymerisation ;
active transport / movement of chromosomes / sliding filaments / movement of vesicles / AW ; [3]

(ii) water ; [1]

(iii) ATP, synthase / synthetase ; R ATPase [1]

Q28.

8 (a) (DNA for) transcription/ codes for mRNA ;

(ribosomes for) translation ;

synthesis of, respiratory enzymes/named enzyme/ inner membrane proteins ; [max 3]

(b)

correct order	letter of stage
1	**V**
2	**S**
3	**U**
4	**W**
5	**R**
6	**Q**
7	**X**
8	**T**

S U W all above **R** ;
S U W in correct order ;

Q X T all below **R** ;
Q X T in correct order ; [4]

STEMUP.CO.UK

(c) hydrolysis / dephosphorylation / exothermic / exergonic ; [1]

(d) anaerobic respiration ;

substrate level phosphorylation (in glycolysis);

at triose phosphate ⟶ pyruvate step ;

(net) gain of 2ATP (per glucose) ; **A** 2 used and 4 produced

pyruvate, reduced / gains hydrogens (from reduced NAD) ;

forming lactate ;

NAD regenerated / $NADH_2$ re-oxidised ;

this allows glycolysis to continue ;

I ethanol pathway [max 5]

[Total:13]

Q29.

STEMUP.CO.UK

8 (a) (i) receptors/hypothalamus, detect change in blood temperature ;

brain ;

(receptor/brain) sends impulses to effector ;

effector carries out response/example of response ;

blood temperature returns to normal ;

negative feedback ; [max 4]

(ii) larger SA:V ratio ;

lose (relatively) more heat ;

ref. more mitochondria to release heat energy ;

cannot carry out behavioural actions to get warm ;

infants cannot shiver ; [max 2]

(b) (i) **A** – ATP synthase/ATP synthetase/stalked particles ; **R** ATPase

B – inner membrane/ crista ; **I** phospholipid bilayer [2]

(ii) arrow going down from intermembrane space to matrix ; [1]

(iii) 1 **and** 3 ; [1]

(iv) water ; [1]

(v) fatty acids ; **A** lipid/fat/triglycerides [1]

[Total:12]

Q30.

5 (a) contains ribose (not deoxyribose) ;

has three phosphate groups (not one) ; [2]

STEMUP.CO.UK

(b) (i) *anaerobic – accept* ***ora*** *for aerobic*

1 *idea that* glucose not completely, broken down / oxidised

or

only glycolysis occurs ;

2 pyruvate / lactate / ethanol, still contains energy ;

3 ETC stops ;

4 (because) no oxygen to act as (final) electron acceptor ;

5 (so) no, Krebs cycle / link reaction / oxidative phosphorylation / chemiosmosis ; [max 3]

(ii) 1 lipid contains (relatively) more, hydrogen atoms / C-H ;

2 detail ; e.g. molecular formula of glucose <u>and</u> a lipid given

3 more reduced, NAD / FAD, produced ;

4 more electrons passed along ETC ;

5 more hydrogen ions pumped across inner mitochondrial membrane/ more hydrogen ions pumped into intermembrane space / steeper proton gradient ; [max 3]

[Total: 8]

Q31.

STEMUP.CO.UK

4 **(a)** *ignore ref. to energy currency*

1 *idea of* synthesis of complex substances or synthesis of named large molecule / anabolic reactions;

2 transport of substances against concentration gradient / active transport ;

3 movement qualified ; e.g. muscle contraction / cilia movement / locomotion

4 AVP ; e.g bioluminescence, electrical discharge, temperature regulation [max 2]

(b) **(i)** *both answers required for one mark*

A adenine **R** adenosine

B ribose / pentose ; [1]

(ii) 1 small ;

2 water soluble ;

3 easily transported around the cell ;

4 easily hydrolysed (to release energy) ;

5 (so) relatively large quantity of energy released / 30.5 $kJ\,mol^{-1}$;

6 *idea of*, rapid turnover / small cellular ATP content is sufficient for cell's requirements ; [max 3]

(c) **(i)** 1 less / decreased (aerobic respiration) ;

2 oxygen, is the final electron acceptor / needed for ETC ;

3 oxidative phosphorylation decreased / chemiosmosis decreased ;

4 regeneration of NAD / Kreb's cycle / link reaction, decreased ;

5 ATP synthesis decreases / ATP synthetase activity decreased ; [max 2]

(ii) more ATP produced (for population growth) ; [1]

(d) **(i)** 1 HB8 always does better than mutant HB8 ;

2 HB8 and mutant HB8 both do better in aerobic than in anaerobic conditions ;

3 data quote to support ;

for mp1
[950×10^6 per cm^3 v 900×10^6 per cm^3] **and** [490×10^6 per cm^3 v 410×10^6 per cm^3] **or** manipulated figures

for mp2
[950×10^6 per cm^3 v 490×10^6 per cm^3] **and** [900×10^6 per cm^3 v 410×10^6 per cm^3] **or** manipulated figures [max 2]

STEMUP.CO.UK

(ii) 1 both grow better in aerobic compared to anaerobic ;

2 *ref. to* significant difference found in mutant HB8 (aerobic compared to anaerobic) ;

3 data quote to support ;

for mp1
[880 × 10^6 per cm^3 v 460 × 10^6 per cm^3] **and** [840 × 10^6 per cm^3 v 50 × 10^6 per cm^3] **or** manipulated figures

for mp2
[840 × 10^6 per cm^3 v 50 × 10^6 per cm^3] **or** [460 × 10^6 per cm^3 v 50 × 10^6 per cm^3] **or** manipulated figures [max 2]

(iii) *idea that* HB8 is a better competitor than mutant HB8 ; **ora**

in mutant HB8 activity of, enzyme/nitrate reductase, is reduced ; [max 1]

[Total: 14]

Section B

1.

9 **(a)** 1 reduced, NAD / FAD ;

2 passed to ETC ;

3 inner membrane / cristae ;

4 hydrogen released (from reduced, NAD / FAD) ; **R** H2

5 split into electrons and protons ;

6 protons in matrix ;

7 electrons pass along, carriers / cytochromes ;

8 ref. redox reactions ;

9 ref. energy gradient ;

10 energy released ; **R** produced

11 protons (pumped) into intermembrane space ;

12 proton gradient ;

13 protons pass through (protein) channels ;

14 ATP synthase / stalked particles ;

15 ATP produced ;

16 chemiosmosis ;

17 electron transferred to oxygen ;

18 addition of proton (to oxygen) to form water / (oxygen) reduced to water ; [9 max]

if candidate mistakenly writes about photosynthesis only allow marking points 7, 8, 9, 10 and 15 to 5 max

STEMUP.CO.UK

(b) *in cytoplasm*

19 NAD, becomes reduced / accepts H ;

20 during glycolysis ;

in plants

21 pyruvate converted to ethanal ;

22 ethanal reduced ;

23 by reduced NAD ;

24 ethanol formed ;

in animals

25 pyruvate converted to lactate ;

26 by reduced NAD ;

27 in, liver / muscles ;

28 allows glycolysis to continue ; [6 max]

allow either 23 ***or*** *26*

[Total: 15]

2.

9 **(a)** 1. reduced, NAD / FAD ;

2. passed to ETC ;

3. inner membrane / cristae ;

4. hydrogen released (from reduced, NAD / FAD) ; **R** H_2

5. split into electrons and protons ;

6. electrons pass along, carriers / cytochromes ;

7. ref. energy gradient ;

8. energy released pumps protons into intermembrane space ;

9. proton gradient ;

10. protons pass through (protein) channels ;

11. ATP synthase / stalked particles ;

12. (ATP produced from) ADP and inorganic phosphate ;

13. electron transferred to oxygen ;

14. addition of proton (to oxygen) to form water / (oxygen) reduced to water ; [8 max]

STEMUP.CO.UK

(b) 15. organisms need energy, to stay alive / for metabolism / AW ;

16. ATP as, (universal) energy currency / described ;

17. light energy for photosynthesis ; **A** light dependent stage

18. light-dependent stage detail ;

19. light-independent stage detail ;

20. chemical energy ;

21. for anabolic reactions ;

22. named reaction; e.g. protein synthesis / starch formation

23. activation of glucose in glycolysis / described ;

24. active transport ;

25. detail; e.g. sodium - potassium pump /movement against a concentration gradient

26. mechanical energy / movement ;

27. detail ; e.g. muscle contraction / spindle

© University of Cambridge International Examinations 2012

28. temperature regulation ;

29. AVP ; e.g. bioluminescence / electrical discharge [7 max]

[Total: 15]

3.

STEMUP.CO.UK

9 (a) *Active transport or anabolic reactions*

1. ATP provides energy (linked to either) ; *ignore ref. to energy currency alone*

active transport

2. movement against concentration gradient ;
3. carrier / transport, protein (in membrane) ; *ignore pump*
4. binds to (specific) ion ;
5. protein changes shape ;

anabolic reactions

6. synthesis of complex substances from simpler ones ;
7. starch / cellulose / glycogen, from, monosaccharides / named monosaccharides / named sugar ;
8. glycosidic bonds ;
9. lipid / triglyceride, from fatty acids and glycerol ;
10. ester bonds ;
11. polypeptides / proteins, from amino acids ;
12. peptide bonds ;
13. other named polymer from suitable monomer ;
14. appropriate named bond ; *5 max*

[7 max]

STEMUP.CO.UK

(b) *general*

15. reduced NAD produced in glycolysis ; ***A*** glycolysis described

16. small amount of ATP produced in glycolysis ;

in yeast cells

17. pyruvate converted to ethanal ;

18. carbon dioxide released / decarboxylation ;

19. ethanal, reduced / accepts H ;

20. by reduced NAD ;

21. ethanol formed ;

in mammalian cells

22. pyruvate converted to lactate ;

23. by reduced NAD ;

© University of Cambridge International Examinations 2012

24. in, liver / muscle, cells ;

25. AVP ;;

26. e.g. reversible in mammal / irreversible in yeast / single step in mammal / more than 1 in yeast / reoxidised NAD allows glycolysis to continue / named enzyme

*only award **either** mp19 **or** mp23* [8 max]

[Total: 15]

4.

STEMUP.CO.UK

6 ***(a) Describe the main features of the Krebs Cycle. [9]***

(b) Explain the role of NAD in aerobic respiration. [6]

(a) 1 matrix;

2 of mitochondrion;

3 acetyl CoA combines with oxaloacetate;

4 to form citrate;

5 4C to 6C;

6 decarboxylation/produce CO_2;

7 dehydrogenation/oxidation;

8 $2CO_2$ released;

9 reduced NAD produced; *accept reduced coenzyme for one mark - annotate 9/10*

10 reduced FAD produced;

11 ATP produced;

12 series of steps/intermediates;

13 enzyme catalysed reactions;

14 oxaloacetate regenerated;

15 AVP; **9 max**

(b) 16 coenzyme;

17 for dehyrogenase;

18 reduced;

19 carries electrons;

20 and protons/H^+/H/hydrogen; **R** H_2/hydrogen molecules

21 from Krebs cycle;

22 and from glycolysis;

23 to cytochromes/electron transfer chain;

24 reoxidised/regenerated;

25 ATP produced;

26 3/2.5 (molecules of ATP) per reduced NAD; **6 max**

Total 15

STEMUP.CO.UK

5.

7 **(a)** Describe the transfer of energy to ATP during photosynthesis. [6]

(b) Describe the process of oxidative phosphorylation. [9]

[Total: 15]

(a)

1 light absorbed by chlorophyll / AW ;
2 ref. photosystems ;
3 ref. harvesting clusters / accessory pigments ;
4 reaction centre / P680 / P700 ;
5 excitation of electrons / AW ;
6 ETC ;
7 idea of different energy levels ;
8 ADP + Pi → ATP ;
9 cyclic / non-cyclic, photophosphorylation ;
10 chemiosmosis / ATP synthase / description ; **6 max**

(b)

11 reduced NAD / FAD ;
12 passed to ETC ;
13 hydrogens removed ; **R** H_2
14 split into H^+ and e^- ;
15 e^- passed to carriers ;
16 H^+ stays in mitochondrial matrix ;
17 oxygen final e^- carrier ;
18 joins with H^+ / reduced ; **R** H_2 / hydrogen
19 forms water ;
20 ref. energy levels of carriers ;
21 energy available to convert ADP and Pi to ATP ;
22 occurs three times (for each reduced NAD) / ref. total yield ;
23 chemiosmosis / ATP synthase / description ; **9 max**

[Total: 15]

6.

STEMUP.CO.UK

9	(a)	1	(glucose) phosphorylated by ATP ;	
		2	raises energy level / overcomes activation energy ;	
		3	hexose bisphosphate ;	
		4	lysis / splitting, of, glucose / hexose ; **R** sugar splitting	
		5	breaks down to two TP ; **A** GALP / GADP / G3P / PGAL	
		6	6C → 2 x 3C ;	
		7	dehydrogenation / description ;	
		8	2 NAD reduced formed (from each TP to pyruvate formed) ;	
		9	4 ATP produced / net gain of 2 ATP ;	
		10	pyruvate produced ;	
		11	reduced NAD → oxidative phosphorylation / redox ; *accept flow diagram*	[7 max]
	(b)	12	nucleotide ;	
		13	adenine + ribose / pentose + three phosphates ;	
		14	loss of phosphate leads to energy release / hydrolysis releases 30.5 kJ ;	
		15	ADP + Pi ↔ ATP (reversible reaction) ;	
		16	synthesised during, glycolysis / Krebs cycle / substrate level phosphorylation ;	
		17	synthesised, using electron carriers / oxidative phosphorylation / photophosphorylation ;	
		18	in, mitochondria / chloroplasts ;	
		19	ATP synthase / ATP synthetase ;	
		20	chemiosmosis / description;	
		21	used by cells as immediate energy donor ;	
		22	link between energy yielding and energy requiring reactions / AW ;	
		23	active transport / muscle contraction / Calvin cycle / protein synthesis ;	[8 max]
				[Total: 15]

7.

STEMUP.CO.UK

9	(a)	1	acetyl CoA combines with oxaloacetate ;	
		2	to form citrate ;	
		3	4C to 6C ;	
		4	decarboxylation / CO_2 released ;	
		5	dehydrogenation / oxidation / release of hydrogen ;	
		6	reduced NAD produced / NAD accepts hydrogen ;	
		7	reduced FAD produced / FAD accepts hydrogen ;	
		8	ATP produced ;	
		9	substrate level phosphorylation ;	
		10	series of, steps / intermediates ; **A** many named steps off a diagram	
		11	enzyme catalysed reactions ;	
		12	oxaloacaetate regenerated ;	
		13	occurs in mitochondrial matrix ;	[9 max]
			accept diagram	
	(b)	14	coenzyme ;	
		15	for dehydrogenase ;	
		16	reduced ;	
		17	carries, electrons and protons / hydrogen / NAD	
		18	from Krebs cycle ;	
		19	and glycolysis ;	
		20	to ETC / electron carrier chain / oxidation ;	
		21	reoxidised / regenerated hydrogen removed ;	
		22	ATP produced ;	[6 max]
				[Total: 15]

8.

STEMUP.CO.UK

10 **(a)** 1 ATP as universal energy currency ;

2 light energy needed for photosynthesis ;

3 ATP used conversion of GP to TP ;

4 ATP used to regenerate RuBP ;

5 (energy needed for) anabolic reactions ;

6 protein synthesis / starch formation / triglyceride formation ;

7 activation energy ;

8 (activate) glucose in glycolysis ;

9 active transport ;

10 example ; e.g. sodium / potassium pump

11 movement / locomotion ;

12 example ; e.g. muscle contraction / cilia beating

13 endocytosis / exocytosis / pinocytosis / bulk transport ;

14 temperature regulation ; [9 max]

(b) 15 idea of lipid > protein > carbohydrate / AW ; **A** lipid has more energy than either protein or carbohydrate

16 comparative figures ; e.g. 39.4, 17.0 and 15.8 *accept any two*

17 kJ g^{-1} / per unit mass ;

18 more hydrogen atoms in molecule, more energy ;

19 lipid have more, hydrogen atoms / C-H bonds ;

20 (most) energy comes from oxidation of hydrogen to water ;

21 using reduced, NAD / FAD ;

22 in ETC ;

23 detail of ETC ;

24 ATP production

[6 max]

[Total: 15]

9.

STEMUP.CO.UK

10 **(a)** 1 nucleotide ;

2 adenine + ribose / pentose + three phosphates ;

3 loss of phosphate leads to energy release / hydrolysis releases 30.5 kJ ;

4 ADP + Pi ⟷ ATP (reversible reaction) ;

5 small packets of energy ;

6 small / water soluble, so can move around cell ;

7 used by cells as immediate energy donor ;

8 link between energy yielding and energy requiring reactions / AW ;

9 high turnover ;

10 two examples of use ; ; e.g. active transport / muscle contraction / Calvin cycle /

11 protein synthesis [8 max]

(b) 12 Pyruvate, cannot enter mitochondrion / remains in the cytoplasm ;

13 becomes, hydrogen acceptor / reduced ;

14 by reduced NAD ;

15 from glycolysis ;

16 converted to lactate ;

17 lactate dehydrogenase ;

18 allows glycolysis to continue ;

19 no, decarboxylation / CO_2 removed ;

20 single step ;

21 reversible reaction / converted back to pyruvate ;

22 by oxidation ;

23 ref. oxygen debt ;

24 ethanol produced ;

accept ora for marking points 19–23 [7 max]

[Total: 15]

10.

STEMUP.CO.UK

11 (a) 1. multicellular ;

2. (cells are) differentiated into tissues ;

3. autotrophic / photosynthetic ;

4. eukaryotic (cells);

5. starch is storage compound ;

6. (some have) chloroplasts / chlorophyll ;

7. cell wall ;

8. made of cellulose ;

9. plasmodesmata ;

10. large (central) vacuole ; [max 7]

(b) 1. 0.5–1.0 μm, diameter / width ;

2. double membrane ;

3. inner membrane folded / cristae ;

4. hold, stalked particles / ATP synthase / ATP synthetase ;

5. site of ETC ;

6. ref. H^+ and intermembrane space ;

7. ATP production ;

8. oxidative phosphorylation / chemiosmosis ;

9. matrix is site of, link reaction / Krebs cycle ;

10. enzymes in matrix ;

11. 70S ribosomes ;

12. (mitochondrial) DNA ; [max 8]

[Total: 15]

STEMUP.CO.UK