CAMBRIDGE INTERNATIONAL AS & A LEVEL CHEMISTRY · 9701

CHEMISTRY



MARK SCHEME

2 ATOMS, MOLECULES AND STOICHIOMETRY

SCORE	PERCENTAGE
/	%

STEMUP

These questions belong to CIE, not to us.

The examination questions, mark schemes and examiner comments in this pack were written and published by Cambridge International (CIE). They are past assessment materials, reproduced here so that students can revise from them.

What we do not own

STEMUP is not affiliated with CIE and does not own, and makes no claim to own, any of the question, mark scheme or examiner report material in this pack. CIE has not endorsed, approved or been involved in the production of it. CIE and the qualification titles used here belong to their respective owners and appear for identification only.

Where to find the originals

CIE publishes its past papers, mark schemes and examiner reports. Always check the current syllabus at cambridgeinternational.org before you revise.

If you hold the rights to any material in this pack and would like it amended or removed, contact admin@stemup.co.uk and we will act on it.

Paper 1 Multiple Choice Answers: Atoms, Molecules and Stoichiometry





Question	Paper	Q. No.	Answer
50	9701_m21_qp_12	4	B
51	9701_w20_qp_11	1	C
52	9701_s21_qp_11	1	C
53	9701_s21_qp_12	1	C
54	9701_w21_qp_12	2	C
55	9701_s20_qp_11	11	C
56	9701_w19_qp_11	2	C
57	9701_s17_qp_11	3	A
58	9701_s17_qp_12	2	B
59	9701_m22_qp_12	3	C
60	9701_m22_qp_12	14	D
61	9701_s21_qp_13	1	A
62	9701_s19_qp_12	3	A
63	9701_m18_qp_12	1	D
64	9701_s18_qp_12	4	D
65	9701_m17_qp_12	2	B
66	9701_w17_qp_11	1	A
67	9701_w17_qp_12	2	C
68	9701_w15_qp_11	1	C
69	9701_m22_qp_12	20	B
70	9701_m22_qp_12	31	C
71	9701_m22_qp_12	32	A
72	9701_m21_qp_12	3	A
73	9701_s21_qp_11	6	B
74	9701_s21_qp_11	27	C
75	9701_s21_qp_13	2	D
76	9701_w21_qp_11	15	A
77	9701_w21_qp_11	28	C
78	9701_w21_qp_12	14	D
79	9701_m20_qp_12	7	B
80	9701_s20_qp_11	4	D
81	9701_s20_qp_11	13	D
82	9701_s20_qp_11	17	C
83	9701_s20_qp_12	2	C
84	9701_s20_qp_12	6	D
85	9701_s20_qp_12	13	D
86	9701_w20_qp_11	2	B
87	9701_w20_qp_12	2	A
88	9701_w20_qp_12	5	D
89	9701_m19_qp_12	2	A
90	9701_s19_qp_11	3	C
91	9701_w19_qp_12	2	A
92	9701_w19_qp_12	9	C
93	9701_m18_qp_12	2	A
94	9701_m18_qp_12	25	A
95	9701_s18_qp_13	5	C

Paper 1 Multiple Choice Answers: Atoms, Molecules and Stoichiometry

Question	Paper	Q. No.	Answer
96	9701_w18_qp_11	3	A
97	9701_s17_qp_12	19	B
98	9701_s17_qp_13	2	D
99	9701_s17_qp_13	3	B
100	9701_w17_qp_12	3	B
101	9701_s16_qp_11	3	C
102	9701_s16_qp_11	17	B
103	9701_s16_qp_12	4	C
104	9701_s16_qp_12	14	D
105	9701_s16_qp_13	3	B
106	9701_w16_qp_12	3	B
107	9701_w16_qp_12	4	D
108	9701_s15_qp_12	2	B
109	9701_w15_qp_11	3	B
110	9701_w15_qp_12	2	C
111	9701_w15_qp_12	13	B

Answers

1. 9701_s17_ms_21 Q: 1

(a)	The mass of a molecule OR the (weighted) average / (weighted) mean mass of the molecules	**1**
	Relative / compared to $\frac{1}{12}$ (the mass) of an atom of carbon–12 OR on a scale in which a carbon–12 atom / isotope has a mass of (exactly) 12 (units)	**1**
(b)(i)	**3**	**1**
(b)(ii)	**8**	**1**
(b)(iii)	$C_3H_8O + 4½O_2 \rightarrow 3CO_2 + 4H_2O$	**1**
(b)(iv)	 **AND** propan–2–ol / 2–propanol	**1**
	 **AND** propan–1–ol / 1–propanol	**1**
	Alternative answers (any two):  **AND** butan–1–ol / 1–butanol  **AND** butan–2–ol / 2–butanol  **AND** (2–)methylpropan–1–ol / (2–)methyl–1–propanol  **AND** (2–)methylpropan–2–ol / (2–)methyl–2–propanol	
(b)(v)	correct conversions of data to SI/consistent units $p = 100\,000$; $V = 20 \times 10^{-6}$; $T = 393$	**1**
	calculation of n ($= pV/RT$) from M1 values $n = \frac{100 \times 10^3 \times 20 \times 10^{-6}}{8.31 \times 393}$	**1**
	calculation of mass m ($= n \times Mr$) **AND** answer correct to **3sf** $m = 6.12 \times 10^{-4} \times 60 = 0.0367$ (g) Alternative answer for using $C_4H_{10}O$: $m = 6.12 \times 10^{-4} \times 74 = 0.0453$ (g)	**1**
	Total:	**10**

STEMUP.CO.UK

2. 9701_m16_ms_22 Q: 2

(a) (i)	$\frac{27.30}{1000} \times 0.020 = \mathbf{5.46 \times 10^{-4}}$ (mol)	[1]
(ii)	**(i)** $\times 6 = \mathbf{3.28 \times 10^{-3}}$ (mol)	[1]
(iii)	**(ii)** $\times \frac{250}{25.00} = \mathbf{3.28 \times 10^{-2}}$ (mol)	[1]
(iv)	M_r of $FeCO_3$ = 55.8 + 12.0 + 3(16.0) = **115.8** **(iii)** $\times M_r(FeCO_3)$ = **3.79** g	[1] [1]
(v)	$\frac{\textbf{(iv)}}{5.00} \times 100\% =$ **75.9%**	[1]
(b) (i)	$2Fe^{3+} + Sn^{2+} \rightarrow 2Fe^{2+} + Sn^{4+}$ species balancing	 [1] [1]
(ii)	**$SnCl_2$(aq)** + **2**$HgCl_2$(aq) → $SnCl_4$(**aq**) + Hg_2Cl_2(**s**) $SnCl_2$ AND 2 state symbols	 [1] [1]

3. 9701_w16_ms_21 Q: 1

(a)	6×10^{-3} (mol)	1	**1**
(b)	$NaOH + HCl \rightarrow NaCl + H_2O$	1	**1**
(c)	6×10^{-3} (mol)	1	**1**
(d)	4×10^{-3} (mol)	1	**1**
(e)	4×10^{-3} (mol)	1	**1**
(f)	1×10^{-3} (mol)	1	**1**
(g)	170	1	**1**
(h)	28(.0) Si/silicon	1 1	**2**
	Total:		**9**

4. 9701_w16_ms_22 Q: 1

(a)	0.04 **OR** 4×10^{-2}	**1**
(b)(i)	$Na_2CO_3 + 2HCl \rightarrow 2NaCl + CO_2 + H_2O$	**1**
(b)(ii)	0.00075 **OR** 7.5×10^{-4}	**1**
(b)(iii)	0.0015 **OR** 1.5×10^{-3}	**1**
(b)(iv)	0.015 **OR** 1.5×10^{-2}	**1**
(b)(v)	0.025 **OR** 2.5×10^{-2}	**1**
(b)(vi)	0.0125 **OR** 1.25×10^{-2} **OR** 0.013 **OR** 1.3×10^{-2}	**1**
(b)(vii)	40 Ca / calcium	**1** **1**
	Total:	**9**

Q1.

6 (a)

$\frac{66.7}{12}$	$\frac{11.1}{1}$	$\frac{22.2}{16}$	
= 5.5	= 11.1	= 1.3875	

Divide by 1.3875

C_4H_8O (1) 48 + 8 + 16 = 72 hence C_4H_8O (1) **[2]**

Q2.

2 **(a)** $1s^2$ $2s^2 2p^6$ $3s^2 3p^3$ [1]

(b) 5 or V [1]

(c) **(i)** $3NaOH + H_3PO_4 \longrightarrow Na_3PO_4 + 3H_2O$ [1]

(ii) (50 x 0.5) / 1000 = 0.025 (moles) [1]

(iii) conseq. on **(i)** 3 x .025 = 0.075 (moles) [1]

(d) **(i)** $P_4S_3 + 8O_2 \longrightarrow P_4O_{10} + 3SO_2$ (or $2P_2O_5$) balanced = 2 marks

OR $+ 6O_2 \longrightarrow P_4O_6 + 3SO_2$ (or $2P_2O_3$) unbalanced = 1 mark

[2]

(ii) $P_4O_{10} + 6H_2O \longrightarrow 4H_3PO_4$ [1]

OR $P_4O_6 + 6H_2O \longrightarrow 4H_3PO_3$

$SO_2 + H_2O \longrightarrow H_2SO_3$ [1]

(if SO_3 then e.c.f.) **Total = [9]**

Q3.

(c) **(i)** (1.2) / (2.4) = 1/20 or 0.05 moles [1]

(ii) $2NH_3 + H_2SO_4 \longrightarrow (NH_4)_2SO_4$ or equivalent [1]

(iii) 0.025 mols of H_2SO_4 are required

Vol. of 0.50 mol dm^{-3} H_2SO_4 required = (0.025 x 1000) / 0.5 = $50cm^3$ [1]

Q4.

STEMUP.CO.UK

(d) (i) $2H_2S + 3O_2 \rightarrow 2H_2O + 2SO_2$ **(1)**
from -2 **(1)** to +4 **(1)**
allow e.c.f. on equation

(ii) 68.2g H_2S react with 3 x 24 dm^3 O_2 **(1)**
8.65g H_2S react with $\frac{3 \times 24 \times 8.65}{68.2}$ = 9.13 dm^3 **(1)**
allow 9.16 dm^3 if H_2S = 68 is used
allow e.c.f on **(d)(i)** **[5]**

Q5.

1 **(a) (i)** ammonia/NH_3 **(1)**

(ii) NH_4^+ **(1)**

(iii) iron(II) hydroxide/$Fe(OH)_2$ **(1)** **[3]**

(b) barium sulphate/$BaSO_4$ **(1)** **[1]**

(c) (i) $FeSO_4$ **(1)**

$(NH_4)_2SO_4$ **(1)**

(ii) $FeSO_4$ = 151.9 (allow 151.8 to 152) **(1)**

$(NH_4)_2SO_4$ = 132.1 (allow 132) **(1)**

(iii) $xH_2O = 392 - (132.1 + 151.9) = 108$ **(1)**

$x = \frac{108}{18} = 6$ **(1)**

allow e.c.f. on candidate's sulphate in **(c)(i)**
e.c.f. answer must be a whole number **[6]**

[Total: 10]

Q6.

(c) (i) $CaC_2 + 2H_2O \rightarrow Ca(OH)_2 + C_2H_2$ **(1)**

(ii) $n(C_2H_2) = n(CaC_2) = 100 \times 46.5$ **(1)**

mass of CaC_2 = 100 x 46.5 x 64 =

= 297 570 g

= 297.6 kg (accept 298 kg)
correct units necessary **(1)**

allow e.c.f. on candidate's answer in **(b)** **[3]**

Q7.

STEMUP.CO.UK

(b) C : H : O – $\frac{40}{2} : \frac{6.7}{1} : \frac{53.3}{16}$ [1]

= 3.33 : 6.7 : 3.33 [1]

= 1 : 2 : 1 **[2]**

Q8.

(d) **(i)** n(Ti) = $\frac{0.72}{47.9}$ = 0.015 (1)

(ii) n(Cl) = $\frac{(2.85 - 0.72)}{35.5}$ = 0.06 (1)

(iii) 0.015 : 0.06 = 1:4
empirical formula of **A** is $TiCl_4$
Allow ecf on answers to **(i)** and/or **(ii)**. (1)

(iv) $Ti + 2Cl_2 \rightarrow TiCl_4$ (1)
Allow ecf on answers to **(iii)**. [4]

Q9.

4 **(a)** C : H : O = $\frac{48.7}{12} : \frac{8.1}{1} : \frac{43.2}{16}$ (1)
= 4.06 : 8.1 : 2.70
= 1.5 : 3 : 1
= 3 : 6 : 2
empirical formula is $C_3H_6O_2$ (1) [2]

(b) **(i)** $M_r = \frac{mRT}{pV} = \frac{0.13 \times 8.31 \times 400}{1.00 \times 10^5 \times 58.0 \times 10^{-6}}$ (1)

= 74.5 (1)

(ii) $C_3H_6O_2$ = 36 + 6 + 32 = 74 (1)
$n(C_3H_6O_2)$ = 74.5
hence molecular formula of **E** is $C_3H_6O_2$ (1) [4]

Q10.

(d) **(i)** C : H : O = $\frac{35.8}{12} : \frac{4.5}{1} : \frac{59.7}{16}$ this mark is for correct use of A_r values (1)
C : H : O = 2.98 : 4.5 : 3.73
C : H : O = 1 : 1.5 : 1.25 this mark is for evidence of correct calculation (1)
gives empirical formula of **W** is $C_4H_6O_5$

(ii) $C_4H_6O_5$ = 12 × 4 + 1 × 6 + 16 × 5 = 134
molecular formula of **W** is $C_4H_6O_5$ (1) [3]

STEMUP.CO.UK

Q11.

(c) (i) $n(Li) = \frac{0.83}{6.9} = 0.12$ (1)

(ii) 2 mol Li $\rightarrow$ 1 mol H_2

0.12 mol Li $\rightarrow \frac{1 \times 0.12}{2} = 0.06$ mol H_2 (1)

volume of $H_2 = 0.06 \times 24.0 = 1.44 dm^3$ (1)

(iii) 2 mol Li $\rightarrow$ 2 mol LiOH

0.12 mol Li $\rightarrow$ 0.12 mol LiOH in 0.50 dm^3 (1)

[LiOH] $= \frac{0.12 \times 1}{0.50} = 0.24$ mol dm^{-3} (1) [5]

Q12.

(c) (i) mass of $C_{14}H_{30}$ burnt

$\frac{8195 \times 10.8}{1000} = 88.506 = 88.5$ t (1)

(ii) mass of CO_2 produced

M_r of $C_{14}H_{30} = (14 \times 12 + 30 \times 1) = 198$ (1)

2 x 198 t of $C_{14}H_{30} \rightarrow 28 \times 44$ t of CO_2

88.5 t of $C_{14}H_{30} \rightarrow \frac{28 \times 44 \times 88.5}{2 \times 198}$ (1)

= 275.3 t of CO_2 (1)

allow 275.4 t if candidate has used 88.506
allow ecf on wrong value for M_r of $C_{14}H_{30}$ [4]

Q13.

(b) (i) $n(NaOH) = \frac{22.5 \times 2.00}{1000} = 0.045$ (1)

(ii) $n(NaOH) = n(HCl) = 0.005$ (1)

(iii) $CH_3CO_2H + NaOH \rightarrow CH_3CO_2Na + H_2O$ (1)

(iv) $n(NaOH) = 0.045 - 0.005 = 0.04$
allow ecf on **(i)** and/or **(ii)** (1) [4]

Q14.

STEMUP.CO.UK

1 Throughout this question, deduct **one mark only** for sig. fig. error.

(a) (i) the volume of solution **A** present in one 'typical ant' is
$7.5 \times 10^{-6} \times 1000 = 7.5 \times 10^{-3}$ cm^3 (1)

(ii) the volume of pure methanoic acid in one 'typical ant' is
$7.5 \times 10^{-3} \times \frac{50}{100} = 3.75 \times 10^{-3}$ gives 3.8×10^{-3} cm^3

allow ecf on **(i)** (1)

(iii) no. of ants = $\frac{1000}{3.8 \times 10^{-3}} = 263157.8947$ gives 2.6×10^5

use of 3.75×10^{-3} gives $266666.6667 = 2.7 \times 10^5$ (1) [3]

(b) (i) the volume of solution **A**, in one ant bite is
$\frac{80}{100} \times 7.5 \times 10^{-3} = 6.0 \times 10^{-3}$ cm^3

allow ecf on **(a)(i)** (1)

the volume of pure methanoic acid in one bite is
$\frac{50}{100} \times 6.0 \times 10^{-3} = 3.0 \times 10^{-3}$ cm^3

allow ecf on first part of **(b)(i)** (1)

(ii) the mass of methanoic acid in one bite is
$3.0 \times 10^{-3} \times 1.2 = 3.6 \times 10^{-3}$ g

allow ecf on **(b)(i)** (1) [3]

(c) (i) $HCO_2H + NaHCO_3 \rightarrow HCO_2Na + H_2O + CO_2$ (1)

(ii) 46 g $HCO_2H \equiv$ 84 g $NaHCO_3$ (1)

5.4×10^{-3} g $HCO_2H \equiv \frac{84 \times 5.4 \times 10^{-3}}{46}$ g $NaHCO_3$
$= 9.860869565 \times 10^{-3}$
$= 9.9 \times 10^{-3}$ g $NaHCO_3$ (1) [3]

[Total: 9]

Q15.

STEMUP.CO.UK

2 **(a)** $(NH_4)_2SO_4 + 2NaOH \rightarrow 2NH_3 + \mathbf{Na_2SO_4} + 2H_2O$

correct products (1)

correctly balanced equation (1) [2]

(b) (i) $NaOH + HCl \rightarrow NaCl + H_2O$ (1)

(ii) $n(HCl) = \frac{31.2}{1000} \times 1.00 = 0.0312 = 0.03$ (1)

(iii) $n(NaOH) = \frac{50.0}{1000} \times 2.00 = 0.10$ (1)

(iv) $n(NaOH)$ used up $= 0.10 - 0.0312 = 0.0688 = 0.07$ (1)

(v) $n[(NH_4)_2SO_4] = \frac{0.0688}{2} = 0.0344 = 0.03$ (1)

(vi) mass of $(NH_4)_2SO_4 = 0.0344 \times 132 = 4.5408 = 4.54$ (1)

(vii) percentage purity $= \frac{4.5408 \times 100}{5.00} = 90.816 = 90.8$ (1) [7]

[Total: 9]

Q16.

(e) $C : H : O = \frac{37.5}{12} : \frac{4.17}{1} : \frac{58.3}{16}$

$= 3.13 : 4.17 : 3.64$ (1)

$= 1 : 1.33 : 1.16$ (1)

$= 6 : 8 : 7$

empirical formula is $C_6H_8O_7$ (1) [3]

[Total: 19]

Q17.

2 **(a) (i)** $Na_2CO_3 + 2HCl \rightarrow 2NaCl + H_2O + CO_2$ (1)

(ii) $n(HCl) = \frac{35.8}{1000} \times 0.100 = 3.58 \times 10^{-3}$ (1)

(iii) $n(Na_2CO_3) = \frac{35.8}{2} \times 10^{-3} = 1.79 \times 10^{-3}$ mol in 25.0 cm^3 (1)

(iv) $n(Na_2CO_3) = 1.79 \times 10^{-3} \times 10 = 1.79 \times 10^{-2}$ mol in 250 cm^3 (1)

(v) mass of $Na_2CO_3 = 1.79 \times 10^{-2} \times 106 = 1.90$g

M_r of $Na_2CO_3 = 106$ (1)

mass of $Na_2CO_3 = 1.90$ g (1) [6]

STEMUP.CO.UK

(b) $n(H_2O)$ in 5.13 g of washing soda $= \frac{5.13 - 1.90}{18} = 1.79 \times 10^{-1}$ mol (1)

$n(Na_2CO_3)$ in 5.13 g of washing soda $= 1.79 \times 10^{-2}$ mol
$n(H_2O) : n(Na_2CO_3) = 10 : 1$ (1)

or
1.90 g Na_2CO_3 are combined with 3.23 g H_2O
106 g Na_2CO_3 are combined with $\frac{3.23 \times 106}{1.90} = 180.2$ g H_2 (1)
this is 10 mol of H_2O (1)

or
1.79×10^{-2} mol $Na_2CO_3.xH_2O \equiv 5.13$ g of washing soda
1 mol $Na_2CO_3.xH_2O \equiv \frac{5.13}{1.79 \times 10^{-2}} = 286.6$ g (1)
$Na_2CO_3 = 106$ and $H_2O = 18$ hence $x = 10$ (1) [2]

[Total: 8]

Q18.

1 **(a) (i)** $NaOH + HCl \rightarrow NaCl + H_2O$ (1)

$(NH_4)_2SO_4 + 2NaOH \rightarrow 2NH_3 + Na_2SO_4 + 2H_2O$ (1)

allow ionic equations in each case

(ii) $n(NaOH) = n(HCl) = \frac{39.2 \times 2.00}{1000} = 0.0784$ (1)

(iii) $n(NaOH) = n(HCl) = \frac{29.5 \times 2.00}{1000} = 0.059$ (1)

(iv) $n(NaOH) = 0.0784 - 0.059 = 0.0194$ (1)

(v) $n[(NH_4)_2SO_4] = \frac{0.0194}{2} = 9.7 \times 10^{-3}$ (1)

(vi) mass of $(NH_4)_2SO_4 = 9.7 \times 10^{-3} \times 132.1 = 1.2814$ g (1)

(vii) % of $(NH_4)_2SO_4 = \frac{1.2814 \times 100}{2.96} = 43.30405405 = 43.3$
give one mark for the correct expression (1)
give one mark for answer given as 43.3 – i.e. to 3 sig. fig. (1)
allow ecf where appropriate [9]

STEMUP.CO.UK

(b) fertiliser in the river causes
excessive growth of aquatic plants/algae **or** algal bloom (1)
when plants and algae die O_2 is used up **or** fish or aquatic life die (1) [2]

(c) manufacture of HNO_3 **or** explosives **or** nylon **or**
as a cleaning agent **or** as a refrigerant
not detergent (1) [1]

[Total:12]

Q19.

2 **(a)** **(i)** $n(H_2SO_4) = \frac{25.0 \times 1.00}{1000} - 0.025$ mol (1)

(ii) $n(NaOH) = \frac{16.2 \times 2.00}{1000} - 0.0324$ mol (1)

(iii) $n(H_2SO_4)$ reacting with NaOH $= \frac{0.0324}{2} - 0.0162$ mol (1)

(iv) $n(H_2SO_4)$ reacting with NH_3 = 0.025 - 0.0162 = 0.0088 mol (1)

(v) $n(NH_3)$ reacting with H_2SO_4 = 2 x 0.0088 = 0.0176 mol (1)

(vi) $n(NaNO_3)$ reacting = $n(NH_3)$ produced = 0.0176 mol (1)

(vii) mass of $NaNO_3$ that reacted = 0.0176 x 85 = 1.496 g (1)

(viii) % of $NaNO_3 = \frac{1.496 \times 100}{1.64} - 91.2195122 - 91.2$

give one mark for the correct expression (1)
give one mark for answer given as 91.2 – i.e to 3 sig. fig. (1)
allow ecf where appropriate

[9]

(b) $NaNO_3$ +5 **and** NH_3 -3 **both** required (1) [1]

[Total: 10]

Q20.

STEMUP.CO.UK

1 **(a)** **(i)** substance that speeds up a chemical reaction (1)
by lowering E_a
or by providing an alternative reaction pathway
or without being used up in the process (1)

(ii) $2H_2O_2 \rightarrow 2H_2O + O_2$ (1) [3]

(b) **(i)** alkanes **or** paraffins (1)

(ii) $2H_2O_2 : O_2$ and $C_{15}H_{32} : 23O_2$ (1)
whence $C_{15}H_{32} : 46H_2O_2$ (1)
allow e.c.f. on **(a)(ii)** [3]

(c) **(i)** $C_{15}H_{32} = 212$ (1)

$$n(C_{15}H_{32}) = \frac{212 \times 10^6}{212} = 1 \times 10^6 \text{ mol}$$

allow e.c.f. on wrong M_r of $C_{15}H_{32}$ (1)

(ii) $n(H_2O_2)$ required $= 46 \times 10^6$ mol (1)
mass of $H_2O_2 = 34 \times 46 \times 10^6$ g $= 1564$ tonnes
final answer must be in tonnes (1)
allow e.c.f. on **(b)(ii)** and **(c)(i)** [4]

(d) they would dissolve (1) [1]

[Total: 11]

Q21.

1 **(a)** the actual number of atoms of each element present (1)

in one molecule of a compound (1) [2]

(b) $C_xH_y + \left(x + \frac{y}{4}\right)O_2 \longrightarrow xCO_2 + \frac{y}{2}H_2O$

xCO_2 (1)

$\frac{y}{2}H_2O$ (1) [2]

(c) **(i)** oxygen/O_2 (1)

(ii) carbon dioxide/CO_2 (1)

(iii) 10 cm^3 (1)

(iv) 20 cm^3 (1) [4]

STEMUP.CO.UK

(d) $C_xH_y + \left(x + \frac{y}{4}\right)O_2 \longrightarrow xCO_2 + \frac{y}{2}H_2O$

$10\ cm^3 \qquad 20\ cm^3 \qquad 10\ cm^3$

1 mol of C_xH_y gives 1 mol of CO_2

whence $x = 1$ (1)

1 mol of C_xH_y reacts with 2 mol of O_2

whence $\left(x + \frac{y}{4}\right) = 2$

and $y = 4$ (1)

molecular formula is CH_4 (1) [3]

[Total: 11]

Q22.

STEMUP.CO.UK

1 (a) (i) mass of C = $\frac{12 \times 0.352}{44}$ = 0.096g (1)

n(C) = $\frac{0.096}{12}$ = 0.008 (1)

(ii) mass of H = $\frac{2 \times 0.144}{18}$ = 0.016g (1)

n(H) = $\frac{0.016}{1}$ = 0.016 (1)

(iii) mass of oxygen = 0.240 – (0.096 + 0.016) = 0.128g (1)

n(O) = $\frac{0.128}{16}$ = 0.008 (1)

allow ecf at any stage [6]

(b) C : H : O = 0.008: 0.016 : 0.008 = 1:2:1

allow C : H : O = $\frac{0.096}{12}$: $\frac{0.016}{1}$: $\frac{0.128}{16}$ = 1:2:1

gives CH_2O (1) [1]

(c) (i) $M_r = \frac{mRT}{pV} = \frac{0.148 \times 8.31 \times 333}{1.01 \times 10^5 \times 67.7 \times 10^{-6}}$ (1)

= 59.89

allow 59.9 or 60 (1)

(ii) $C_2H_4O_2$ (1) [3]

Q23.

2 (a) (i) mass of C = $\frac{12 \times 1.32}{44}$ = 0.36g (1)

n(C) = $\frac{0.36}{12}$ = 0.03 (1)

(ii) mass of H = $\frac{2 \times 0.54}{18}$ = 0.06 g (1)

n(H) = $\frac{0.06}{1}$ = 0.06 (1)

(iii) yes **because** 0.03 mol of C are combined with 0.06 mol of H **or**
C : H ratio is 1 : 2 **or**
empirical formula is CH_2 (1) [5]

STEMUP.CO.UK

(b) (i) C : H : O = $\frac{64.86}{12} : \frac{13.50}{1} : \frac{21.64}{16}$ (1)

= 5.41: 13.50 : 1.35

= 4 : 10 : 1

gives $C_4H_{10}O$ (1)

Q24.

1 **(a)** $ZnCO_3$ $Zn(OH)_2$ ZnO
not Zn **or** other compounds of Zn (any 2) [2]

(b) (i) to ensure all of the water of crystallisation had been driven off **or** to be at constant mass (1)

(ii) mass of $ZnSO_4$ = 76.34 – 74.25 = 2.09 g (1)

M_r $ZnSO_4$ = 65.4 + 32.1 + (4 × 16.0) = 161.5

allow use of Zn = 65 and/or S = 32 to give values between 161 and 161.5 (1)

$n(ZnSO_4) = \frac{2.09}{161.5} = 0.01294 = 1.29 \times 10^{-2}$

$ZnSO_4$ = 161 gives 1.30×10^{-2} (1)

(iii) mass of H_2O driven off = 77.97 – 76.34 = 1.63 g (1)

$n(H_2O) = \frac{1.63}{18} = 0.0905 = 9.1 \times 10^{-2}$ (1)

STEMUP.CO.UK

(iv) 1.29×10^{-2} mol $ZnSO_4$ are combined with 9.1×10^{-2} mol H_2O

1 mol $ZnSO_4$ is combined with $\frac{9.1 \times 10^{-2}}{1.29 \times 10^{-2}}$

$= 7.054 \equiv 7$ mol H_2O

answer must be expressed as a whole number
allow ecf on candidate's answers to **(b)(ii)** and **(b)(iii)** (1) [7]

(c) (i) $n(Zn) = n\ (CH_3CO_2)_2Zn.2H_2O$ (1)

$n(Zn) = \frac{0.015}{65.4} = 2.290 \times 10^{-4}$

$= 2.29 \times 10^{-4}$ (1)

mass of crystals $= 2.29 \times 10^{-4} \times 219.4 = 0.0502655$ g
$= 0.05$ g $= 50$ mg (1)

(ii) concentration of $(CH_3CO_2)_2Zn.2H_2O = \frac{2.29 \times 10^{-4}}{0.005} = 0.0458$

$= 4.58 \times 10^{-2}$ mol dm^{-3} (1)

allow correct answers if Zn = 65 is used [4]

[Total: 13]

Q25.

1 In this question, numerical answers should be given to three significant figures.

(a) (i) $C_6H_{12}O_6 + 6O_2 \rightarrow 6CO_2 + 6H_2O$ (1)

(ii) $M_r\ C_6H_{12}O_6 = 180$ (1)
180 g $C_6H_{12}O_6 \rightarrow 6$ mol CO_2

1200 g $C_6H_{12}O_6 \rightarrow \frac{6 \times 200}{180}$ mol CO_2

$= 40.0$ mol to 3 sf

allow ecf on wrong equation and/or wrong M_r (1)

(iii) 6.82×10^9 people will produce $6.82 \times 10^9 \times 40.0$ mol CO_2

$= 2.728 \times 10^{11}$ mol CO_2 (1)

2.728×10^{11} mol $CO_2 \equiv 2.728 \times 10^{11} \times 44 = 1.20032 \times 10^{13}$ g

$= 1.20 \times 10^7$ tonnes CO_2 to 3 sf (1) [5]

allow ecf on answer from **(ii)**

STEMUP.CO.UK

(b) (i) $2C_8H_{18} + 25O_2 \rightarrow 16CO_2 + 18H_2O$ **or**

$C_8H_{18} + 12½O_2 \rightarrow 8CO_2 + 9H_2O$ (1)

(ii) $M_r\ C_8H_{18} = (8 \times 12) + (18 \times 1) = 114$ (1)

mass of 4.00 dm^3 of octane = $4000 \times 0.70 = 2800$ g (1)

$n(C_8H_{18}) = \frac{2800}{114} = 24.56140351$ mol in 4.00 dm^3

= 24.6 mol to 3 sf (1)

(iii) 2 mol C_8H_{18} produce 16×44 g CO_2

24.6 mol C_8H_{18} produce $\frac{16 \times 44 \times 24.6}{2}$ g CO2

= 8659.2 g CO_2

= 8660 g CO_2 to 3 sf (1) [5]

(c) 6.82×10^9 people produce 1.20×10^7 tonnes CO_2 per day

8660 g CO_2 produced when car travels 100 km

when travelling 1 km, car produces $\frac{8660}{100} = 8.66 \times 10^{-1}$ g

$= 8.66 \times 10^{-5}$ tonnes (1)

to produce 1.20×10^7 tonnes CO_2 car must travel

$\frac{1.20 \times 10^7}{8.66 \times 10^{-5}}$

$= 1.385681293 \times 10^{11} = 1.39 \times 10^{11}$ km to 3 sf (1) [2]

(d) possible pollutants and the damage they cause

CO	NO_x		SO_2	H_2O	C	unburned C_8H_{18}
	NO	NO2				
toxic	toxic	toxic	toxic			
	global warming	respiratory problems	respiratory problems	global warming	respiratory problems	respiratory problems
	photochemical smog	acid rain	acid rain			

compound (1)
damage (1) [2]

[Total: 14]

Q26.

STEMUP.CO.UK

4 (a) (i) $H_2X + 2NaOH \rightarrow Na_2X + 2H_2O$ (1)

(ii) $n(OH^-) = \frac{21.6 \times 0.100}{1000} = 2.16 \times 10^{-3}$ mol (1)

(iii) $n(\mathbf{R}) = n(H_2X) = \frac{2.16 \times 10^{-3}}{2}$

$= 1.08 \times 10^{-3}$ mol in 25.0 cm^3 (1)

(iv) $n(\mathbf{R}) = 1.08 \times 10^{-3} \times \frac{250}{25.0} = 0.0108$ mol in 250 cm^3 (1)

(v) 0.0108 mol of **R** = 1.25 g of **R**

1 mol of **R** $= \frac{1.25 \times 1}{0.0108} = 115.7 = 116$ g (1) [5]

(b) (i) M_r of **S** = 116
M_r of **T** = 134
M_r of **U** = 150 **all three** needed (1)

(ii) **S** (1) [2]

STEMUP.CO.UK