

CHEMISTRY



EXAM QUESTION PACK

6 ELECTROCHEMISTRY

SCORE

/

PERCENTAGE

%

STEMUP

These questions belong to CIE, not to us.

The examination questions, mark schemes and examiner comments in this pack were written and published by Cambridge International (CIE). They are past assessment materials, reproduced here so that students can revise from them.

What we do not own

STEMUP is not affiliated with CIE and does not own, and makes no claim to own, any of the question, mark scheme or examiner report material in this pack. CIE has not endorsed, approved or been involved in the production of it. CIE and the qualification titles used here belong to their respective owners and appear for identification only.

Where to find the originals

CIE publishes its past papers, mark schemes and examiner reports. Always check the current syllabus at [cambridgeinternational.org](https://www.cambridgeinternational.org) before you revise.

If you hold the rights to any material in this pack and would like it amended or removed, contact admin@stemup.co.uk and we will act on it.



Chapter 6

Electrochemistry


6.1 Redox processes: electron transfer and changes in oxidation number (oxidation state)

281. 9701_m22_qp_12 Q: 10

Two half-equations are shown.



The equation for the reaction between manganate(VII) ions and sulfite ions is shown.




Which statements are correct?

- 1 $u = x = z$
- 2 Manganese is reduced to oxidation state +4.
- 3 Sulfur is oxidised from oxidation state +4 to +6.

A 1, 2 and 3 **B** 1 and 2 only **C** 1 and 3 only **D** 2 and 3 only

282. 9701_m21_qp_12 Q: 8

VO_2Cl reacts with NaI under acidic conditions.




The oxidation state of Cl is -1 in VO_2Cl and in VOCl_2 .

Which row about this reaction is correct?

	vanadium	iodine
A	is oxidised	is oxidised
B	is oxidised	is reduced
C	is reduced	is oxidised
D	is reduced	is reduced

283. 9701_s21_qp_11 Q: 9

When hydrogen iodide is reacted with concentrated sulfuric acid, several reactions occur, including:




Which row gives the change in oxidation number of iodine and of sulfur in this reaction?

	change in oxidation number of iodine	change in oxidation number of sulfur
A	-1	$+6$
B	-1	$+8$
C	$+1$	-6
D	$+1$	-8

284. 9701_s21_qp_12 Q: 9

The equation for a redox reaction is shown.



Which species is being oxidised in this reaction?

- A** Sn^{2+} **B** Cl^- **C** Hg^+ **D** Hg^{2+}

6.1. REDOX PROCESSES: ELECTRON TRANSFER AND CHANGES IN OXIDATION NUMBER (OXIDATION STATE)

285. 9701_w21_qp_11 Q: 9

Zinc atoms can be oxidised to Zn^{2+} ions by dichromate(VI) ions in acid solution. Chromium is reduced to Cr^{3+} in this reaction.

Which equation is correct?

- A** $\text{Cr}_2\text{O}_7^{2-} + \text{Zn} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + \text{Zn}^{2+} + 7\text{H}_2\text{O}$
- B** $\text{Cr}_2\text{O}_7^{2-} + \text{Zn} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 3\text{Zn}^{2+} + 7\text{H}_2\text{O}$
- C** $\text{Cr}_2\text{O}_7^{2-} + 3\text{Zn} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 3\text{Zn}^{2+} + 7\text{H}_2\text{O}$
- D** $2\text{Cr}_2\text{O}_7^{2-} + 3\text{Zn} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 3\text{Zn}^{2+} + 7\text{H}_2\text{O}$

286. 9701_w21_qp_12 Q: 10


In a catalytic converter in the exhaust system of a car, carbon monoxide is oxidised to carbon dioxide, and nitrogen monoxide is reduced to nitrogen.

What are the changes in oxidation number of carbon and nitrogen in these two processes?

	carbon	nitrogen
A	−2	+2
B	−1	+1
C	+1	−1
D	+2	−2

287. 9701_m20_qp_12 Q: 5

In the redox reaction shown, how do the oxidation states of vanadium and sulfur change?



	vanadium		sulfur	
	from	to	from	to
A	+1	+3	0	−2
B	+1	+3	+4	+6
C	+5	+3	0	−2
D	+5	+3	+4	+6

288. 9701_m20_qp_12 Q: 10

When the equation is correctly balanced, what is the value of c ?



- A** 3 **B** 4 **C** 5 **D** 6

289. 9701_s20_qp_11 Q: 2

Cobalt can form the positive ion $\text{Co}(\text{NH}_3)_4\text{Cl}_2^+$.


What is the oxidation number of cobalt in this ion?

- A** +1 **B** +2 **C** +3 **D** +6
-

290. 9701_s20_qp_11 Q: 8

In this question you should use changes in oxidation numbers to balance a chemical equation.

Acidified potassium dichromate(VI) solution can oxidise a solution of V^{2+} ions. The equation for this reaction is shown.



What is the ratio $a : b$ in the correctly balanced equation?

- A** 1 : 1 **B** 1 : 2 **C** 2 : 1 **D** 4 : 1
-

291. 9701_s20_qp_12 Q: 5

In this question you should use changes in oxidation numbers to balance a chemical equation.

The following reaction occurs when MnO_2 is warmed with dilute H_2SO_4 .



What is the ratio of $c : d$ in the correctly balanced equation?

- A** 1 : 1 **B** 1 : 2 **C** 2 : 3 **D** 3 : 2
-

292. 9701_s20_qp_12 Q: 10

In which reaction does an element undergo the largest change in oxidation number?

- A** $\text{Cl}_2 + 2\text{OH}^- \rightarrow \text{OCl}^- + \text{Cl}^- + \text{H}_2\text{O}$
B $3\text{Cl}_2 + 6\text{OH}^- \rightarrow \text{ClO}_3^- + 5\text{Cl}^- + 3\text{H}_2\text{O}$
C $\text{Cr}_2\text{O}_7^{2-} + 6\text{Fe}^{2+} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 6\text{Fe}^{3+} + 7\text{H}_2\text{O}$
D $3\text{MnO}_4^{2-} + 4\text{H}^+ \rightarrow \text{MnO}_2 + 2\text{MnO}_4^- + 2\text{H}_2\text{O}$
-

293. 9701_s20_qp_13 Q: 10

When solid KClO_3 is heated in the absence of air, a mixture of two chlorine compounds in the mole ratio of 3 : 1 is formed. Chlorine is the only element whose oxidation number changes in this reaction.

What could be the oxidation numbers of chlorine in the two compounds that are formed?

- A** +3 and -1 **B** +6 and +4 **C** +7 and -1 **D** +7 and +1
-


294. 9701_w20_qp_11 Q: 8

In which reaction does the greatest change in the oxidation number of sulfur occur?

- A** $\text{S(s)} + \text{O}_2\text{(g)} \rightarrow \text{SO}_2\text{(g)}$
- B** $\text{SO}_2\text{(g)} + \frac{1}{2}\text{O}_2\text{(g)} \rightleftharpoons \text{SO}_3\text{(g)}$
- C** $\text{SO}_3\text{(g)} + \text{H}_2\text{SO}_4\text{(l)} \rightarrow \text{H}_2\text{S}_2\text{O}_7\text{(l)}$
- D** $\text{H}_2\text{S}_2\text{O}_7\text{(l)} + \text{H}_2\text{O(l)} \rightarrow 2\text{H}_2\text{SO}_4\text{(l)}$

295. 9701_w20_qp_11 Q: 9

The first stage in the chloride process for the manufacture of titanium consists of the following reaction.



What is reduced in this reaction?


- A** carbon
- B** chlorine
- C** oxygen
- D** titanium

296. 9701_w20_qp_12 Q: 1

What is the average oxidation number of sulfur in each compound?

	$\text{Ca}(\text{HSO}_3)_2$	$\text{Na}_2\text{S}_2\text{O}_3$
A	4	2
B	4	4
C	6	2
D	6	4

297. 9701_w20_qp_12 Q: 9

Chlorine dioxide, ClO_2 , reacts with sodium hydroxide in the reaction shown.




Which statement correctly describes this redox reaction?

- A** Chlorine atoms are oxidised and oxygen atoms are reduced.
- B** Chlorine atoms are reduced and oxygen atoms are oxidised.
- C** Some chlorine atoms are oxidised and some chlorine atoms are reduced.
- D** Some oxygen atoms are oxidised and some oxygen atoms are reduced.

298. 9701_m19_qp_12 Q: 10

Acidified potassium manganate(VII) reacts with iron(II) ethanedioate, FeC_2O_4 .

The reactions taking place are shown.



How many moles of iron(II) ethanedioate react with **one** mole of potassium manganate(VII)?

- A** 0.60 **B** 1.67 **C** 2.50 **D** 5.00
-

299. 9701_s19_qp_11 Q: 9

Ethanedioic acid, $\text{HO}_2\text{CCO}_2\text{H}$, can be oxidised by KMnO_4 in dilute sulfuric acid. The products of this reaction are carbon dioxide, water, potassium sulfate and manganese(II) sulfate.

In this reaction each ethanedioic acid molecule loses two electrons as it is oxidised. A half-equation for this process is shown.



How many water molecules are produced when five ethanedioic acid molecules are oxidised by KMnO_4 in dilute sulfuric acid?

- A** 5 **B** 8 **C** 10 **D** 16
-

300. 9701_s19_qp_12 Q: 9

X is either chlorine or an oxide of chlorine.

X reacts with water, under suitable conditions, to form the two acids HCl and HClO_3 in the mole ratio of 1 (HCl): 5 (HClO_3).

What could be X?

- A** Cl_2 **B** Cl_2O **C** ClO_2 **D** Cl_2O_7
-

6.1. REDOX PROCESSES: ELECTRON TRANSFER AND CHANGES IN OXIDATION NUMBER (OXIDATION STATES)

301. 9701_s19_qp_13 Q: 1

Manganese and nitrogen can show a range of different oxidation states.

Calculate the sum of the oxidation states of Mn and N in each row of the table.

In which row is this sum the smallest?

	manganese-containing species	nitrogen-containing species
A	MnCl_4	N_2
B	MnCO_3	NO_2^-
C	K_2MnO_4	NH_4^+
D	$\text{Mn}(\text{OH})_3$	NH_2OH

302. 9701_s19_qp_13 Q: 8

Ethanol can be oxidised to ethanal by dilute acidified dichromate(VI) ions.

The oxidation reaction equation is $\text{C}_2\text{H}_5\text{OH} \rightarrow \text{C}_2\text{H}_4\text{O} + 2\text{H}^+ + 2\text{e}^-$.

The reduction reaction equation is $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^- \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$.

Which equation is correct?

- A** $\text{Cr}_2\text{O}_7^{2-} + 8\text{H}^+ + 3\text{C}_2\text{H}_5\text{OH} \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O} + 3\text{C}_2\text{H}_4\text{O}$
- B** $\text{Cr}_2\text{O}_7^{2-} + 12\text{H}^+ + \text{C}_2\text{H}_5\text{OH} \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O} + \text{C}_2\text{H}_4\text{O}$
- C** $\text{Cr}_2\text{O}_7^{2-} + 12\text{H}^+ + 3\text{C}_2\text{H}_5\text{OH} \rightarrow 2\text{Cr}^{3+} + 6\text{H}_2\text{O} + 3\text{C}_2\text{H}_4\text{O}$
- D** $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 3\text{C}_2\text{H}_5\text{OH} \rightarrow 2\text{Cr}^{3+} + 6\text{H}_2\text{O} + 3\text{C}_2\text{H}_4\text{O}$


303. 9701_w19_qp_11 Q: 6

What is the oxidation number of sulfur in each species?

	H_2S	SO_2	H_2SO_3
A	-2	+4	+4
B	-2	+4	+6
C	+2	-4	+4
D	+2	-4	+6

304. 9701_w19_qp_11 Q: 9

In the chemical equation, w, x, y and z are all whole numbers.




When the equation is balanced, what are w, x and y?

	w	x	y
A	1	1	2
B	2	2	2
C	2	3	8
D	3	2	2

305. 9701_w19_qp_12 Q: 8

$(\text{NH}_4)_2\text{Cr}_2\text{O}_7$ decomposes when heated.




Which element is oxidised and which element is reduced?

	oxidised	reduced
A	chromium	nitrogen
B	hydrogen	chromium
C	nitrogen	chromium
D	nitrogen	hydrogen

306. 9701_w19_qp_12 Q: 15

Chlorate(V) ions, ClO_3^- , are produced in the redox reaction between chlorine and hot aqueous sodium hydroxide. Oxidation numbers can be used to help balance the equation for this reaction.



What are the values of v, x and y in the balanced equation?

	v	x	y
A	2	3	1
B	3	4	2
C	3	5	1
D	7	12	2

307. 9701_m18_qp_12 Q: 10

Which reaction is **not** a redox reaction?

- A** $\text{Mg} + 2\text{HNO}_3 \rightarrow \text{Mg}(\text{NO}_3)_2 + \text{H}_2$
B $2\text{Mg}(\text{NO}_3)_2 \rightarrow 2\text{MgO} + 4\text{NO}_2 + \text{O}_2$
C $\text{SO}_2 + \text{NO}_2 \rightarrow \text{SO}_3 + \text{NO}$
D $\text{SO}_3 + \text{H}_2\text{O} \rightarrow \text{H}_2\text{SO}_4$
-

308. 9701_m18_qp_12 Q: 11

The reaction between sulfur dioxide and oxygen is reversible.



Which conditions of pressure and temperature favour the **reverse** reaction?

	pressure	temperature
A	high	high
B	high	low
C	low	high
D	low	low


309. 9701_m18_qp_12 Q: 12

Which statement about the effect of a catalyst on a reversible reaction is correct?

- A** The activation energy of the forward reaction stays the same.
B The composition of the equilibrium mixture stays the same.
C The rate of the backward reaction stays the same.
D The value of the equilibrium constant changes.
-

310. 9701_s18_qp_11 Q: 8

Ethanedioate ions, $\text{C}_2\text{O}_4^{2-}$, react with a suitable reagent to form CO_2 . A half-equation for this reaction is shown.



Which row is correct?

	oxidation state of carbon in $\text{C}_2\text{O}_4^{2-}$	type of reaction
A	+3	oxidation
B	+3	reduction
C	+5	oxidation
D	+5	reduction

311. 9701_s18_qp_11 Q: 9

Oxidation numbers should be used to answer this question.

A redox reaction takes place between hydroxylammonium ions, $[\text{NH}_3\text{OH}]^+$, and acidified iron(III) ions, Fe^{3+} . The products are iron(II) ions, Fe^{2+} , H^+ ions, water and a compound of nitrogen.

The mole ratio of reacting hydroxylammonium ions to reacting iron(III) ions is 1 : 2.

Which nitrogen-containing compound could be formed in the reaction?

A NH_3 **B** N_2O **C** NO **D** NO_2

312. 9701_s18_qp_12 Q: 8

Sulfur reacts with concentrated nitric acid in a redox reaction.




What are the changes in oxidation number of sulfur and of nitrogen in this reaction?

	sulfur	nitrogen
A	+2	-3
B	+2	-1
C	+4	-3
D	+4	-1

6.1. REDOX PROCESSES: ELECTRON TRANSFER AND CHANGES IN OXIDATION NUMBER (OXIDATION STATE)

313. 9701_s18_qp_13 Q: 8

Xenon hexafluoride, XeF_6 , reacts with water.




Which statement is correct?

- A** Hydrogen is reduced in this reaction.
- B** Hydrogen is the only element oxidised in this reaction.
- C** The only element oxidised in this reaction is xenon.
- D** This is not a redox reaction.

314. 9701_w18_qp_11 Q: 7

Nitric acid is known to take part in the oxidation of atmospheric sulfur dioxide. One possible reaction is shown.



Which row shows the correct changes in oxidation numbers of nitrogen and sulfur?

	nitrogen	sulfur
A	−3	+3
B	−2	+2
C	−2	+3
D	−1	+2

315. 9701_w18_qp_11 Q: 8

A transition metal ion, M^{2+} , reacts with acidified dichromate(VI) ions to form M^{4+} ions, Cr^{3+} ions, and H_2O .

Which equation correctly represents this reaction?

- A** $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + \text{M}^{2+} \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O} + \text{M}^{4+}$
- B** $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 2\text{M}^{2+} \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O} + 2\text{M}^{4+}$
- C** $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 3\text{M}^{2+} \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O} + 3\text{M}^{4+}$
- D** $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{M}^{2+} \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O} + 6\text{M}^{4+}$

316. 9701_w18_qp_12 Q: 8

Ammonium metavanadate, NH_4VO_3 , can be used to make a solution containing VO_2Cl , which contains chloride ions.

What is the **change** in the oxidation number of vanadium in this reaction?

- A** −1
- B** 0
- C** +1
- D** +2

317. 9701_m17_qp_12 Q: 8

HOCl(aq) is the molecule that kills bacteria when chlorine is added to water.

The following reaction produces this molecule.



Which statement about this reaction is correct?

- A** Chlorine is both oxidised and reduced.
 - B** Chlorine is oxidised but not reduced.
 - C** Hydrogen is both oxidised and reduced.
 - D** Hydrogen is oxidised but not reduced.
-

318. 9701_s17_qp_11 Q: 8

Solutions containing chlorate(I) ions are used as household bleaches and disinfectants. These solutions decompose on heating as shown.




Which oxidation state is shown by chlorine in each of these three ions?

	ClO^-	ClO_3^-	Cl^-
A	+1	+3	-1
B	-1	+3	+1
C	+1	+5	-1
D	-1	+5	+1

319. 9701_s17_qp_11 Q: 9

When K_2MnO_4 is dissolved in water, the following reaction occurs.




What are the values of a and c in the balanced chemical equation?

	a	c
A	2	1
B	3	2
C	4	3
D	5	4

320. 9701_s17_qp_12 Q: 8

A reaction sequence is shown.





In each stage of this sequence the sulfur is oxidised, reduced or neither oxidised nor reduced.

Which row is correct?

	1	2	3	4
A	neither	oxidised	reduced	reduced
B	oxidised	neither	reduced	reduced
C	oxidised	neither	reduced	oxidised
D	oxidised	oxidised	reduced	oxidised

321. 9701_s17_qp_13 Q: 4

Two conversions are shown.



Which similar feature do these two conversions have?

- A** change in oxidation state of an element
- B** decrease in bond angle
- C** formation of a lone pair of electrons
- D** loss of a π bond

322. 9701_s17_qp_13 Q: 7

Vanadium reacts with dilute sulfuric acid to form $\text{V}_2(\text{SO}_4)_3$ and hydrogen gas.

What happens to vanadium atoms in this reaction?

- A** They lose three electrons and are oxidised.
- B** They lose three electrons and are reduced.
- C** They lose two electrons and are oxidised.
- D** They lose two electrons and are reduced.

323. 9701_s17_qp_13 Q: 8

$\text{Na}_2\text{S}_2\text{O}_3$ reacts with HCl as shown.



When calculating the oxidation number of sulfur in $\text{Na}_2\text{S}_2\text{O}_3$, the **average** oxidation number of the two sulfur atoms should be found.

What is the oxidation number of sulfur in each of $\text{Na}_2\text{S}_2\text{O}_3$, SO_2 , and S ?

	$\text{Na}_2\text{S}_2\text{O}_3$	SO_2	S
A	+2	+2	+1
B	+2	+4	0
C	+4	+4	0
D	+5	+4	0

324. 9701_w17_qp_11 Q: 6

One of the reactions in a lead / acid cell is shown.



Which statement about this reaction is correct?

- A** Lead is both oxidised and reduced.
- B** Lead is neither oxidised nor reduced.
- C** Lead is oxidised only.
- D** Lead is reduced only.

325. 9701_w17_qp_12 Q: 8

Which statement is **always** correct for an oxidation reaction?

- A** It involves the gain of oxygen by an element.
- B** For one reactant to be oxidised a different reactant must be reduced.
- C** The element or ion being oxidised will gain electrons.
- D** The oxidation number of the element being oxidised will increase.


326. 9701_m16_qp_12 Q: 1

Which compound contains two different elements with identical oxidation states?


- A** HClO
- B** $\text{Mg}(\text{OH})_2$
- C** Na_2SO_4
- D** NH_4Cl

327. 9701_s16_qp_12 Q: 9

In the treatment of domestic water supplies, chlorine is added to the water to form HClO .



The HClO reacts further to give ClO^- ions.



Both HClO and ClO^- kill bacteria by oxidation.

What is the change in oxidation number of chlorine when forming the ClO^- ion from aqueous chlorine?

- A** -1 **B** 0 **C** +1 **D** +2

328. 9701_s16_qp_12 Q: 17

Nitrogen(II) oxide, NO , nitrogen(IV) oxide, NO_2 , carbon monoxide, CO , and unburnt hydrocarbons are present in the exhaust gases of internal combustion engines. When catalytic converters are used to remove these compounds from the exhaust gases, redox reactions occur.

What happens to each compound in the catalytic converter?

	NO	NO_2	CO	hydrocarbons
A	oxidised	oxidised	reduced	oxidised
B	oxidised	oxidised	oxidised	oxidised
C	reduced	reduced	oxidised	oxidised
D	reduced	reduced	reduced	reduced

329. 9701_w16_qp_11 Q: 3


The reaction between acidified dichromate(VI) ions, $\text{Cr}_2\text{O}_7^{2-}$, and aqueous Fe^{2+} ions results in the dichromate(VI) ions being reduced to Cr^{3+} ions.

What is the correct equation for this reaction?

- A** $\text{Cr}_2\text{O}_7^{2-} + \text{Fe}^{2+} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + \text{Fe}^{3+} + 7\text{H}_2\text{O}$
B $\text{Cr}_2\text{O}_7^{2-} + 2\text{Fe}^{2+} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 2\text{Fe}^{3+} + 7\text{H}_2\text{O}$
C $\text{Cr}_2\text{O}_7^{2-} + 3\text{Fe}^{2+} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 3\text{Fe}^{3+} + 7\text{H}_2\text{O}$
D $\text{Cr}_2\text{O}_7^{2-} + 6\text{Fe}^{2+} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 6\text{Fe}^{3+} + 7\text{H}_2\text{O}$

330. 9701_w16_qp_11 Q: 9

Sodium chromate(VI), Na_2CrO_4 , is manufactured by heating chromite, FeCr_2O_4 , with sodium carbonate in an oxidising atmosphere. Chromite contains $\text{Cr}_2\text{O}_4^{2-}$ ions.




What happens in this reaction?

- A** Chromium and iron are the only elements oxidised.
- B** Chromium, iron and carbon are oxidised.
- C** Only chromium is oxidised.
- D** Only iron is oxidised.

331. 9701_s15_qp_11 Q: 4

In oxygen difluoride, OF_2 , fluorine has an oxidation number of -1 . OF_2 will react with sulfur dioxide according to the following equation.






What is oxidised and what is reduced in this reaction?

	fluorine	oxygen in OF_2	sulfur
A	oxidised	oxidised	reduced
B	oxidised	reduced	oxidised
C	reduced	oxidised	reduced
D	reduced	reduced	oxidised

332. 9701_s15_qp_12 Q: 10

Photochromic glass, used for sunglasses, darkens when exposed to bright light and becomes more transparent again when the light is less bright. The darkness of the glass is related to the concentration of silver atoms.

The following reactions are involved.



Which statement about these reactions is correct?

- A** Cu^+ and Cu^{2+} ions act as catalysts.
- B** Cu^+ ions act as an oxidising agent in reaction 2.
- C** Reaction 3 increases the darkness of the glass.
- D** Silver atoms are reduced in reaction 3.

333. 9701_s15_qp_13 Q: 5

In which reaction does an element have the largest change in oxidation number?

- A** $\text{Cr}_2\text{O}_7^{2-} + 6\text{Fe}^{2+} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 6\text{Fe}^{3+} + 7\text{H}_2\text{O}$
- B** $3\text{OCl}^- \rightarrow \text{ClO}_3^- + 2\text{Cl}^-$
- C** $5\text{Fe}^{2+} + \text{MnO}_4^- + 8\text{H}^+ \rightarrow 5\text{Fe}^{3+} + \text{Mn}^{2+} + 4\text{H}_2\text{O}$
- D** $\text{PbO}_2 + \text{Sn}^{2+} + 4\text{H}^+ \rightarrow \text{Sn}^{4+} + \text{Pb}^{2+} + 2\text{H}_2\text{O}$

334. 9701_s15_qp_13 Q: 16

Transition elements and their compounds are widely used as catalysts.

What is the identity and what is the oxidation number of the element present in the catalyst used in the Contact process?

	element	oxidation number
A	iron	0
B	iron	+3
C	vanadium	0
D	vanadium	+5

335. 9701_w15_qp_11 Q: 19

Element X forms a pollutant oxide Y. Y can be further oxidised to Z. Two students made the following statements.

Student P 'The molecule of Y contains lone pairs of electrons.'

Student Q 'The oxidation number of X increases by 1 from Y to Z.'

X could be carbon or nitrogen or sulfur.

Which student(s) made a correct statement?

- A** P only
- B** Q only
- C** both P and Q
- D** neither P nor Q

336. 9701_w15_qp_12 Q: 5

Calcium forms an ionic compound with carbon, called calcium carbide. The oxidation number of carbon in calcium carbide is -1 .

Calcium carbide is readily hydrolysed by water giving two products only.

What could be the formulae of calcium carbide and the two products of hydrolysis?

	calcium carbide	products
A	Ca_2C	CaO and C_2H_4
B	Ca_2C	$\text{Ca}(\text{OH})_2$ and C_2H_2
C	CaC_2	CaO and C_2H_4
D	CaC_2	$\text{Ca}(\text{OH})_2$ and C_2H_2

Paper 1 Multiple Choice Answers: Electrochemistry

Question	Paper	Q. No.	Answer
281	9701_m22_qp_12	10	A
282	9701_m21_qp_12	8	C
283	9701_s21_qp_11	9	D
284	9701_s21_qp_12	9	A
285	9701_w21_qp_11	9	C
286	9701_w21_qp_12	10	D
287	9701_m20_qp_12	5	D
288	9701_m20_qp_12	10	D
289	9701_s20_qp_11	2	C
290	9701_s20_qp_11	8	B
291	9701_s20_qp_12	5	D
292	9701_s20_qp_12	10	B
293	9701_s20_qp_13	10	C
294	9701_w20_qp_11	8	A
295	9701_w20_qp_11	9	B
296	9701_w20_qp_12	1	A
297	9701_w20_qp_12	9	C
298	9701_m19_qp_12	10	B
299	9701_s19_qp_11	9	B
300	9701_s19_qp_12	9	C
301	9701_s19_qp_13	1	D
302	9701_s19_qp_13	8	A
303	9701_w19_qp_11	6	A
304	9701_w19_qp_11	9	D
305	9701_w19_qp_12	8	C
306	9701_w19_qp_12	15	C
307	9701_m18_qp_12	10	D
308	9701_m18_qp_12	11	C
309	9701_m18_qp_12	12	B
310	9701_s18_qp_11	8	A
311	9701_s18_qp_11	9	B
312	9701_s18_qp_12	8	D
313	9701_s18_qp_13	8	D
314	9701_w18_qp_11	7	B
315	9701_w18_qp_11	8	C
316	9701_w18_qp_12	8	B
317	9701_m17_qp_12	8	A
318	9701_s17_qp_11	8	C
319	9701_s17_qp_11	9	B
320	9701_s17_qp_12	8	C
321	9701_s17_qp_13	4	B
322	9701_s17_qp_13	7	A
323	9701_s17_qp_13	8	B
324	9701_w17_qp_11	6	A
325	9701_w17_qp_12	8	D
326	9701_m16_qp_12	1	A

Paper 1 Multiple Choice Answers: Electrochemistry (continued)

Question	Paper	Q. No.	Answer
327	9701_s16_qp_12	9	C
328	9701_s16_qp_12	17	C
329	9701_w16_qp_11	3	D
330	9701_w16_qp_11	9	A
331	9701_s15_qp_11	4	B
332	9701_s15_qp_12	10	A
333	9701_s15_qp_13	5	C
334	9701_s15_qp_13	16	D
335	9701_w15_qp_11	19	A
336	9701_w15_qp_12	5	D

Chapter 6


Electrochemistry

6.1 Redox processes: electron transfer and changes in oxidation number

23. 9701_s20_qp_21 Q: 2

(a) The equation shown in (a)(i) describes the reaction which occurs when aqueous potassium iodide is added to aqueous copper(II) sulfate. A white precipitate of copper(I) iodide forms in a brown solution of iodine and potassium sulfate.

(i) Balance the equation and include state symbols.



[2]

The table gives the oxidation numbers of iodine in the different species in the equation.

iodine-containing species	oxidation number of iodine
KI	−1
CuI	−1
I ₂	0

(ii) Deduce the oxidation number of copper in CuSO₄ and CuI.

- oxidation number of copper in CuSO₄
- oxidation number of copper in CuI

[1]

(iii) Describe the type of reaction shown by the equation in (a)(i). Explain your answer in terms of electron transfer.

.....

 [2]

(b) In the reaction described in (a)(i), a student uses 17.43 g of CuSO₄·yH₂O. By further titration of the reaction products the student concludes that the total amount of CuSO₄ in the sample is 0.0982 mol.

Use the *Data Booklet* to complete the table to calculate the value of y, where y is an integer. Show your working.

mass of 0.0982 mol CuSO ₄	 g
amount of H ₂ O in 17.43 g of CuSO ₄ ·yH ₂ O	 mol H ₂ O
value of y	y = ...

[4]

[Total: 9]

24. 9701_s19_qp_23 Q: 2

Iodine is an element in Group 17 of the Periodic Table.

- (a) (i) At room temperature, iodine solid has a lattice structure.

Describe the arrangement of the iodine molecules within the solid.

.....
 [1]

- (ii) List **all** of the forces of attraction present in solid iodine and identify which of these are overcome when solid iodine is heated to produce iodine vapour.

force(s) of attraction present


.....

force(s) of attraction overcome

.....
 [2]

Iodine reacts with aluminium to form a white solid, Al_2I_6 .

- (b) The diagram shows the arrangement of the outer electrons within a molecule of Al_2I_6 .



- (i) How many co-ordinate (dative covalent) bonds are made when a molecule of Al_2I_6 is formed from its atoms?

..... [1]

- (ii) Describe how co-ordinate (dative covalent) bonds form within this molecule.

.....

 [1]

(c) In a reaction between hydrogen iodide and concentrated sulfuric acid, the products are hydrogen sulfide, sulfur, iodine and water.

(i) Write an equation for this reaction.

You may wish to use oxidation numbers to help you.

..... [2]


(ii) Explain, with reference to oxidation numbers, why this reaction is a redox reaction.

.....
.....
..... [2]

[Total: 9]

25. 9701_s16_qp_22 Q: 3

Acidified potassium dichromate(VI) can oxidise ethanedioic acid, $\text{H}_2\text{C}_2\text{O}_4$.
The relevant half-equations are shown.



- (a) State the overall equation for the reaction between acidified dichromate(VI) ions and ethanedioic acid.

..... [2]

- (b) In an experiment a 0.242g sample of hydrated ethanedioic acid, $\text{H}_2\text{C}_2\text{O}_4 \cdot x\text{H}_2\text{O}$, was reacted with a $0.0200 \text{ mol dm}^{-3}$ solution of acidified potassium dichromate(VI).

32.0 cm^3 of the acidified potassium dichromate(VI) solution was required for complete oxidation of the ethanedioic acid.

- (i) Calculate the amount, in moles, of dichromate(VI) ions used to react with the sample of ethanedioic acid.

amount = ... mol [1]

- (ii) Calculate the amount, in moles, of ethanedioic acid in the sample.

amount = ... mol [1]

- (iii) Calculate the relative molecular mass, M_r , of the hydrated ethanedioic acid.

M_r = ... [1]

- (iv) Calculate the value of x in $\text{H}_2\text{C}_2\text{O}_4 \cdot x\text{H}_2\text{O}$.

x = ... [1]

[Total: 6]

23. 9701_s20_ms_21 Q: 2

(a)(i)	$2\text{CuSO}_4(\text{aq}) + 4\text{KI}(\text{aq}) \rightarrow 2\text{CuI}(\text{s}) + (\text{+})\text{I}_2(\text{aq}) + 2\text{K}_2\text{SO}_4(\text{aq})$ M1 correct balancing M2 correct state symbols	2						
(a)(ii)	Oxidation state of copper in CuSO_4 (+)2 AND Oxidation state of copper in CuI (+)1	1						
(a)(iii)	M1 redox	1						
	M2 iodide ions – lost electron(s) AND copper ions – gained electron(s)	1						
(b)	<table><tr><td>Mass of 0.0982mol CuSO_4 in 17.43g $\text{CuSO}_4 \cdot y\text{H}_2\text{O}$</td><td>M1 calculate M_r CuSO_4 using A_r from data booklet $63.5 + 32.1 + 64.0 = 159.6$ M2 use M_r to calculate mass of CuSO_4 $(0.0982 \times M1) = 15.67272\text{g}$</td></tr><tr><td>number of water in 17.43g of $\text{CuSO}_4 \cdot y\text{H}_2\text{O}$</td><td>M3 calculate the mass amount of water in sample AND use this value to calculate the amount of water present $(17.43 - 15.67) / 18 = 0.097778 \text{ mol}$</td></tr><tr><td>value of y</td><td>M4 use the ratio of M2: 0.0982 to find y $(\text{mol H}_2\text{O} \div \text{mol CuSO}_4) = 1$</td></tr></table>	Mass of 0.0982mol CuSO_4 in 17.43g $\text{CuSO}_4 \cdot y\text{H}_2\text{O}$	M1 calculate M_r CuSO_4 using A_r from data booklet $63.5 + 32.1 + 64.0 = 159.6$ M2 use M_r to calculate mass of CuSO_4 $(0.0982 \times M1) = 15.67272\text{g}$	number of water in 17.43g of $\text{CuSO}_4 \cdot y\text{H}_2\text{O}$	M3 calculate the mass amount of water in sample AND use this value to calculate the amount of water present $(17.43 - 15.67) / 18 = 0.097778 \text{ mol}$	value of y	M4 use the ratio of M2: 0.0982 to find y $(\text{mol H}_2\text{O} \div \text{mol CuSO}_4) = 1$	4
Mass of 0.0982mol CuSO_4 in 17.43g $\text{CuSO}_4 \cdot y\text{H}_2\text{O}$	M1 calculate M_r CuSO_4 using A_r from data booklet $63.5 + 32.1 + 64.0 = 159.6$ M2 use M_r to calculate mass of CuSO_4 $(0.0982 \times M1) = 15.67272\text{g}$							
number of water in 17.43g of $\text{CuSO}_4 \cdot y\text{H}_2\text{O}$	M3 calculate the mass amount of water in sample AND use this value to calculate the amount of water present $(17.43 - 15.67) / 18 = 0.097778 \text{ mol}$							
value of y	M4 use the ratio of M2: 0.0982 to find y $(\text{mol H}_2\text{O} \div \text{mol CuSO}_4) = 1$							

24. 9701_s19_ms_23 Q: 2

(a)(i)	held in regular / uniform arrangement	1
(a)(ii)	M1 covalent (bonds) AND (temporary) induced dipoles M2 (temporary) induced dipoles	2
(b)(i)	2	1
(b)(ii)	iodine (atom / s) donates a pair of electrons (to the A-I covalent bond / s).	1
(c)(i)	$2\text{H}_2\text{SO}_4 + 14\text{HI} \rightarrow 7\text{I}_2 + 8\text{H}_2\text{O} + \text{H}_2\text{S} + \text{S}$ M1 correct species M2 correctly balanced equation	2
(c)(ii)	<i>explain with ref to ox no's why the reaction in (c)(i) is a redox reaction</i> M1 I (oxidation number increases) from -1 to 0 = oxidation / reducing agent M2 S (oxidation number decreases) from (+) 6 to 0 OR -2 = reduction / oxidising agent	2

25. 9701_s16_ms_22 Q: 3

(a)	$\text{Cr}_2\text{O}_7^{2-} + 8\text{H}^+ + 3\text{H}_2\text{C}_2\text{O}_4 \rightarrow 2\text{Cr}^{3+} + 6\text{CO}_2 + 7\text{H}_2\text{O}$ M1 = species M2 = balancing	[1] [1]
(b) (i)	$(0.02 \times 32.0 / 1000) = 6.40 \times 10^{-4}$	[1]
(ii)	$(6.4 \times 10^{-4} \times 3) = 1.92 \times 10^{-3}$	[1]
(iii)	$(0.242 / 1.92 \times 10^{-3}) = 126(.0)$	[1]
(iv)	$(126 - 90 = 36; 36 / 18 = 2 \text{ hence}) x = 2$	[1]

1.

During electrolysis of brine in a diaphragm cell, chlorine, hydrogen and sodium hydroxide are produced.

What is the molar ratio of these products?

	chlorine	hydrogen	sodium hydroxide
A	1	1	1
B	1	1	2
C	2	1	1
D	2	2	1

[2002 M/J (16)]

2.

In the industrial electrolysis of brine to manufacture chlorine, the diaphragm used is a porous screen which allows the flow of electrolytes but keeps other chemicals separate.

Which substance needs to be kept separate from the chlorine by the diaphragm?

- A hydrogen
B sodium hydroxide
C sodium chloride
D water

[2002 O/N (18)]

3.

Use of the Data Booklet is relevant to this question.

In the commercial electrolysis of brine, the products are chlorine, hydrogen and sodium hydroxide.


What is the maximum yield of each of these products when 58.5 kg of sodium chloride are electrolysed as brine?

	yield of chlorine / kg	yield of hydrogen / kg	yield of sodium hydroxide / kg
A	35.5	1	40
B	35.5	2	40
C	71	1	40
D	71	2	80

[2004 M/J (18)]

4.

Strontium metal can be obtained by the electrolysis of molten strontium bromide, SrBr_2 , using the apparatus shown in the diagram.




Why is an atmosphere of argon used around the cathode?

- A** The argon keeps the strontium molten.
- B** The argon stops the molten strontium rising too high in the tube.
- C** A thin film of a compound of strontium and argon forms on the surface protecting the freshly formed metal.
- D** Without the argon strontium oxide would form in the air.

[2004 O/N (15)]

5.

A weedkiller can be prepared by heating a bleach solution.



What are the oxidation states of chlorine in these three compounds?

- | | | | |
|----------|----|----|----|
| A | -1 | -1 | +5 |
| B | +1 | -1 | +5 |
| C | +1 | -1 | +7 |
| D | +2 | +1 | +7 |

[2004 O/N (16)]

6.


In the extraction of aluminium by electrolysis, why is it necessary to dissolve aluminium oxide in molten cryolite?

- A to reduce the very high melting point of the electrolyte
- B cryolite provides the ions needed to carry the current
- C cryolite reacts with the aluminium oxide to form ions
- D molten aluminium oxide alone would not conduct electricity

[2005 M/J (8)]

7.

A cheap carbon monoxide detector for a gas heater consists of a patch containing palladium chloride crystals. When carbon monoxide is present, the crystals turn from orange to black as the following reaction takes place.



Which is the element whose oxidation number **decreases** in this reaction?

- A carbon
- B chlorine
- C hydrogen
- D palladium

[2005 M/J (9)]

8.


During the electrolysis of brine using a diaphragm cell, which reaction occurs at the cathode?

- A $2\text{Cl}^-(\text{aq}) \rightarrow \text{Cl}_2(\text{g}) + 2\text{e}^-$
- B $4\text{OH}^-(\text{aq}) \rightarrow \text{O}_2(\text{g}) + 2\text{H}_2\text{O(l)} + 4\text{e}^-$
- C $2\text{H}_2\text{O(l)} + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$
- D $\text{Na}^+(\text{aq}) + \text{e}^- \rightarrow \text{Na(s)}$

[2005 O/N (8)]

9.

The nickel-cadmium rechargeable battery is based upon the following overall reaction.



What is the oxidation number of nickel at the beginning and at the end of the reaction?

	beginning	end
A	+1.5	+2
B	+2	+3
C	+3	+2
D	+3	+4

[2006 M/J (9)]

10.

When copper is extracted from its ores, the metal is not pure enough for electrical uses. The impure copper, which contains small amounts of silver and gold, is purified by electrolysis. During this process a 'sludge' forms beneath the anode which is found to contain silver and traces of gold.

Why is silver found in this sludge?

- A Silver is less electropositive than copper.
- B Silver is more dense than copper and falls off the cathode.
- C Silver reacts with the electrolyte to form an insoluble chloride salt.
- D Silver reacts with the electrolyte to form an insoluble sulfate salt.

[2009 M/J (9)]

11.

Which conversion involves a reduction of chromium?




- A $\text{CrO}_4^{2-} \rightarrow \text{CrO}_3$
- B $\text{CrO}_4^{2-} \rightarrow \text{Cr}_2\text{O}_7^{2-}$
- C $\text{CrO}_2\text{Cl}_2 \rightarrow \text{CrO}_4^{2-}$
- D $\text{CrO}_2\text{Cl}_2 \rightarrow \text{Cr}_2\text{O}_3$

[2009 O/N-11 (7)]

12.

Photochromic glass, used for sunglasses, darkens when exposed to bright light and becomes more transparent again when the light is less bright. The depth of colour of the glass is related to the concentration of silver atoms.

The following reactions are involved.



Which statement about these reactions is correct?

- A Cu^+ and Cu^{2+} ions act as catalysts.
- B Cu^+ ions act as an oxidising agent in reaction 2.
- C Reaction 2 is the one in which light is absorbed.
- D Ag^+ ions are oxidised in reaction 1.

[2009 O/N-11 (11)]

13.

How does concentrated sulfuric acid behave when it reacts with sodium chloride?

- A as an acid only
- B as an acid and oxidising agent
- C as an oxidising agent only
- D as a reducing agent only

[2010 M/J-11 (13)]

14.


What happens when chlorine is bubbled through aqueous potassium iodide?

- A Chlorine is oxidised to chlorate(V) ions.
- B Chlorine is oxidised to chloride ions.
- C Iodide ions are oxidised to iodine.
- D There is no observable reaction.

[2010 O/N-12 (16)]

15.

Strontium metal can be obtained by the electrolysis of molten strontium bromide, SrBr_2 , using the apparatus shown in the diagram.



Why is an atmosphere of argon used around the cathode?

- A A thin film of a compound of strontium and argon forms on the surface protecting the freshly formed metal.
- B The argon keeps the strontium molten.
- C The argon stops the molten strontium rising too high in the tube.
- D Without the argon, strontium oxide would form in the air.

[2010 O/N-12 (17)]

16.

In the extraction of aluminium by the electrolysis of molten aluminium oxide, why is cryolite added to the aluminium oxide?

- A to ensure the aluminium is not oxidised
- B to ensure the anode is not oxidised
- C to lower the melting point of the aluminium oxide
- D to prevent corrosion of the cathode

[2011 M/J-11 (3)]

17.

Aluminium is the most abundant metal in the Earth's crust. The extraction of aluminium is carried out by the electrolysis of aluminium oxide dissolved in molten cryolite.


Which material is used for each of the electrodes in this electrolysis?

	anode	cathode
A	aluminium	carbon
B	carbon	carbon
C	carbon	steel
D	steel	aluminium

[2011 M/J-12 (3)]

18.

When chlorine and aqueous sodium hydroxide are heated together the following overall reaction occurs.




What are the oxidation numbers for chlorine in each of the following species?

	Cl_2	NaCl	NaClO_3
A	0	+1	-5
B	+2	-1	+3
C	0	-1	+5
D	-2	+1	-3

[2011 O/N-11 (7)]

19.

The amount of titanium dioxide in an ore can be determined by using the following reaction.



Which element increases in oxidation number in this reaction?

- A bromine
- B fluorine
- C oxygen
- D titanium

[2011 O/N-12 (11)]

20.

Chlorine can be manufactured from brine in a diaphragm cell.

Which row represents the correct electrodes?

	nature of anode	nature of cathode
A	graphite	titanium
B	steel	titanium
C	titanium	graphite
D	titanium	steel

[2011 O/N-12 (15)]

21.

Sodium iodide reacts with concentrated sulfuric acid. The equation which represents one of the reactions that takes place is shown.




Which species has been oxidised in this reaction?

- A H^+
- B I^-
- C Na^+
- D SO_4^{2-}

[2011 O/N-12 (16)]

22.

The diagram shows a cell for the manufacture of aluminium.



Which statement is **incorrect**?

- A Aluminium ions are oxidised in this process.
- B Aluminium is liberated at the cathode by the reaction $Al^{3+} + 3e^- \rightarrow Al$.
- C The cryolite acts as a solvent.
- D The graphite anode burns away.

[2012 M/J-11 (11)]

23.

Aluminium is extracted by the electrolysis of a molten mixture containing aluminium oxide. By a similar method, magnesium is extracted by the electrolysis of a molten mixture containing magnesium chloride.


Which statement about the extraction of magnesium is correct?

- A Magnesium ions travel to the anode and are oxidised to magnesium metal.
- B Magnesium ions travel to the anode and are reduced to magnesium metal.
- C Magnesium ions travel to the cathode and are oxidised to magnesium metal.
- D Magnesium ions travel to the cathode and are reduced to magnesium metal.

[2012 O/N-11 (7)]

24.

The diagram shows an electrolytic cell for the extraction of aluminium.



Which statement is correct?

- A Aluminium ions are oxidised in this process.
- B Aluminium is liberated at the anode by the reaction $Al^{3+} + 3e^- \rightarrow Al$.
- C Cryolite is purified aluminium oxide.
- D The graphite anode burns away.

[2013 M/J-11 (3)]

25.

During the electrolysis of molten aluminium oxide to produce aluminium, using carbon electrodes, two consecutive reactions occur at the anode, each producing a different gas.

How does the oxidation number of oxygen change in these reactions?

- A decreases by 2, then increases by 2
- B increases by 2, then decreases by 2
- C increases by 2, then decreases by 4
- D no change, then decreases by 2

[2013 M/J-12 (1)]

26.

In the redox reaction shown, how do the oxidation states of vanadium and sulfur change?



	vanadium		sulfur	
	from	to	from	to
A	+1	+3	0	-2
B	+1	+3	+4	+6
C	+5	+3	0	-2
D	+5	+3	+4	+6

[2013 M/J-13 (1)]

27.

A space shuttle's upward thrust came from the following reaction between aluminium and ammonium perchlorate.



Which statements about this reaction are correct?

- 1 Aluminium is oxidised.
- 2 Chlorine is reduced.
- 3 Nitrogen is oxidised.

[2013 M/J-13 (31)]

28.

At the age of 17, in a woodshed in Ohio, Charles Martin Hall discovered the commercial process for the production of aluminium metal by the electrolysis of a mixture of bauxite, Al_2O_3 , and cryolite, Na_3AlF_6 .

What is the main purpose of the cryolite?

- A Al_2O_3 is covalent, and AlF_6^{3-} ions interact with it to produce Al^{3+} ions which can be discharged at the cathode.
- B Cryolite is a base, forming NaAlO_2 with bauxite, enabling aluminium to be discharged at the anode.
- C Cryolite minimises the release of O^{2-} ions at the graphite anodes, which are otherwise burnt away to CO .
- D Cryolite reduces the melting point of the bauxite.

[2013 O/N-11 (1)]

29.


In which reaction does a single nitrogen atom have the greatest change in oxidation number?

- A $4\text{NH}_3 + 5\text{O}_2 \rightarrow 4\text{NO} + 6\text{H}_2\text{O}$
- B $3\text{NO}_2 + \text{H}_2\text{O} \rightarrow 2\text{HNO}_3 + \text{NO}$
- C $2\text{NO} + \text{O}_2 \rightarrow 2\text{NO}_2$
- D $4\text{NH}_3 + 6\text{NO} \rightarrow 5\text{N}_2 + 6\text{H}_2\text{O}$

[2013 O/N-11 (2)]

30.

Strontium metal can be obtained by the electrolysis of molten strontium bromide, SrBr_2 , using the apparatus shown in the diagram.



Why is an atmosphere of argon used around the cathode?

- A A thin film of a compound of strontium and argon forms on the surface protecting the freshly formed metal.
- B The argon keeps the strontium molten.
- C The argon stops the molten strontium rising too high in the tube.
- D Without the argon, strontium oxide would form in the air.

[2013 O/N-11 (12)]

31.

Ammonium nitrate, NH_4NO_3 , can decompose explosively when heated.



What are the changes in the oxidation numbers of the two nitrogen atoms in NH_4NO_3 when this reaction proceeds?

- A -2, -4 B +2, +6 C +4, -6 D +4, -4

[2013 O/N-13 (1)]

32.

In the extraction of aluminium by electrolysis, why is it necessary to dissolve aluminium oxide in molten cryolite?

- A to reduce the very high melting point of the electrolyte
B cryolite is a base; aluminium oxide is amphoteric
C cryolite reacts with the aluminium oxide to form ions
D molten aluminium oxide alone would not conduct electricity

[2013 O/N-13 (2)]

33.

The electrolysis of brine using the diaphragm cell is an important industrial process.


What happens at the anode?

- A Chloride ions are oxidised.
B Hydrogen gas is produced.
C Hydroxide ions are formed.
D The electrode reaction is $2\text{Cl}^-(\text{aq}) + 2\text{e}^- \rightarrow \text{Cl}_2(\text{g})$.

[2014 M/J-13 (5)]

34.

Many crude oils contain H_2S . During refining, by the Claus process, the H_2S is converted into solid sulfur, which is then removed.



Which statements about the Claus process are correct?

- 1 H_2S is oxidised in reaction I.
2 SO_2 oxidises H_2S in reaction II.
3 Hydrogen is oxidised in reaction II.

[2014 M/J-13 (33)]

35.

In which reaction does hydrogen behave as an oxidising agent?

- A $\text{H}_2 + \text{Cl}_2 \rightarrow 2\text{HCl}$
B $\text{C}_2\text{H}_4 + \text{H}_2 \rightarrow \text{C}_2\text{H}_6$
C $\text{N}_2 + 3\text{H}_2 \rightarrow 2\text{NH}_3$
D $2\text{Na} + \text{H}_2 \rightarrow 2\text{NaH}$

[2014 O/N-12 (2)]

36.

Impure copper is purified by electrolysis. The electrolyte used in this process is aqueous copper(II) sulfate.

Which reaction takes place at the anode?


- A $\text{Cu}(\text{s}) \rightarrow \text{Cu}^{2+}(\text{aq}) + 2\text{e}^-$
B $\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$
C $2\text{OH}^-(\text{aq}) \rightarrow \frac{1}{2}\text{O}_2(\text{g}) + \text{H}_2\text{O}(\text{l}) + 2\text{e}^-$
D $2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$

[2014 O/N-13 (4)]

37.

In a solution that contains both Br_2 and Cl_2 , a process takes place that produces BrO_3^- ions.

The process is represented by the following equations.



Which statements about these reactions are correct?

- 1 Chlorine is reduced in equation 2.
- 2 Bromine is oxidised in both equation 1 and equation 2.
- 3 Bromine is reduced in both equation 1 and equation 2.

[2014 O/N-13 (31)]

1. Nov/2020/Paper_41/No.3

- (a) Identify the substances liberated at the anode and at the cathode during the electrolysis of aqueous sodium sulfate, $\text{Na}_2\text{SO}_4(\text{aq})$.

anode

cathode

[1]

- (b) When molten sodium chloride is electrolysed, chlorine is liberated at the anode and sodium is liberated at the cathode.

A sample of molten sodium chloride is electrolysed for 1.50 hours using a current of 4.50 A.

Calculate the volume of chlorine and the mass of sodium that are liberated under room conditions.

volume of chlorine = ... dm^3

mass of sodium = ... g

[4]

- (c) The equation representing the standard electrode potential, E° , for the reduction of $\text{MnO}_4^-(\text{aq})$ to $\text{Mn}^{2+}(\text{aq})$ in acid solution is given.



- (i) Draw a diagram of the apparatus that would be used to measure the E° value of this half-cell. Your diagram should be fully labelled to identify all apparatus, substances and conditions.

[4]

- (ii) Use the *Data Booklet* to identify a substance that could be used to oxidise Mn^{2+} ions to MnO_4^- ions under standard conditions.

Write an equation for the reaction.

.....

.....

..... [2]

[Total: 11]

- (a) Identify the substances liberated at the anode and at the cathode during the electrolysis of saturated $KCl(aq)$.

at the anode

at the cathode

[1]

- (b) When dilute sulfuric acid is electrolysed, oxygen is liberated at the anode.

Dilute sulfuric acid is electrolysed for 15.0 minutes using a current of 0.750 A.

Calculate the volume of oxygen that is liberated under room conditions.

volume of oxygen = ... cm^3 [3]

- (c) The halogens chlorine, bromine and iodine differ in their strengths as oxidising agents. These strengths are indicated by the E° values for these halogens.

- (i) Give the E° values for chlorine, bromine and iodine acting as oxidising agents.

..... [1]

- (ii) Deduce which of chlorine, bromine and iodine will react with a solution of $Sn^{2+}(aq)$ under standard conditions.

Explain your answer. Include a relevant equation in your explanation.

.....

.....

..... [3]

(iii) An excess of chlorine is added to a solution of acidified $\text{Mn}^{2+}(\text{aq})$ under standard conditions.

Give the formula of the product of this reaction that contains manganese.

..... [1]

(d) An electrochemical cell can be made by connecting an $\text{Fe}^{3+}/\text{Fe}^{2+}$ half-cell to an $\text{S}_2\text{O}_8^{2-}/\text{SO}_4^{2-}$ half-cell under standard conditions.

(i) Calculate the standard cell potential of this electrochemical cell.

$E_{\text{cell}}^{\circ} = \dots\dots\dots \text{V}$ [1]

(ii) State the material that should be used as the electrode in each half-cell.

in the $\text{Fe}^{3+}/\text{Fe}^{2+}$ half-cell

in the $\text{S}_2\text{O}_8^{2-}/\text{SO}_4^{2-}$ half-cell

[1]

(iii) Describe **one** change to each half-cell that would **increase** the value of the cell potential. The temperature should remain at 298 K.

$\text{Fe}^{3+}/\text{Fe}^{2+}$ half-cell

.....

$\text{S}_2\text{O}_8^{2-}/\text{SO}_4^{2-}$ half-cell

.....

[1]

[Total: 12]

Gold is an unreactive metal that can only be oxidised under specific conditions.

(a) The standard electrode potential, $E^\ominus$, of $\text{Au}^{3+}(\text{aq})/\text{Au}(\text{s})$ is +1.50 V.

(i) Define the term *standard electrode potential*.

.....
.....
..... [2]

(ii) Draw a fully labelled diagram of the apparatus that should be used to measure the standard cell potential, $E^\ominus_{\text{cell}}$, of $\text{Au}^{3+}(\text{aq})/\text{Au}(\text{s})$ and $\text{HNO}_3(\text{aq})/\text{NO}(\text{g})$.

Include all necessary chemicals.

[4]

Some relevant half-equations and their standard electrode potentials are given.

	half-equation	E°/V
1	$\text{Au}^{3+}(\text{aq}) + 3\text{e}^- \rightleftharpoons \text{Au}(\text{s})$	+1.50
2	$[\text{AuCl}_4]^- (\text{aq}) + 3\text{e}^- \rightleftharpoons \text{Au}(\text{s}) + 4\text{Cl}^- (\text{aq})$	+1.00
3	$\text{NO}_3^- (\text{aq}) + 4\text{H}^+ (\text{aq}) + 3\text{e}^- \rightleftharpoons \text{NO}(\text{g}) + 2\text{H}_2\text{O}(\text{l})$	+0.96

- (iii) Write an ionic equation to show the spontaneous reaction that occurs when an electric current is drawn from the cell in (a)(ii).

..... [1]

- (iv) Calculate the E°_{cell} of the reaction in (a)(iii).

$E^\circ_{\text{cell}} = \dots\dots\dots \text{V}$ [1]

- (v) Gold can be oxidised by a mixture of concentrated hydrochloric acid and concentrated nitric acid, known as *aqua regia*. Concentrated hydrochloric acid is 12mol dm^{-3} . Concentrated nitric acid is 16mol dm^{-3} .

Explain why *aqua regia* is able to dissolve gold.

In your answer, state and explain what effect the use of concentrated hydrochloric acid and concentrated nitric acid have on the E values of half-equations 2 and 3.

.....

.....

.....

.....

.....

..... [3]

- (b) Aqueous gold(III) chloride, AuCl_3 , reacts with aqueous hydrogen peroxide, H_2O_2 , under certain conditions, forming Au, O_2 and HCl .

A student carries out separate experiments using different initial concentrations of AuCl_3 and H_2O_2 . The initial rate of each reaction is measured.

The table shows the results that are obtained.

experiment	$[\text{AuCl}_3]$ / mol dm^{-3}	$[\text{H}_2\text{O}_2]$ / mol dm^{-3}	rate of production of $\text{O}_2(\text{g})$ / $\text{dm}^3 \text{ minute}^{-1}$
1	0.05	0.50	7.66×10^{-2}
2	0.10	0.50	1.53×10^{-1}
3	0.15	1.00	4.60×10^{-1}

- (i) Write an equation for the reaction of AuCl_3 with H_2O_2 .

..... [1]

- (ii) Determine the rate equation of the reaction.

Show your reasoning, quoting data from the table.

.....

 [3]

- (iii) Use the results of experiment 2 to calculate the value of the rate constant, k , for this reaction.

Include the units of k .


rate constant, k = ...

units = ...

[2]

(c) AlF_3 is an ionic compound.

The Born–Haber cycle for the formation of AlF_3 is shown.



(i) Name the enthalpy changes labelled ΔH_4 and ΔH_6 .

$\Delta H_4 =$

$\Delta H_6 =$

[2]

(ii) Use the data in the table and data from the *Data Booklet* to calculate the lattice energy of AlF_3 .

process	enthalpy change / kJ mol^{-1}
$\text{Al}(\text{s}) \rightarrow \text{Al}(\text{g})$	+326
$\text{Al}(\text{g}) \rightarrow \text{Al}^{3+}(\text{g})$	+5137
$\text{F}(\text{g}) \rightarrow \text{F}^{-}(\text{g})$	−328
$\text{Al}(\text{s}) + 1.5\text{F}_2(\text{g}) \rightarrow \text{AlF}_3(\text{s})$	−1504

lattice energy of $\text{AlF}_3 =$ kJ mol^{-1} [2]

(iii) Scandium fluoride, ScF_3 , is an ionic compound.

Use data from the *Data Booklet* to suggest how the lattice energy of AlF_3 compares with the lattice energy of ScF_3 .

Explain your answer.

.....
.....
..... [2]

(d) AlF_3 is sparingly soluble in water. The concentration of its saturated solution at 298 K is $6.5 \times 10^{-2} \text{ mol dm}^{-3}$.

(i) Write an expression for the solubility product, K_{sp} , of AlF_3 .

$K_{\text{sp}} =$ [1]


(ii) Calculate the numerical value of K_{sp} for AlF_3 at 298 K.

$K_{\text{sp}} =$ [1]

[Total: 25]

4. June/2020/Paper_41/No.3

The overall reaction for photosynthesis is shown.



Water is oxidised in this process according to the following half-equation.



- (a) (i) Use these equations to deduce the half-equation for the reduction of carbon dioxide in this process.

[2]

- (ii) Draw a fully labelled diagram of the apparatus that should be used to measure the standard electrode potential, E° , of $\text{O}_2(\text{g})$ in half-equation 1 under standard conditions. Include all necessary chemicals.

[4]

- (iii) For the cell drawn in (a)(ii), use the *Data Booklet* to calculate the E°_{cell} and deduce which electrode is positive.

$$E^\circ_{\text{cell}} = \dots\dots\dots \text{V}$$

identity of the positive electrode = $\dots\dots\dots$
[1]

(a) Silver carbonate, Ag_2CO_3 , is sparingly soluble in water. The numerical value of the solubility product, K_{sp} , for silver carbonate is 6.3×10^{-12} at 25°C .

(i) Write an expression for the solubility product, K_{sp} , of Ag_2CO_3 , and state its units.

$$K_{\text{sp}} =$$

units = ...
[2]

(ii) Calculate the equilibrium concentration of Ag^+ in a saturated solution of Ag_2CO_3 at 25°C .

$$[\text{Ag}^+] = \dots\dots\dots \text{mol dm}^{-3} \quad [1]$$

(iii) Solid Ag_2CO_3 is stirred at 25°C with $0.050 \text{ mol dm}^{-3} \text{ AgNO}_3$ until no more Ag_2CO_3 dissolves.

Calculate the concentration of carbonate ions, $[\text{CO}_3^{2-}]$, in this solution.


$$[\text{CO}_3^{2-}] = \dots\dots\dots \text{mol dm}^{-3} \quad [1]$$

(iv) An electrochemical cell is set up to measure the electrode potential, E , for the Ag^+/Ag half-cell using the saturated $\text{Ag}_2\text{CO}_3(\text{aq})$ with a standard hydrogen electrode.

Use the *Data Booklet*, your answer to (a)(ii), and the Nernst equation to calculate the electrode potential, E , for this Ag^+/Ag half-cell.

$$E \text{ for } \text{Ag}^+/\text{Ag} \text{ half-cell} = \dots\dots\dots \text{V} \quad [2]$$

- (a) Manganese(IV) oxide, MnO_2 , catalyses the decomposition of hydrogen peroxide, H_2O_2 , as shown.



The mechanism involves the formation of the intermediate species, Mn^{2+} , in the first step which is subsequently used up in the second step.

State and use relevant electrode potentials, $E^\ominus$, to construct **two** equations to show how MnO_2 can catalyse this reaction.

.....

.....

.....

.....

.....

equation 1

equation 2

[3]

- (b) The equation for the decomposition of hydrogen peroxide without a catalyst is shown.






Under certain conditions this reaction is found to be first order with respect to hydrogen peroxide, with a rate constant, k , of $2.0 \times 10^{-6} \text{ s}^{-1}$ at 298 K.

Calculate the initial rate of decomposition of a 0.75 mol dm^{-3} hydrogen peroxide solution at 298 K.

initial rate = ... $\text{mol dm}^{-3} \text{ s}^{-1}$ [1]

- (c) A four-step mechanism is suggested for the reaction between hydrogen peroxide and iodide ions in an acidic solution.



Step 1 is the rate-determining step.

- (i) State what is meant by the term *rate-determining step*.

.....
..... [1]

- (ii) Use this mechanism to construct a balanced equation for this reaction.

..... [1]

- (iii) Deduce the order of reaction with respect to each of the following.

H_2O_2 = ... I^- = ... H^+ = ... [1]

[Total: 7]

(a) (i) Define the term *electron affinity*.

.....

.....

..... [2]

(ii) Define the term *lattice energy*.


.....

.....

..... [2]

1. Nov/2021/Paper_41/No.1

When dilute sulfuric acid is electrolysed, water is split into hydrogen and oxygen.



A current of $x\text{A}$ is passed through the solution for 14.0 minutes. 462 cm^3 of hydrogen are produced at the cathode, measured under room conditions.

(a) Calculate the number of hydrogen molecules produced during the electrolysis.

number of hydrogen molecules = ... [2]

(b) Calculate the total number of electrons transferred to produce this number of hydrogen molecules.

total number of electrons = ... [1]

(c) Calculate the quantity of charge, in coulombs, of the total number of electrons calculated in **(b)**.

quantity of charge = ... C [1]

(d) Calculate the current, x , passed during this experiment.

$x = \dots\dots\dots\text{A}$ [1]

(e) The standard entropies, S° , of three species are given in the table.

species	$S^\circ / \text{JK}^{-1} \text{mol}^{-1}$
$\text{H}_2\text{O}(\text{l})$	+70
$\text{H}_2(\text{g})$	+131
$\text{O}_2(\text{g})$	+205

(i) Calculate ΔS° for the reaction $2\text{H}_2\text{O}(\text{l}) \rightarrow 2\text{H}_2(\text{g}) + \text{O}_2(\text{g})$.

$$\Delta S^\circ = \dots\dots\dots \text{JK}^{-1} \text{mol}^{-1} \quad [1]$$

(ii) ΔH° for the reaction $2\text{H}_2\text{O}(\text{l}) \rightarrow 2\text{H}_2(\text{g}) + \text{O}_2(\text{g})$ is $+572 \text{kJ mol}^{-1}$.

Calculate ΔG° for this reaction at 298 K.

$$\Delta G^\circ = \dots\dots\dots \text{kJ mol}^{-1} \quad [2]$$

(iii) Predict the effect of increasing temperature on the spontaneity of this reaction.
Explain your answer.

.....
..... [1]

[Total: 9]

(a) Define the term *electron affinity*.

.....
 [2]

(b) Write an equation for the process corresponding to the **second** ionisation energy of calcium. Include state symbols.

..... [1]

Some data relating to calcium and oxygen are listed. Select relevant data from this list for your answers to parts (c), (d) and (e).

process	value / kJ mol ⁻¹
first ionisation energy of oxygen	+1310
second ionisation energy of oxygen	+3390
first electron affinity of oxygen	-142
second electron affinity of oxygen	+844
enthalpy change for $\frac{1}{2}\text{O}_2(\text{g}) + 2\text{e}^- \rightarrow \text{O}^{2-}(\text{g})$	+951
enthalpy change for $\text{Ca}(\text{s}) \rightarrow \text{Ca}^{2+}(\text{g}) + 2\text{e}^-$	+1933
lattice energy of CaO(s)	-3517

(c) Oxygen exists as O₂ molecules.

Use the data in this question to calculate a value for the bond energy of the O=O bond. Show all your working.

bond energy = ... kJ mol⁻¹ [3]

(d) (i) Suggest why the first electron affinity of oxygen is negative.

.....
..... [1]

(ii) Suggest why the second electron affinity of oxygen is positive.

.....
..... [1]

(e) Calculate the enthalpy of formation of calcium oxide, CaO(s) .

enthalpy of formation = ... kJ mol^{-1} [2]

(f) The lattice energy of lithium fluoride, LiF(s) , is $-1022 \text{ kJ mol}^{-1}$.

Identify the factor that causes the lattice energy of calcium oxide to be more exothermic than that of lithium fluoride. Explain why this factor causes the difference in lattice energies.

.....
..... [2]

[Total: 12]

An excess of sodium iodide is added to a solution of copper(II) sulfate. Iodine and a white precipitate of copper(I) iodide are formed.

(a) Write an equation for the reaction that occurs.

..... [1]

(b) (i) Explain why the copper(II) sulfate solution is coloured.

.....

 [4]

(ii) Suggest why the precipitate of copper(I) iodide is white.

.....
 [1]

(c) Use suitable E° values from the *Data Booklet* to predict whether iodide ions can reduce Cu^{2+} to Cu^+ under standard conditions. Explain your answer.

.....

 [2]


(d) An excess of sodium iodide is added to copper(II) sulfate solution. Copper(I) iodide forms as a precipitate. After precipitation, $[\text{Cu}^+]$ is much lower than 1.0 mol dm^{-3} .

Use this information and your answer to (c) to explain how the relevant electrode potentials change and hence why I^- ions can reduce Cu^{2+} ions.


.....

 [2]

Pure water is a very poor conductor of electricity. However, when hydrogen chloride gas is dissolved in water, ions are formed and a current flows during electrolysis.



The overall change after electrolysis is that hydrogen chloride gas is converted into hydrogen and chlorine.



When a current of 3.10A is passed through the solution for Y minutes, 351 cm³ of chlorine are produced at the anode, measured under room conditions.

(a) Calculate the number of chlorine molecules produced during the electrolysis.

number of chlorine molecules = ... [2]

(b) Calculate the total number of electrons transferred to produce this number of chlorine molecules.

total number of electrons = ... [1]

(c) Calculate the quantity of charge, in coulombs, of the total number of electrons calculated in (b).

quantity of charge = ... C [1]

(d) Calculate the time, Y, in minutes, for which the current flows.

Y = ... minutes [1]

(a) (i) Define the term *transition element*.

.....
..... [1]

(ii) State how the melting point and density of iron compare to those of calcium.

.....
..... [1]

(b) (i) Define the term *standard cell potential*, $E_{\text{cell}}^{\ominus}$.

.....
.....
..... [2]

(ii) Draw a fully labelled diagram of the apparatus that can be used to measure the cell potential of a cell composed of a Cu(II)/Cu electrode and an Fe(III)/Fe(II) electrode. Include all necessary reactants.

(c) The reaction between $\text{S}_2\text{O}_8^{2-}(\text{aq})$ and $\text{I}^-(\text{aq})$ is catalysed by adding a few drops of $\text{Fe}^{3+}(\text{aq})$.

(i) Use equations to show the catalytic role of Fe^{3+} in this reaction.

[2]


(ii) $\text{Fe}^{3+}(\text{aq})$ can oxidise $\text{I}^-(\text{aq})$, whereas $[\text{Fe}(\text{CN})_6]^{3-}(\text{aq})$ cannot oxidise $\text{I}^-(\text{aq})$.

Use E° values to explain these observations.

.....
.....
.....
..... [2]

(d) When aqueous solutions of $\text{S}_2\text{O}_8^{2-}$ and tartrate ions are mixed the reaction proceeds very slowly. However, this reaction proceeds quickly in the presence of an $\text{Fe}^{3+}(\text{aq})$ catalyst. The overall equation for this reaction is shown.

tartrate ions



(i) Suggest why this reaction is slow without the Fe^{3+} catalyst.


.....
..... [1]

(ii) Use the overall equation to deduce the half-equation for the oxidation of tartrate ions, $\text{C}_4\text{H}_4\text{O}_6^{2-}$, to carbon dioxide, CO_2 , and methanoate ions, HCO_2^- .



- (e) (i) Complete the following table to show the structures of the organic products formed when tartaric acid reacts separately with each reagent. Identify each type of reaction.

tartaric acid



reagent	structure of organic product	type of reaction
an excess of LiAlH_4		
an excess of CH_3COCl		

[3]

- (ii) Tartaric acid reacts with the amine 1-phenylethylamine, $\text{C}_6\text{H}_5\text{CH}(\text{NH}_2)\text{CH}_3$, to form an ionic salt.

Draw the structure of the salt formed in this reaction. Include the charges on the ions.

[1]

[Total: 17]

- (a) Samples of $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$ are reacted separately with an excess of aqueous sodium hydroxide or with an excess of aqueous ammonia.

Give the following information about these reactions.

- (i) reaction 1: $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$ with an excess of aqueous of sodium hydroxide

colour and state of the copper-containing species

ionic equation

type of reaction

[3]

- (ii) reaction 2: $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$ with an excess of aqueous ammonia

colour and state of the copper-containing species

ionic equation

type of reaction

[3]

- (b) Copper(I) oxide is added to hot dilute sulfuric acid. A blue solution, **X**, and a red-brown solid, **Y**, form.

Suggest the identities of **X** and **Y**. Name the type of reaction.

X

Y

type of reaction

[2]

[Total: 8]

(a) (i) Define the term *standard electrode potential*.

.....


.....

..... [2]

Three redox systems, **A**, **B** and **C**, are shown. The ligand 1,2-diaminoethane, $\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2$, is represented by en.

A	$[\text{Ru}(\text{H}_2\text{O})_6]^{3+} + \text{e}^- \rightleftharpoons [\text{Ru}(\text{H}_2\text{O})_6]^{2+}$
B	$[\text{Ru}(\text{NH}_3)_6]^{3+} + \text{e}^- \rightleftharpoons [\text{Ru}(\text{NH}_3)_6]^{2+}$
C	$[\text{Ru}(\text{en})_3]^{3+} + \text{e}^- \rightleftharpoons [\text{Ru}(\text{en})_3]^{2+}$

Two electrochemical cells are set up to compare the standard electrode potentials, E° , of three half-cells. The diagrams show the relative potential of each electrode.



(ii) Use this information to complete the table by adding the labels **A**, **B** and **C** to deduce the order of E° for the three half-cells.

E°	redox system
most negative	
↑	
least negative	


[1]

(iii) The complex $[\text{Ru}(\text{en})_3]^{3+}$ shows stereoisomerism. The ligand en is bidentate.

Draw three-dimensional diagrams to show the two isomers of $[\text{Ru}(\text{en})_3]^{3+}$. Represent the

ligand en by using 

Name the type of stereoisomerism.



type of stereoisomerism [3]

- (b) (i) An electrochemical cell consists of a Br_2/Br^- half-cell and a Ag^+/Ag half-cell, under standard conditions.

Use the *Data Booklet* to calculate the $E_{\text{cell}}^\ominus$. Deduce the direction of electron flow in the wire through the voltmeter between these two half-cells.

$E_{\text{cell}}^\ominus = \dots\dots\dots \text{V}$

direction of electron flow from to [1]

- (ii) Water is added to the Ag^+/Ag half-cell in (b)(i).


Suggest the effect of this addition on the $E_{\text{cell}}^\ominus$. Place a tick (✓) in the appropriate box.

less positive	no change	more positive

Explain your answer.

.....
.....
..... [2]

- (c) Silver bromide, AgBr, dissolves in an aqueous solution of $\text{S}_2\text{O}_3^{2-}$ ions to form the complex ion $[\text{Ag}(\text{S}_2\text{O}_3)_2]^{3-}$. The $\text{S}_2\text{O}_3^{2-}$ ions act as monodentate ligands.



- (i) Define the term *ligand*.

.....
 [1]

- (ii) Write an expression for the equilibrium constant, K_c , for equilibrium 1.

$K_c =$

[1]

- (iii) Some additional data are given about the dissolution of AgBr in $\text{S}_2\text{O}_3^{2-}(\text{aq})$.

equilibrium constant	numerical value
solubility product, K_{sp} , of AgBr	5.4×10^{-13}
stability constant, K_{stab} , of $[\text{Ag}(\text{S}_2\text{O}_3)_2]^{3-}$	2.9×10^{13}

Use your answer to (c)(ii) and these data to calculate K_c for equilibrium 1. Include the units for K_c .

$K_c =$ units [2]

- (d) The numerical values for the stability constants, K_{stab} , of two other silver(I) complexes are given.

silver(I) complex	numerical value of K_{stab}
$[\text{Ag}(\text{CN})_2]^{-}$	5.3×10^{18}
$[\text{Ag}(\text{NH}_3)_2]^{+}$	1.6×10^7

An aqueous solution containing Ag^{+} is added to a solution containing equal concentrations of $\text{CN}^{-}(\text{aq})$, $\text{NH}_3(\text{aq})$ and $\text{S}_2\text{O}_3^{2-}(\text{aq})$. The mixture is left to reach equilibrium.

Deduce the relative concentrations of $[\text{Ag}(\text{CN})_2]^{-}$, $[\text{Ag}(\text{NH}_3)_2]^{+}$ and $[\text{Ag}(\text{S}_2\text{O}_3)_2]^{3-}$ present in the resulting mixture. Explain your answer.

..... > >
 highest concentration lowest concentration

 [2]

[Total: 15]

1. June/2022/Paper_41/No.2(d, e)

- (d) Manganese(VII) oxide, Mn_2O_7 , can be made by treatment of KMnO_4 with concentrated sulfuric acid (reaction 1).

Mn_2O_7 readily decomposes at room temperature to form manganese(IV) oxide and a colourless diatomic gas (reaction 2).

Construct equations for **both** the reactions described.

reaction 1

reaction 2

[2]

(e) Aqueous manganese(II) ions show similar chemical properties to aqueous copper(II) ions when reacted separately with NaOH(aq) and with concentrated HCl.

- (i) Write the ionic equation, and state the type of reaction, for the reaction of $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$ with NaOH(aq).

ionic equation

type of reaction

[2]

- (ii) Write the ionic equation, and state the type of reaction, for the reaction of $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$ with concentrated HCl.

ionic equation

type of reaction

[2]

- (iii) Table 2.1 lists relevant electrode potentials for some electrode reactions.

Table 2.1

electrode reaction	E°/V
$\text{Mn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Mn}$	-1.18
$\text{Cl}_2 + 2\text{e}^- \rightleftharpoons 2\text{Cl}^-$	+1.36
$2\text{HOCl} + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{Cl}_2 + 2\text{H}_2\text{O}$	+1.64
$\text{MnO}_2 + 4\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{Mn}^{2+} + 2\text{H}_2\text{O}$	+1.23
$\text{MnO}_4^- + 4\text{H}^+ + 3\text{e}^- \rightleftharpoons \text{MnO}_2 + 2\text{H}_2\text{O}$	+1.67

Suggest the formula of the manganese species formed when $\text{Mn}^{2+}(\text{aq})$ reacts with Cl_2 .

State the type of reaction.

formula of manganese species formed

type of reaction

[1]

(c) (i) Define standard electrode potential, E° .

.....

.....

..... [1]

(ii) A salt bridge is used in an electrochemical cell.


State the function of the salt bridge. Explain your answer.

.....

..... [1]

(iii) Complete the diagram of the apparatus that can be used to measure the E° of the $\text{Cr}_2\text{O}_7^{2-}(\text{aq})$, $\text{H}^+(\text{aq})/\text{Cr}^{3+}(\text{aq})$ electrode against the standard hydrogen electrode.

Your diagram should be fully labelled to identify all apparatus, substances and conditions.



[3]

(iv) The E° of the $\text{Cr}_2\text{O}_7^{2-}(\text{aq})$, $\text{H}^+(\text{aq})/\text{Cr}^{3+}(\text{aq})$ electrode is +1.33 V.

Label the negative electrode and the direction of electron flow in the external circuit when the current flows in your diagram in (c)(iii). [1]

- (d) Table 3.1 lists relevant electrode potentials for some electrode reactions for use in (d)(i) and (d)(ii).

Table 3.1

electrode reaction	E°/V
$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^- \rightleftharpoons 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	+1.33
$\text{CH}_3\text{CHO} + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{CH}_3\text{CH}_2\text{OH}$	-0.61
$\text{CH}_3\text{COOH} + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{CH}_3\text{CHO} + \text{H}_2\text{O}$	-0.94
$\text{O}_2 + 4\text{H}^+ + 4\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}$	+1.23

- (i) Ethanal is oxidised to ethanoic acid in the presence of $\text{Cr}_2\text{O}_7^{2-}$ ions.

Construct the ionic equation for the oxidation of ethanal to ethanoic acid using dichromate(VI) in acid conditions. Calculate the E°_{cell} for this reaction.

ionic equation

$$E^\circ_{\text{cell}} = \dots\dots\dots \text{V} \quad [2]$$

- (ii) In an ethanol-oxygen fuel cell, $\text{CH}_3\text{CH}_2\text{OH}(\text{l})$ and $\text{O}_2(\text{g})$ are in contact with two inert electrodes immersed in an acidic solution.

The cell reaction for the oxidation of ethanol by oxygen is shown.



Calculate ΔG° , in kJ mol^{-1} , for the oxidation of ethanol by oxygen.

$$\Delta G^\circ = \dots\dots\dots \text{kJ mol}^{-1} \quad [2]$$

- (a) Complete Table 5.1 to predict the substance liberated at each electrode during electrolysis of the indicated electrolyte with inert electrodes.

Table 5.1

electrolyte	substance liberated at the anode	substance liberated at the cathode
$\text{PbBr}_2(\text{l})$		
concentrated $\text{NaCl}(\text{aq})$		
$\text{Cu}(\text{NO}_3)_2(\text{aq})$		

[3]

- (b) An electrolytic cell is set up to determine a value for the Avogadro constant, L . The electrolyte is dilute sulfuric acid and both electrodes are copper.

When a current of 0.600 A is passed through the acid for 30.0 minutes, the anode decreases in mass by 0.350 g.

- (i) State the relationship between the Faraday constant, F , and the Avogadro constant, L .

..... [1]

- (ii) Use the experimental information in (b) and data from the table on page 23 to calculate a value for the Avogadro constant, L .

Show all working.

Avogadro constant, $L =$ [4]

[Total: 8]

1. Nov/2022/Paper_41/No.3

Data should be selected from Table 3.1 in order to answer some parts of this question.

Table 3.1


electrode reaction	E° / V
$\text{Cl}_2 + 2\text{e}^- \rightleftharpoons 2\text{Cl}^-$	+1.36
$2\text{HOCl} + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{Cl}_2 + 2\text{H}_2\text{O}$	+1.64
$\text{ClO}^- + \text{H}_2\text{O} + 2\text{e}^- \rightleftharpoons \text{Cl}^- + 2\text{OH}^-$	+0.89
$\text{Sn}^{4+} + 2\text{e}^- \rightleftharpoons \text{Sn}^{2+}$	+0.15
$\text{Sn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Sn}$	-0.14
$\text{V}^{2+} + 2\text{e}^- \rightleftharpoons \text{V}$	-1.20
$\text{V}^{3+} + \text{e}^- \rightleftharpoons \text{V}^{2+}$	-0.26
$\text{VO}^{2+} + 2\text{H}^+ + \text{e}^- \rightleftharpoons \text{V}^{3+} + \text{H}_2\text{O}$	+0.34
$\text{VO}_2^+ + 2\text{H}^+ + \text{e}^- \rightleftharpoons \text{VO}^{2+} + \text{H}_2\text{O}$	+1.00

(a) Standard electrode potentials are measured under standard conditions.

(i) Describe the standard conditions used in the $\text{Sn}^{4+}/\text{Sn}^{2+}$ half-cell.

.....
 [1]

(ii) Complete the diagram below to show how $E^\circ(\text{Sn}^{4+}/\text{Sn}^{2+})$ can be measured experimentally. Your diagram should be fully labelled to identify all apparatus and substances.



[3]

(iii) Equal volumes of $1.0 \text{ mol dm}^{-3} \text{ Sn}^{2+}(\text{aq})$ and $1.0 \text{ mol dm}^{-3} \text{ Cl}^{-}(\text{aq})$ are mixed.

Use relevant E° values to explain whether a reaction occurs between these two ions.

.....
.....
.....
..... [2]

(iv) Equal volumes of 1.0 mol dm^{-3} of $\text{Sn}^{2+}(\text{aq})$ and acidified $1.0 \text{ mol dm}^{-3} \text{ VO}^{2+}(\text{aq})$ are mixed.

Write an equation for the reaction that takes place in the resulting mixture.

..... [2]

(b) A solution of $\text{SnCl}_2(\text{aq})$ is electrolysed for a measured time using a steady current.

A mass of 2.95 g of tin metal is produced at the cathode.

$\text{Al}_2\text{O}_3(\text{l})$ is electrolysed for the same time by the same current.

Calculate the mass of aluminium metal produced at the cathode. Give your answer to **three** significant figures. Show your working.

mass of aluminium metal = ... g [2]

[Total: 10]

Data should be selected from Table 3.1 in order to answer some parts of this question.

Table 3.1

electrode reaction	E°/V
$\text{Mg}^{2+} + 2\text{e}^- \rightleftharpoons \text{Mg}$	-2.38
$\text{Mn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Mn}$	-1.18
$\text{Mn}^{3+} + \text{e}^- \rightleftharpoons \text{Mn}^{2+}$	+1.49
$\text{MnO}_2 + 4\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{Mn}^{2+} + 2\text{H}_2\text{O}$	+1.23
$\text{MnO}_4^- + \text{e}^- \rightleftharpoons \text{MnO}_4^{2-}$	+0.56
$\text{MnO}_4^- + 4\text{H}^+ + 3\text{e}^- \rightleftharpoons \text{MnO}_2 + 2\text{H}_2\text{O}$	+1.67
$\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightleftharpoons \text{Mn}^{2+} + 4\text{H}_2\text{O}$	+1.52


- (a) An electrochemical cell can be constructed from a Mg^{2+}/Mg half-cell and a $\text{MnO}_4^-/\text{Mn}^{2+}$ half-cell. The standard cell potential of this cell can be calculated using the standard electrode potentials of the two half-cells.

- (i) Define standard electrode potential. Include details of the standard conditions used.

.....

 [2]

- (ii) Complete the diagram below to show an electrochemical cell constructed from a Mg^{2+}/Mg half-cell and a $\text{MnO}_4^-/\text{Mn}^{2+}$ half-cell. Label your diagram.



[3]

- (iii) Use a positive (+) sign and a negative (–) sign to identify the polarity of each of the two electrodes in your diagram.

Use an arrow and the symbol 'e' to show the direction of electron flow in the external circuit. [1]

(iv) Calculate the standard cell potential, $E_{\text{cell}}^{\ominus}$, of this cell.

$E_{\text{cell}}^{\ominus} = \dots\dots\dots \text{V}$ [1]

(v) Construct an equation for the cell reaction.

$\dots\dots\dots$ [1]

(vi) Predict how the cell reaction will change, if at all, when the solution in the Mg^{2+}/Mg half-cell is diluted by the addition of a large volume of water. Explain your answer.

$\dots\dots\dots$

$\dots\dots\dots$

$\dots\dots\dots$ [1]

(b) A molten magnesium salt is electrolysed for 15.0 minutes by a constant current.

4.75×10^{22} magnesium atoms are produced at the cathode.


Calculate the value of the current used.

current = $\dots\dots\dots \text{A}$ [2]

[Total: 11]

1. Nov/2023/Paper_9701/41/No.3(e, f)

(e) The $E^\ominus$ values for two electrode reactions are given.



(i) An electrochemical cell is constructed with the following half-cells (electrodes):

- an acidified solution of H_2O_2 , a platinum wire
- Cr^{2+} mixed with Cr^{3+} , a platinum wire.

Identify the positive half-cell and calculate the standard cell potential, $E_{\text{cell}}^\ominus$.


positive half-cell $E_{\text{cell}}^\ominus =$ V

[1]

(ii) Calculate the value of $\Delta G^\ominus$ for the cell reaction that occurs, per mole of H_2O_2 .

$\Delta G^\ominus =$ kJ mol^{-1} [2]

(f) The $E^\ominus$ values for two electrode reactions are given.



An electrochemical cell is constructed with the following half-cells.

half-cell 1 an acidified solution of H_2O_2 under standard conditions, a platinum wire

half-cell 2 a solution containing $0.020\text{ mol dm}^{-3}\text{Co}^{3+}$ and $2.0\text{ mol dm}^{-3}\text{Co}^{2+}$, a platinum wire

- (i) Use the Nernst equation to calculate the value of E , the electrode potential of half-cell 2 under these conditions.

$E = \dots\dots\dots\text{ V}$ [2]

- (ii) Write an equation for the cell reaction that occurs in this cell under these conditions.


$\dots\dots\dots$ [1]

Some electrode potentials are shown in Table 3.1.

Table 3.1

electrode reaction	$E^\ominus/\text{V}$
$\text{V}^{2+} + 2\text{e}^- \rightleftharpoons \text{V}$	-1.20
$\text{V}^{3+} + \text{e}^- \rightleftharpoons \text{V}^{2+}$	-0.26
$\text{VO}^{2+} + 2\text{H}^+ + \text{e}^- \rightleftharpoons \text{V}^{3+} + \text{H}_2\text{O}$	+0.34
$\text{VO}_2^+ + 2\text{H}^+ + \text{e}^- \rightleftharpoons \text{VO}^{2+} + \text{H}_2\text{O}$	+1.00
$\text{Fe}^{2+} + 2\text{e}^- \rightleftharpoons \text{Fe}$	-0.44
$\text{Fe}^{3+} + 3\text{e}^- \rightleftharpoons \text{Fe}$	-0.04
$\text{Fe}^{3+} + \text{e}^- \rightleftharpoons \text{Fe}^{2+}$	+0.77
$2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2$	0.00
$\text{ClO}^- + \text{H}_2\text{O} + 2\text{e}^- \rightleftharpoons \text{Cl}^- + 2\text{OH}^-$	+0.89

- (a) (i) Complete the diagram to show a standard hydrogen electrode. Label your diagram. Identify all substances. You do **not** need to state standard conditions.



[1]

- (ii) An electrochemical cell is set up using an $\text{Fe}^{3+}/\text{Fe}^{2+}$ electrode and a standard hydrogen electrode.

Identify the positive electrode in the electrochemical cell and the direction of electron flow in the external circuit.

positive electrode

Electrons flow from the electrode to the electrode.

[1]

(b) The vanadium-containing species in the electrode reactions given in Table 3.1 are V, V^{2+} , V^{3+} , VO^{2+} and VO_2^+ .

- (i) Identify **one** vanadium-containing species that does **not** react with Fe^{2+} ions under standard conditions.

Use data from Table 3.1 to explain your answer.

.....
..... [1]

- (ii) Identify **all** the vanadium-containing species that will react with Fe^{2+} ions under standard conditions.

..... [1]

- (iii) Write an equation for **one** of the possible reactions identified in (ii).

..... [1]

- (c) Another electrochemical cell is set up using an Fe^{3+}/Fe^{2+} electrode and an alkaline ClO^-/Cl^- electrode.

The concentration of Fe^{3+} is 1000 times greater than the concentration of Fe^{2+} in the Fe^{3+}/Fe^{2+} electrode. All other conditions are standard.

- (i) Use the Nernst equation to calculate the E value of the Fe^{3+}/Fe^{2+} electrode.

Show your working.

$E =$ V [2]

- (ii) Write an equation for the reaction that occurs in the cell, under these conditions.

..... [1]

- (d) Another electrochemical cell is set up using an Fe^{2+}/Fe electrode and an alkaline ClO^-/Cl^- electrode under standard conditions.

Calculate the value of $\Delta G^\ominus$ for the cell.

$\Delta G^\ominus =$ kJ mol^{-1} [3]

- (e) A solution of iron(II) sulfate, $\text{FeSO}_4(\text{aq})$ is electrolysed with iron electrodes. Under the conditions used, no gas is evolved at the cathode.

A current of 0.640A is passed for 17.0 minutes. The mass of the cathode increases by 0.185g.

Use these results to calculate an experimental value for the Avogadro constant, L .

Show your working.

$$L = \dots\dots\dots \text{mol}^{-1} \quad [3]$$

- (f) Iron(II) chloride, FeCl_2 , is oxidised by chlorine to form iron(III) chloride, FeCl_3 , under standard conditions.




Table 3.2

species	$S^\ominus / \text{JK}^{-1} \text{mol}^{-1}$
$\text{Cl}_2(\text{g})$	223
$\text{FeCl}_2(\text{s})$	120
$\text{FeCl}_3(\text{s})$	142

- (i) Use Table 3.2 and other data to calculate the Gibbs free energy change, $\Delta G^\ominus$, for this reaction.

Show your working.

$$\Delta G^\ominus = \dots\dots\dots \text{kJ mol}^{-1} \quad [3]$$

(ii) Predict whether this reaction becomes more or less feasible at a higher temperature.

Explain your answer.

The reaction becomes feasible.

explanation

.....

[1]

[Total: 18]



3. June/2023/Paper_9701/42/No.1(c_d)

(c) Potassium sulfite, K_2SO_3 , is used as a food additive.

The concentration of sulfite ions, SO_3^{2-} , can be determined by titration using aqueous acidified manganate(VII) ions, MnO_4^- .

- A 250 cm^3 solution contains 3.40 g of impure K_2SO_3 .
- 25.0 cm^3 of this solution requires 22.40 cm^3 of $0.0250\text{ mol dm}^{-3}$ acidified MnO_4^- to reach the end-point. All the SO_3^{2-} ions are oxidised. None of the other species in the impure K_2SO_3 are oxidised.

The reaction occurs as shown by the two half-equations.



- (i) Give the ionic equation for the reaction between SO_3^{2-} and acidified MnO_4^- .

..... [1]

- (ii) Calculate the percentage purity of the sample of K_2SO_3 .
Show your working.

percentage purity of $\text{K}_2\text{SO}_3 = \dots\dots\dots$ [3]

- (d) Potassium disulfite, $\text{K}_2\text{S}_2\text{O}_5$, is another food additive. The disulfite ion, $\text{S}_2\text{O}_5^{2-}$, has the displayed formula shown in Fig. 1.2.




Fig. 1.2

Deduce the geometry (shape) around the $\text{S}(\alpha)$ atom in $\text{S}_2\text{O}_5^{2-}$.

geometry around $\text{S}(\alpha) \dots\dots\dots$ [1]

4. June/2023/Paper_9701/42/No.9

(a) Define standard cell potential, $E_{\text{cell}}^{\ominus}$.

.....

..... [1]

(b) An electrochemical cell is set up to measure $E_{\text{cell}}^{\ominus}$ of a cell consisting of an $\text{Fe}^{3+}/\text{Fe}^{2+}$ half-cell and a Cl_2/Cl^- half-cell.

Draw a labelled diagram of this electrochemical cell.

Include all necessary substances. It is **not** necessary to state conditions used.

[3]

(c) The cell reaction for the electrochemical cell in (b) is shown.



Calculate $\Delta G^{\ominus}$, in kJ mol^{-1} , for this cell reaction.

$\Delta G^{\ominus} = \dots\dots\dots \text{kJ mol}^{-1}$ [2]

(d) Another experiment is set up using the same electrochemical cell.

In this experiment the Fe^{2+} concentration is 0.15 mol dm^{-3} . All other concentrations remain at their standard values.

The Nernst equation is shown.

$$E = E^\ominus + (0.059/z) \log \frac{[\text{oxidised species}]}{[\text{reduced species}]}$$

- (i) Use the Nernst equation to calculate the electrode potential, E , for the $\text{Fe}^{3+}/\text{Fe}^{2+}$ half-cell in this experiment.

$$[E^\ominus: \text{Fe}^{3+}/\text{Fe}^{2+} = +0.77 \text{ V}]$$

$$E = \dots\dots\dots \text{ V [1]}$$

- (ii) Use your answer to (d)(i) to calculate E_{cell} for this electrochemical cell.

$$E_{\text{cell}} = \dots\dots\dots \text{ V [1]}$$

[Total: 8]

5. March/2023/Paper_9701/42/No.2(a, b)


Hypophosphorous acid is an inorganic acid.

The conjugate base of hypophosphorous acid is H_2PO_2^- .

(a) Give the formula of hypophosphorous acid.

..... [1]

(b) H_2PO_2^- is a strong reducing agent. It can be used to reduce metal cations without the need for electrolysis.



(i) In an experiment, an alkaline $\text{HPO}_3^{2-}/\text{H}_2\text{PO}_2^-$ half-cell is constructed with $[\text{H}_2\text{PO}_2^-] = 0.050\text{ mol dm}^{-3}$.

All other ions are at their standard concentration.

Predict how the value of E of this half-cell differs from its $E^\ominus$ value.

Explain your answer.

.....

 [2]

(ii) The Cr^{3+}/Cr half-cell has a standard electrode potential of -0.74 V .

An electrochemical cell consists of an alkaline $\text{HPO}_3^{2-}/\text{H}_2\text{PO}_2^-$ half-cell and a Cr^{3+}/Cr half-cell.

Calculate the standard cell potential, $E_{\text{cell}}^\ominus$.

$E_{\text{cell}}^\ominus = \dots\dots\dots \text{ V}$ [1]

- (iii) Complete the diagram in Fig. 2.1 to show how the standard electrode potential of the Cr^{3+}/Cr half-cell can be measured relative to that of the standard hydrogen electrode.

Identify the chemicals, conditions and relevant pieces of apparatus.




Fig. 2.1

[3]


- (iv) Label Fig. 2.1 to show:

- which is the positive electrode
- the direction of electron flow in the external circuit.

[1]

- (v) H_2PO_2^- reduces Ni^{2+} to Ni in alkaline conditions.

Use equation 1 to construct the ionic equation for this reaction.



..... [1]

1. Nov/2021/Paper_12/No.32

Four solutions, each of concentration 0.1 mol dm^{-3} , were tested with a pH meter. The results are shown.

solution	formula of acid or base	pH
acid 1	$\text{CH}_3\text{CO}_2\text{H}$	4
acid 2	HNO_3	1
base 1	CH_3NH_2	11
base 2	NaOH	14

Which statements explain these results?

- 1 Acid 2 has a lower pH than acid 1 because it is more soluble.
- 2 Base 2 has a higher concentration of hydroxide ions in solution than base 1.
- 3 Acid 1 dissociates less than acid 2.

2. Nov/2021/Paper_13/No.9

Zinc atoms can be oxidised to Zn^{2+} ions by dichromate(VI) ions in acid solution. Chromium is reduced to Cr^{3+} in this reaction.

Which equation is correct?

- A** $\text{Cr}_2\text{O}_7^{2-} + \text{Zn} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + \text{Zn}^{2+} + 7\text{H}_2\text{O}$
- B** $\text{Cr}_2\text{O}_7^{2-} + \text{Zn} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 3\text{Zn}^{2+} + 7\text{H}_2\text{O}$
- C** $\text{Cr}_2\text{O}_7^{2-} + 3\text{Zn} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 3\text{Zn}^{2+} + 7\text{H}_2\text{O}$
- D** $2\text{Cr}_2\text{O}_7^{2-} + 3\text{Zn} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 3\text{Zn}^{2+} + 7\text{H}_2\text{O}$

3. Nov/2021/Paper_13/No.17

Z is a compound of sodium, chlorine and oxygen.

It contains 45.1% by mass of oxygen.

Z is prepared by reacting sodium hydroxide with chlorine.

Which row shows the conditions used for the reaction and the oxidation state of chlorine in Z?

	reaction conditions	oxidation state of Cl in Z
A	cold dilute NaOH	+1
B	cold dilute NaOH	+5
C	hot concentrated NaOH	+1
D	hot concentrated NaOH	+5

(c) Hydrogen sulfide gas, $\text{H}_2\text{S}(\text{g})$, is slightly soluble in water. It acts as a weak acid in aqueous solution.


(i) State the meaning of *weak acid*.

.....
 [1]

(ii) Give the formula of the conjugate base of H_2S .

..... [1]

(iii) $\text{H}_2\text{S}(\text{aq})$ reacts slowly with oxygen dissolved in water. The reaction is represented by the following equation.



Explain, with reference to oxidation numbers, why this reaction is a redox reaction.

.....

 [2]

Phosphorus is a reactive Period 3 element.

(a) Phosphorus has several allotropes. Details of two allotropes are given.

allotrope of phosphorus	formula	melting point/°C
white	P ₄	44
red	P	590

(i) White phosphorus and red phosphorus both have covalent bonding.

Suggest the types of structure shown by white phosphorus (P₄) and red phosphorus (P).

Explain why red phosphorus (P) has a higher melting point than white phosphorus (P₄).

structure of P₄

structure of P

explanation


.....
.....

[3]

(ii) Red phosphorus (P) forms when white phosphorus (P₄) is exposed to sunlight.




Use this information to draw a reaction pathway diagram to show the formation of red phosphorus (P) from white phosphorus (P₄).



[1]

(b) Some reactions of $P_4(s)$ are shown in the reaction scheme.



(i) State the oxidation number of phosphorus in P_4O_{10} .

..... [1]


(ii) Deduce the identity of **Q** and hence construct chemical equations for reactions 1 and 2.

reaction 1 $PCl_5 + \dots H_2O \rightarrow \dots$


reaction 2 $P_4O_{10} + \dots H_2O \rightarrow \dots$ [2]

(c) Triphenylphosphine is used in a type of reaction known as a *Wittig reaction*.

triphenylphosphine




where





(i) Give the empirical formula of triphenylphosphine.

..... [1]

In a Wittig reaction, an aldehyde reacts with a halogenoalkane to form an alkene. The conversion is shown in the following unbalanced equation.



Compound **H** can be made from propanal, $\text{C}_2\text{H}_5\text{CHO}$. Stage 3 in the reaction scheme is a Wittig reaction.





(ii) State the types of reaction that occur in stages 1 and 2.

stage 1

stage 2


[2]

(iii) Draw the structures of **G** and **H** in the boxes provided.



[2]

(d) Identify the organic products formed when compound **J**, shown below, is heated with hot concentrated acidified manganate(VII) ions.




[2]

[Total: 14]

6. **March/2021/Paper_12/No.8**

VO_2Cl reacts with NaI under acidic conditions.




The oxidation state of Cl is -1 in VO_2Cl and in VOCl_2 .

Which row about this reaction is correct?

	vanadium	iodine
A	is oxidised	is oxidised
B	is oxidised	is reduced
C	is reduced	is oxidised
D	is reduced	is reduced

7. **June/2021/Paper_11/No.9**

When hydrogen iodide is reacted with concentrated sulfuric acid, several reactions occur, including:





Which row gives the change in oxidation number of iodine and of sulfur in this reaction?

	change in oxidation number of iodine	change in oxidation number of sulfur
A	-1	$+6$
B	-1	$+8$
C	$+1$	-6
D	$+1$	-8


8. **June/2021/Paper_11/No.25**

When an organic compound is oxidised, any oxygen atom gained by the organic molecule is considered to be from a water molecule also producing $2\text{H}^+ + 2\text{e}^-$. Any hydrogen atom lost may be considered to be lost as $\text{H}^+ + \text{e}^-$.

These changes can be represented by the following two equations.



Compound X is oxidised by heating under reflux with hot, acidified potassium dichromate(VI) for one hour. The half-equation for the reduction reaction is shown.




Under these conditions, one mole of potassium dichromate(VI) oxidises three moles of X.

What could X be?

- A propanal
- B propan-1-ol
- C propan-1,2-diol
- D propan-1,3-diol

9. **June/2021/Paper_12/No.9**

The equation for a redox reaction is shown.




Which species is being oxidised in this reaction?

- A Sn^{2+} B Cl^- C Hg^+ D Hg^{2+}

10. **June/2021/Paper_13/No.32**

The equation shows the decomposition of three moles of an ion containing chromium in an acid solution.



Which statements are correct?

- 1 One mole of CrO_4^{3-} is reduced.
- 2 Two moles of CrO_4^{3-} are oxidised.
- 3 Three moles of electrons are transferred.

11. June/2021/Paper_13/No.35

Which reagents produce a solution of sodium chlorate(V)?


- 1 chlorine and hot concentrated sodium hydroxide solution
- 2 chlorine and cold dilute sodium hydroxide solution
- 3 chlorine dissolved in water at room temperature

12. June/2021/Paper_23/No.1(f)

(f) Sodium chlorate(I), NaClO , oxidises dilute hydrochloric acid to form three products. The products which contain chlorine have chlorine species with oxidation number -1 or 0 .

No other species changes its oxidation number during the reaction.

Use this information to complete the ionic equation.



1. June/2022/Paper_11/No.17


Solid sodium iodide reacts with concentrated sulfuric acid to form more than one product that contains sulfur.

What is the lowest oxidation number of sulfur in these products?

- A** -2 **B** 0 **C** +4 **D** +6

2. June/2022/Paper_12/No.11

NCl_3 reacts with H_2O .



The oxidation state of nitrogen does not change in this reaction.

Which statement is correct?

- A** Chlorine is reduced.
B Chlorine is oxidised.
C Hydrogen is both oxidised and reduced.
D This is not a redox reaction.

3. June/2022/Paper_12/No.12

In which row do the oxidation numbers of vanadium increase?

	smallest	→	largest
A	VO_4^{3-}	VO_3^-	VO_2^+
B	VO^{2+}	V_2O_3	VO_4^{3-}
C	V_2O_3	VO^{2+}	VO_3^-
D	VO_4^{3-}	VO_2^+	VO^{2+}

4. **June/2022/Paper_13/No.11**

NH_4NO_3 decomposes into N_2O and H_2O on heating.

Which statements are correct?

- 1 The ammonium ion is behaving as a reducing agent.
- 2 The nitrate(V) ion is behaving as an oxidising agent.
- 3 It is a redox reaction.
- 4 It is a disproportionation reaction.

- A** 1, 2, 3 and 4
B 1, 2 and 3 only
C 3 and 4 only
D 3 only

5. **June/2022/Paper_13/No.12**

A student adds 3 mol of acidified $\text{K}_2\text{Cr}_2\text{O}_7$ to an excess of I^- ions.

The chromium is all reduced to Cr^{3+} and I^- ions are oxidised to I_2 .

The I_2 released is reduced back to I^- ions by X mol of $\text{S}_2\text{O}_3^{2-}$ ions.

1 mol of I_2 is reduced by 2 mol of $\text{S}_2\text{O}_3^{2-}$ ions.

What is the value of X?

- A** 3 **B** 6 **C** 9 **D** 18

1. Nov/2022/Paper_11/No.11

Ammonium ions are converted into nitrate ions by bacteria.

What is the change in the oxidation number of nitrogen?

- A** -6 **B** +6 **C** +8 **D** +9

2. Nov/2022/Paper_11/No.12

Sodium dichromate(VI), $\text{Na}_2\text{Cr}_2\text{O}_7$, reacts with hydrogen peroxide, H_2O_2 , producing Cr^{3+} ions, water and oxygen.

What is the correctly balanced ionic equation for this reaction?

- A** $\text{Cr}_2\text{O}_7^{2-} + 2\text{H}^+ + \text{H}_2\text{O}_2 \rightarrow 2\text{Cr}^{3+} + 2\text{H}_2\text{O} + 4\text{O}_2$
B $\text{Cr}_2\text{O}_7^{2-} + 8\text{H}^+ + 3\text{H}_2\text{O}_2 \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O} + 3\text{O}_2$
C $\text{Cr}_2\text{O}_7^{2-} + 8\text{H}^+ + 6\text{H}_2\text{O}_2 \rightarrow 2\text{Cr}^{3+} + 10\text{H}_2\text{O} + 6\text{O}_2$
D $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 3\text{H}_2\text{O}_2 \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O} + 3\text{O}_2$

3. Nov/2022/Paper_12/No.5

Two moles of VO_2^+ ions react with one mole of zinc atoms in the presence of dilute acid. The products include Zn^{2+} ions and an ion, Y. Ion Y contains vanadium. Only zinc and vanadium change oxidation state in the reaction.

What is ion Y?

- A** VO_3^- **B** VO^+ **C** VO^{2+} **D** VO_2^{2+}

4. Nov/2022/Paper_22/No.1(b)

- (b) Complete Table 1.1 to show the total numbers of protons and electrons in the molecular ions NH_4^+ , CO_3^{2-} and PO_4^{3-} .

Table 1.1


molecular ion	total number of protons	total number of electrons
NH_4^+		
CO_3^{2-}		
PO_4^{3-}		

[3]

1. Nov/2023/Paper_ 9701/11/No.11

HOCl(aq) is the molecule that kills bacteria when chlorine is added to water.

The following reaction produces this molecule.




Which statement about this reaction is correct?

- A Chlorine is both oxidised and reduced.
- B Chlorine is oxidised but **not** reduced.
- C Hydrogen is both oxidised and reduced.
- D Hydrogen is oxidised but **not** reduced.

2. Nov/2023/Paper_ 9701/12/No.12

Chlorine reacts with sodium bromide.




Which words correctly describe this reaction?

- 1 redox
- 2 displacement
- 3 disproportionation

- A 1, 2 and 3
- B 1 and 2 only
- C 1 only
- D 2 only

3. Nov/2023/Paper_ 9701/12/No.13

The equation for the reaction between aqueous copper ions and aqueous iodide ions is as follows.



What is the change in oxidation state of copper?

- A +2 to -1
- B +2 to 0
- C +2 to +1
- D +4 to +2

4. Nov/2023/Paper_9701/21/No.3(a)

Potassium chlorate, KClO_3 , is widely used as an oxidising agent and to make $\text{O}_2(\text{g})$.


(a) Define oxidising agent.

.....
..... [1]

5. June/2023/Paper_9701/12/No.11

LiAlH_4 contains AlH_4^- ions in which aluminium has an oxidation state of +3.

LiAlH_4 reacts with water, as shown.



In this reaction, each of the four water molecules produces one hydroxide ion. It does this by losing one H^+ ion, which reacts with the LiAlH_4 .


Which changes in oxidation number occur in this reaction?

- A Al increases by 1 and H decreases by 1.
- B H decreases by 2 and also increases by 1.
- C H increases by 1 and also decreases by 1.
- D O decreases by 1 and H increases by 1.

6. June/2023/Paper_9701/12/No.12

The vanadium salt, VOSO_4 , is soluble in water and reacts readily with powdered manganese in dilute sulfuric acid.

The equation for the reaction is shown.




Which statement about this reaction is correct?

- A Hydrogen is oxidised in the reaction.
- B Manganese is the reducing agent in this reaction.
- C Sulfuric acid is the oxidising agent in this reaction.
- D The oxidation state of the vanadium changes from +5 to +3.

7. [June/2023/Paper_9701/13/No.2](#)

The ionic equation shows iodide ions reacting with manganate(VII) ions in acidic solution.



The letters u , v , w , x , y and z all represent whole numbers. Two or more of u , v , w , x , y and z are the same as each other.

What is the lowest possible value of v ?

- A** 2 **B** 8 **C** 10 **D** 16

8. [June/2023/Paper_9701/13/No.6](#)

Electronegativity differences can be used to help determine the oxidation number of an atom in different species. A number of rules are used which include:

- The more electronegative atom is given a negative oxidation number.
- Hydrogen is more electronegative than Group 1 metals.
- Oxygen is more electronegative than hydrogen.

Which row is correct?

	equation of reaction	redox reaction	disproportionation reaction
A	$2\text{CrO}_4^{2-} + 2\text{H}^+ \rightarrow \text{Cr}_2\text{O}_7^{2-} + \text{H}_2\text{O}$	✓	✗
B	$\text{NaH} + \text{H}_2\text{O} \rightarrow \text{NaOH} + \text{H}_2$	✓	✓
C	$3\text{MnO}_4^{2-} + 4\text{H}^+ \rightarrow \text{MnO}_2 + 2\text{MnO}_4^-$	✓	✓
D	$\text{VO}_3^- + 2\text{H}^+ \rightarrow \text{VO}_2^+ + \text{H}_2\text{O}$	✓	✗

9. [June/2023/Paper_9701/13/No.14](#)

Which particle contains nitrogen in the same oxidation state as in the ion $\text{N}_2\text{O}_2^{2-}$?

- A** NH_2F **B** N_2O_4 **C** NO_3^- **D** HNF_2

10. March/2023/Paper_9701/12/No.6

The ore psilomelane may be considered to have the general formula $\text{Ba}(\text{Mn}^{x+})(\text{Mn}^{y+})_8\text{O}_{16}(\text{OH})_4$.

In this general formula, $x+$ and $y+$ are the two different oxidation states of manganese in psilomelane.

What could be the values of x and y ?

	x	y
A	2	4
B	6	4
C	6	3
D	7	3

11. March/2023/Paper_9701/12/No.7

Silicon reacts with a mixture of calcium oxide and magnesium oxide at 1200°C .



Which statement about this reaction is correct?

- A** Calcium is reduced and silicon is neither oxidised nor reduced.
- B** Magnesium is reduced and calcium is neither oxidised nor reduced.
- C** Magnesium is reduced and silicon is neither oxidised nor reduced.
- D** Silicon is reduced and calcium is neither oxidised nor reduced.