
CHEMISTRY

9701/41

Paper 4 A Level Structured Questions

May/June 2018

MARK SCHEME

Maximum Mark: 100

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2018 series for most Cambridge IGCSE™, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

IGCSE™ is a registered trademark.

This document consists of **13** printed pages.



Cambridge Assessment
International Education

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Question	Answer	Marks
1(a)	$4\text{Na(s)} + \text{O}_2\text{(g)} \rightarrow 2\text{Na}_2\text{O(s)}$	
	balanced with all formulae correct	1
	state symbols	1
1(b)	giant ionic	1
	strong bond / attraction between AND positive and negative ions / anions and cations / Na^+ and O^{2-} / oppositely charged ions	1
1(c)(i)	the reaction produces sodium hydroxide / hydroxide ions / OH^- ions	1
	the hydroxide ions can receive / accept H^+ ions / protons	1
1(c)(ii)	Calculation of Na_2O moles $3.10 \text{ g} / 62$ OR 0.05	1
	Calculation of $[\text{OH}^-]$ $0.05 \times (2 / 0.400) = 0.25 \text{ mol dm}^{-3}$	1
	Calculation of pH $-\log 0.25 = 0.60$ $14 - 0.60 = 13.40$	1
1(d)	use of (2×109) or 218 and (2×494) or 988	1
	use of (0.5×496) or 248	1
	use of 416, 142, 844	1
	evaluation of expression correctly $\Delta H_{\text{lat}} = -416 - (2 \times 109) - (0.5 \times 496) - (2 \times 494) - (-142 + 844) = -2572$	1
1(e)	the lattice energy of Na_2S is less exothermic	1
	the sulfide ion is larger than the oxide ion / S^{2-} larger than O^{2-} / ionic radii quoted 0.184 nm and 0.140 nm AND less attraction (between the ions)/bonds are weaker	1


Question	Answer	Marks
2(a)	change in amount / mass / concentration of reactant / product per time	1
2(b)	decrease in volume or pressure	1
2(c)	$8.13 \times 10^4 / 81280 / 81300$	1
	$\text{mol}^{-2} \text{dm}^6 \text{s}^{-1}$	1
2(d)	$\sqrt{(0.00231 / (0.0046 \times 81280))} = 2.49 \times 10^{-3}$	1
2(e)	2, 1, 3	1
2(f)(i)	2	1
2(f)(ii)	the total of steps 1 and 2 / the components of 2 are two NO and one H ₂	1
2(g)(i)	time for amount or mass or concentration to halve	1
2(g)(ii)	0.02 at start and 0.01 after 2 seconds	1
	0.005 after 4 seconds and 0.0025 after 6 seconds	1
2(h)(i)	NO + $\frac{1}{2}$ O ₂ → NO ₂ or NO + O ₂ → NO ₂ + $\frac{1}{2}$ O ₂ AND NO ₂ + SO ₂ → NO + SO ₃	1
2(h)(ii)	(NO is) regenerated / reformed	1
2(h)(iii)	SO ₃ + H ₂ O → H ₂ SO ₄ AND acid rain or consequence of this described	1

PUBLISHED



Question	Answer	Marks												
3(a)	<table border="1"> <tr> <td></td><td>anode</td><td>cathode</td></tr> <tr> <td>AgNO₃ (aq)</td><td>oxygen / O₂</td><td>silver / Ag</td></tr> <tr> <td>saturated NaCl (aq)</td><td>chlorine / Cl₂</td><td>hydrogen / H₂</td></tr> <tr> <td>CuSO₄ (aq)</td><td>oxygen / O₂</td><td>copper / Cu</td></tr> </table>		anode	cathode	AgNO ₃ (aq)	oxygen / O ₂	silver / Ag	saturated NaCl (aq)	chlorine / Cl ₂	hydrogen / H ₂	CuSO ₄ (aq)	oxygen / O ₂	copper / Cu	3
	anode	cathode												
AgNO ₃ (aq)	oxygen / O ₂	silver / Ag												
saturated NaCl (aq)	chlorine / Cl ₂	hydrogen / H ₂												
CuSO ₄ (aq)	oxygen / O ₂	copper / Cu												
3(b)(i)	2I ⁻ → I ₂ + 2e ⁻	1												
	Ca ²⁺ + 2e ⁻ → Ca	1												
3(b)(ii)	- Ca / Calcium reduced and I / iodine oxidised - Oxidation number of calcium decreases from 2 to 0 - Oxidation number of iodine increases from -1 to 0 2 points = 1 mark 3 points = 2 marks	2												
3(b)(iii)	- metal / grey / silvery - purple AND vapour / gas / fumes - amount of melt decreases any 2 points for 1 mark	1												
3(c)	2 × 60 × 60 × 0.8 = 5760 C AND 5760 / 96500 = 0.060 (0.0597) F	1												
	1.11 / 55.8 = 0.020 (0.0199) mol of Fe	1												
	0.06 / 0.02 = 3 ∴ Fe ³⁺ or +3 or 3	1												




PUBLISHED


Question	Answer	Marks
4(a)	calcium – red flame	1
	barium – green flame	1
4(b)	- the temperature increases down the group - ionic radius increases / charge density decreases down the group - decreasing distortion / polarisation or decreasing weakening of bonds - of the anion / the CO_3^{2-} ion 2 points = 1 mark 3 points = 2 marks 4 points = 3 marks	3
4(c)	the gas is colourless / looks like air / cannot be seen	1
	appearance of solid doesn't change / white solid becomes white solid	1



Question	Answer	Marks
5(a)	$3s^2 3p^6 3d^{10} 4s^1$	1
	$3s^2 3p^6 3d^{10}$	1
5(b)	$\text{Cl}-\text{Cu}-\text{Cl}$	1
	one minus charge	1
5(c)		1
	octahedral and 90° or 180° labelled correctly on diagram as appropriate	1


Question	Answer	Marks
5(d)	reaction 1: blue ppt / blue solid	1
	$[\text{Cu}(\text{H}_2\text{O})_6]^{2+} + 2\text{OH}^- \rightarrow \text{Cu}(\text{OH})_2 + 6\text{H}_2\text{O}$ or $[\text{Cu}(\text{H}_2\text{O})_6]^{2+} + 2\text{OH}^- \rightarrow \text{Cu}(\text{OH})_2(\text{H}_2\text{O})_4 + 2\text{H}_2\text{O}$ or $[\text{Cu}(\text{H}_2\text{O})_6]^{2+} + 2\text{NH}_3 \rightarrow \text{Cu}(\text{OH})_2(\text{H}_2\text{O})_4 + 2\text{NH}_4^+$	1
	reaction 2: deep / dark / royal blue solution	1
	$\text{Cu}(\text{OH})_2 + 4\text{NH}_3 + 2\text{H}_2\text{O} \rightarrow [\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+} + 2\text{OH}^-$ or $\text{Cu}(\text{OH})_2 + 4\text{NH}_3 \rightarrow [\text{Cu}(\text{NH}_3)_4]^{2+} + 2\text{OH}^-$ or $\text{Cu}(\text{OH})_2(\text{H}_2\text{O})_4 + 4\text{NH}_3 \rightarrow [\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+} + 2\text{OH}^- + 2\text{H}_2\text{O}$	1
5(e)(i)	ligand exchange / displacement / replacement / substitution	1
5(e)(ii)	$K_{\text{stab}} = \frac{[\text{CuEDTA}^{2-}]}{[\text{Cu}(\text{H}_2\text{O})_6]^{2+}][\text{EDTA}^{4-}]}$	1
5(e)(iii)	stable / more stable than $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$	1
5(f)(i)	donates lone pairs / forms dative / co-ordinate bonds to (central) metal atom / metal ion	1
	donates two lone pairs / forms two (dative or coordinate) bonds	1
5(f)(ii)	$[\text{Zr}(\text{C}_2\text{O}_4)_4]^{4-}$	1
	not octahedral because 8 dative bonds to Zr or not octahedral because not 6 dative bonds to Zr or not octahedral because co-ordination number is 8 / is not 6	1

Question	Answer	Marks
6(a)(i)	D 2-chloropropane	1
	E hydrogen chloride	1
6(a)(ii)	(Friedel-Crafts) alkylation	1
6(b)(i)	$AlCl_3$ or $FeCl_3$	1
6(b)(ii)		1
6(b)(iii)	sunlight or UV OR $T > 100\text{ }^{\circ}\text{C}$	1
6(b)(iv)		1

Question	Answer	Marks
6(c)	reaction with hot $\text{KMnO}_4(\text{aq})$ 	1
	reaction with $\text{H}_2 + \text{Ni}$, high pressure 	1
6(d)		
	attacking species is NO_2^+	1
	curly arrow starting within hexagon and going to NO_2^+	1
	correct intermediate	1
	2nd curly arrow from C-H bond into ring	1

Question	Answer	Marks
7(a)	$2\text{C}_3\text{H}_7\text{OH} + 2\text{Na} \rightarrow 2\text{C}_3\text{H}_7\text{ONa} + \text{H}_2$	1
7(b)(i)	propanoic acid, phenol, propan-1-ol	1
7(b)(ii)	- propan-1-ol: O-H bond strengthened by positive inductive effect of alkyl group OR propoxide ion is destabilised by positive inductive effect of alkyl group - phenol: O-H bond weakened by negative inductive effect of ring OR phenoxide ion is stabilised by delocalisation of oxygen lone pair into ring - propanoic acid: O-H bond weakened by negative inductive effect of C=O OR propanoate ion is stabilised by delocalisation of minus charge by C=O 1 mark for a correct explanation, max 2 marks	2
7(c)	Tollens' reagent or Fehling's reagent	1
	methanoic acid gives a silver mirror/solid with Tollen's reagent OR red / orange ppt / solid with Fehlings' reagent	1
7(d)	PCl_5 or PCl_3 (+heat) or SOCl_2 (added to propanoic acid)	1
	product of first step: 	1
	add product of first step to phenol in NaOH	1
7(e)(i)	propanoic acid	1
7(e)(ii)	propan-1-ol would have peak at 0.5–6.0 because of OH group	1
	propanal would have peak at 9.3–10.5 because of CHO / aldehyde	1

Question	Answer	Marks
8(a)	correct chiral centre labelled only  $C_{15}H_{22}O_4$	1
8(b)	$C_{15}H_{22}O_4$	1
8(c)(i)		1
8(c)(ii)	CO_2	1
	oxidation / oxidative cleavage	1
8(c)(iii)	CH_3COCO_2H	1

Question	Answer	Marks
9(a)	$\text{C}_8\text{H}_{11}\text{O}_3\text{N}$	1
9(b)	yes, as it has a chiral C atom	1
9(c)(i)	(phenyldiazonium ion is stabilised because) positive charge is delocalised by ring / positive charge is spread over ring	1
9(c)(ii)		1
	N_2	1

CHEMISTRY

9701/42

Paper 4 A Level Structured Questions

May/June 2018

MARK SCHEME

Maximum Mark: 100

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2018 series for most Cambridge IGCSE™, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

IGCSE™ is a registered trademark.

This document consists of **14** printed pages.



Cambridge Assessment
International Education

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

PUBLISHED

Question	Answer	Marks
1(a)	simple molecular / simple covalent	1
	weak London forces / id-id forces / VDW forces or London forces / id-id forces / VDW forces AND small amount of energy to break	1
1(b)(i)	$\text{SiCl}_4 + 2\text{H}_2\text{O} \rightarrow \text{SiO}_2 + 4\text{HCl}$ or $\text{SiCl}_4 + 4\text{H}_2\text{O} \rightarrow \text{Si(OH)}_4 + 4\text{HCl}$	1
1(b)(ii)	white solid	1
	steamy fumes / white fumes / misty fumes	1
1(b)(iii)	moles of $\text{SiCl}_4 = 0.8505 / 170.1 = 0.005$	1
	conc of $\text{H}^+ (0.005) \times 4 / 0.8 = 0.025$	1
	$\text{pH} = -\log(0.025) = 1.6$	1
1(c)(i)	$-225.7 = 239.0 - (18.7 + 2x)$	1
	$x = +223$	1
1(c)(ii)	decrease in number of moles of gas / more moles of gas on left / reactants (ora)	1
1(d)	use of $\Delta G = \Delta H - T\Delta S$ with $\Delta G = 0$ / $\Delta G > 0$ or $T = \Delta H / \Delta S$ or $T = (640\,000 / 225.7)$	1
	2836 / 2840 (2835.6)	1


Question	Answer	Marks
2(a)	colorimetry / (change) in colour / less light transmission / measure absorbance	1
2(b)	Exp 1 and 2: rate $\times$ 1.75 and $[H_2] \times 1.75$ (when $[ICl]$ no change) or calculation e.g.: order = $(0.007 / 0.004) / (1.75 / 1.00) = 1$ or Exp 1 and 3: rate $\times$ 2.5 and $[H_2] \times 2.5$ (when $[ICl]$ no change) or Exp 2 and 3: rate $\times 10 / 7(1.43)$ and $[H_2] \times 10 / 7(1.43)$ (when $[ICl]$ no change)	1
2(c)	Exp 4 and 5: rate $\times$ 1.4 and $[ICl] \times 1.4$ (when $[H_2]$ no change) or calculation	1
2(d)	(rate=) $k[ICl][H_2]$	1
2(e)	62 500 or 6.25×10^4	1
2(f)	$ICl + H_2 \rightarrow HCl + HI$ or $ICl + H_2 \rightarrow IClH_2$ or $ICl + H_2 \rightarrow \frac{1}{2}I_2 + ClH_2$	1
	$HI + ICl \rightarrow HCl + I_2$ or $IClH_2 + ICl \rightarrow 2HCl + I_2$ or $ClH_2 + ICl \rightarrow 2HCl + \frac{1}{2}I_2$	1


PUBLISHED

Question	Answer	Marks
2(g)(i)	part mark 1: plot a graph of concentration of $[H_2]$ against time part mark 2: constant half-life (showing it is 1st order) part mark 3: draw tangent AND determine gradient (on conc vs time graph) or draw two tangents to determine two gradients (rate) (on conc vs time graph) part mark 4: if conc 1 (at time 1) / conc 2 (at time 2) = gradient 1 / gradient 2 part mark 5: plot a graph of rate against concentration of $[H_2]$ part mark 6: gives a straight-line through the origin of graph for part mark 5 2 parts = 1 mark 3 parts = 2 marks 4 parts = 3 marks	3
2(g)(ii)	[IC $\ddot{I}$] doesn't change or [IC $\ddot{I}$] only changes slightly	1
2(h)	provides an alternative route of lower activation energy / E_a or to lower E_a and more molecules with $E \geq E_a$	1


Question	Answer	Marks												
3(a)	<table border="1"> <tr> <td></td><td>anode</td><td>cathode</td></tr> <tr> <td>NaOH (aq)</td><td>oxygen / O₂</td><td>hydrogen / H₂</td></tr> <tr> <td>dilute CuCl₂ (aq)</td><td>oxygen / O₂</td><td>copper / Cu</td></tr> <tr> <td>conc MgCl₂ (aq)</td><td>chlorine / Cl₂</td><td>hydrogen / H₂</td></tr> </table>		anode	cathode	NaOH (aq)	oxygen / O₂	hydrogen / H₂	dilute CuCl ₂ (aq)	oxygen / O₂	copper / Cu	conc MgCl ₂ (aq)	chlorine / Cl₂	hydrogen / H₂	3
	anode	cathode												
NaOH (aq)	oxygen / O₂	hydrogen / H₂												
dilute CuCl ₂ (aq)	oxygen / O₂	copper / Cu												
conc MgCl ₂ (aq)	chlorine / Cl₂	hydrogen / H₂												
3(b)(i)	2Br ⁻ → Br ₂ + 2e ⁻ or 2Br ⁻ – 2e ⁻ → Br ₂	1												
	Zn ²⁺ + 2e ⁻ → Zn	1												
	Zinc / Zn ²⁺ reduced and Br ⁻ / bromide oxidised	1												
3(b)(ii)	liquid / molten metal or orange- brown / reddish brown vapour / gas (at anode) or amount of melt / electrolyte decreases	1												
3(c)	• 50 × 60 × 1.2 or 3600 C (calculation of number of Coulombs) • 3600 / 96 500 or 0.0373 F (calculation of number of Faradays) • 0.0373 F / 2 or 0.01865 / 0.0187 mol H₂ (use of stoichiometry) • 0.01865 × 24 000 = 448–449 (Use of 24 000 & answer to 3sf) 2 points = 1 mark 3 points = 2 marks 4 points = 3 marks	3												

Question	Answer	Marks
4(a)(i)	$2\text{Ca}(\text{NO}_3)_2 \rightarrow 2\text{CaO} + 4\text{NO}_2 + \text{O}_2$	1
	correct state symbols: (s), (s), (g), (g)	1
4(a)(ii)	brown gas / fumes / vapour	1
	white solid / residue (stays the same)	1
4(b)	solubility decreases (down the group)	1
	ΔH_{latt} and ΔH_{hyd} decrease / both become less exothermic / less negative	1
	ΔH_{latt} decreases / becomes less exothermic by a smaller extent ora	1
	ΔH_{sol} becomes less exothermic / less negative ora	1



Question	Answer	Marks
5(a)	$3s^2 3p^6 3d^7 4s^2$ or $3s^2 3p^6 4s^2 3d^7$	1
	$3s^2 3p^6 3d^7$	1
5(b)(i)	An ion formed by a central metal atom / ion surrounded by / bonded to (one or more) ligand(s)	1
5(b)(ii)	charge is 2–	1
		1

Question	Answer	Marks
5(b)(iii)	d-orbitals are split into two groups of orbitals	1
	absorption of light	1
	an electron is excited / (during) electron promotion (to a higher d-orbital)	1
5(b)(iv)	blue light is not absorbed or absorb (wavelength / frequency of) red and yellow / orange / green or absorbs least in the blue region	1
5(c)	(blue goes) pink	1
	ligand exchange / ligand substitution	1
5(d)(i)	F_2 or $S_2O_8^{2-}$	1
5(d)(ii)	+1.05 or +0.19	1
5(d)(iii)	$2Co^{2+} + F_2 \rightarrow 2Co^{3+} + 2F^-$ or $2Co^{2+} + S_2O_8^{2-} \rightarrow 2Co^{3+} + 2SO_4^{2-}$	1
5(e)	 trans 3D diagram – two $O_2N-Co-NO_2$ bond angles are 90° and the other is 180°	1
	cis 3D diagram – three $O_2N-Co-NO_2$ bond angles are 90° to each other	1
	cis-trans / geometric(al)	1

Question	Answer	Marks
6(a)	<u>any two from</u> K / potassium or KOH / potassium hydroxide or K_2O / potassium oxide	2
	correct products: (K) hydrogen, (KOH) water, (K_2O) water	1
6(b)(i)	bond circled between N = N	1
6(b)(ii)	phenylamine and HNO_2	1
	T=10 °C or below and diazonium ion / salt formed or structure of diazonium ion as $[C_6H_5N_2^+]$	1
	add 2-naphthol in aqueous NaOH / alkali	1
6(c)(i)	dilute / aqueous nitric acid / $HNO_3(aq)$ (at room temp.)	1
	<u>any two from</u> concentrated (acid) needed sulfuric acid / H_2SO_4 needed higher T needed ora	1
6(c)(ii)	p-orbital(s) / lone pair on oxygen / OH group delocalises into / over ring	1

Question	Answer	Marks
6(c)(iii)		
	first curly arrow	1
	structure of intermediate	1
	2nd curly arrow	1
6(c)(iv)	4-nitrophenol	1


Question	Answer	Marks
7(a)	$C_4H_8N_2O_3$	1
7(b)	amide, amine, carboxylic acid 2 correct = 1 mark 3 correct = 2 marks	2

Question	Answer	Marks
7(c)		
	peptide / amide bond / unit labelled or circled	1
	(val-asp or asp-val) rest of the dipeptide structure is correct	1
7(d)(i)	resists change in pH / pH kept within a small range	1
	when small amount of acid or alkali / base is added	1
7(d)(ii)	$\text{H}_2\text{NCHRCO}_2\text{H} + \text{H}^+ \rightarrow \text{H}_3\text{N}^+\text{CHRCO}_2\text{H}$	1
	$\text{H}_2\text{NCHRCO}_2\text{H} + \text{OH}^- \rightarrow \text{H}_2\text{NCHRCO}_2^- + \text{H}_2\text{O}$	1
7(e)	 each structure [1]	2

PUBLISHED

Question	Answer	Marks
7(f)	$\text{H}_2\text{NCH}(\text{CH}_2\text{CONH}_2)\text{CO}_2\text{H} + \text{H}_2\text{O} \rightarrow \text{H}_2\text{NCH}(\text{CH}_2\text{CO}_2\text{H})\text{CO}_2\text{H} + \text{NH}_3$ scores or $\text{H}_2\text{NCH}(\text{CH}_2\text{CONH}_2)\text{CO}_2\text{H} + \text{H}_2\text{O} + 2\text{H}^+ \rightarrow \text{H}_3\text{N}^+\text{CH}(\text{CH}_2\text{CO}_2\text{H})\text{CO}_2\text{H} + \text{NH}_4^+$	2

Question	Answer	Marks
8(a)	0, 2, 1	1
8(b)	6	1
8(c)	4	1
8(d)(i)	4	1
8(d)(ii)	range δ 25–5	1
	range δ 190–220	1
	one peak in first range and three peaks in second range	1
8(d)(iii)	1	1
8(d)(iv)	singlet	1
	neighbouring / adjacent (carbon) atom has no protons / H	1
8(e)(i)	CO_2	1
	(further) oxidation / oxidative cleavage	1

Question	Answer	Marks
8(e)(ii)	 The chemical structure shows a bicyclic system consisting of a six-membered ring fused to a five-membered ring. The six-membered ring has a ketone group (=O) at the 1-position and a methyl group at the 2-position. The five-membered ring has a methyl group at the 2-position and is connected to a side chain at the 3-position. The side chain is a 4-hydroxy-4-methylpentyl group, consisting of a five-carbon chain with a methyl group and a hydroxyl group (-OH) on the fourth carbon.	1

CHEMISTRY

9701/43

Paper 4 A Level Structured Questions

May/June 2018

MARK SCHEME

Maximum Mark: 100

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2018 series for most Cambridge IGCSE™, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

IGCSE™ is a registered trademark.

This document consists of **13** printed pages.



Cambridge Assessment
International Education

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.


Question	Answer	Marks
1(a)	$4\text{Na(s)} + \text{O}_2\text{(g)} \rightarrow 2\text{Na}_2\text{O(s)}$	
	balanced with all formulae correct	1
	state symbols	1
1(b)	giant ionic	1
	strong bond / attraction between AND positive and negative ions / anions and cations / Na^+ and O^{2-} / oppositely charged ions	1
1(c)(i)	the reaction produces sodium hydroxide / hydroxide ions / OH^- ions	1
	the hydroxide ions can receive / accept H^+ ions / protons	1
1(c)(ii)	Calculation of Na_2O moles $3.10 \text{ g} / 62$ OR 0.05	1
	Calculation of $[\text{OH}^-]$ $0.05 \times (2 / 0.400) = 0.25 \text{ mol dm}^{-3}$	1
	Calculation of pH $-\log 0.25 = 0.60$ $14 - 0.60 = 13.40$	1
1(d)	use of (2×109) or 218 and (2×494) or 988	1
	use of (0.5×496) or 248	1
	use of 416, 142, 844	1
	evaluation of expression correctly $\Delta H_{\text{lat}} = -416 - (2 \times 109) - (0.5 \times 496) - (2 \times 494) - (-142 + 844) = -2572$	1
1(e)	the lattice energy of Na_2S is less exothermic	1
	the sulfide ion is larger than the oxide ion / S^{2-} larger than O^{2-} / ionic radii quoted 0.184 nm and 0.140 nm AND less attraction (between the ions)/bonds are weaker	1

Question	Answer	Marks
2(a)	change in amount / mass / concentration of reactant / product per time	1
2(b)	decrease in volume or pressure	1
2(c)	$8.13 \times 10^4 / 81280 / 81300$	1
	$\text{mol}^{-2} \text{dm}^6 \text{s}^{-1}$	1
2(d)	$\sqrt{(0.00231 / (0.0046 \times 81280))} = 2.49 \times 10^{-3}$	1
2(e)	2, 1, 3	1
2(f)(i)	2	1
2(f)(ii)	the total of steps 1 and 2 / the components of 2 are two NO and one H ₂	1
2(g)(i)	time for amount or mass or concentration to halve	1
2(g)(ii)	0.02 at start and 0.01 after 2 seconds	1
	0.005 after 4 seconds and 0.0025 after 6 seconds	1
2(h)(i)	NO + $\frac{1}{2}$ O ₂ → NO ₂ or NO + O ₂ → NO ₂ + $\frac{1}{2}$ O ₂ AND NO ₂ + SO ₂ → NO + SO ₃	1
2(h)(ii)	(NO is) regenerated / reformed	1
2(h)(iii)	SO ₃ + H ₂ O → H ₂ SO ₄ AND acid rain or consequence of this described	1



PUBLISHED




Question	Answer	Marks												
3(a)	<table border="1"> <tr> <td></td><td>anode</td><td>cathode</td></tr> <tr> <td>AgNO₃ (aq)</td><td>oxygen / O₂</td><td>silver / Ag</td></tr> <tr> <td>saturated NaCl (aq)</td><td>chlorine / Cl₂</td><td>hydrogen / H₂</td></tr> <tr> <td>CuSO₄ (aq)</td><td>oxygen / O₂</td><td>copper / Cu</td></tr> </table>		anode	cathode	AgNO ₃ (aq)	oxygen / O ₂	silver / Ag	saturated NaCl (aq)	chlorine / Cl ₂	hydrogen / H ₂	CuSO ₄ (aq)	oxygen / O ₂	copper / Cu	3
	anode	cathode												
AgNO ₃ (aq)	oxygen / O ₂	silver / Ag												
saturated NaCl (aq)	chlorine / Cl ₂	hydrogen / H ₂												
CuSO ₄ (aq)	oxygen / O ₂	copper / Cu												
3(b)(i)	2I ⁻ → I ₂ + 2e ⁻	1												
	Ca ²⁺ + 2e ⁻ → Ca	1												
3(b)(ii)	- Ca / Calcium reduced and I / iodine oxidised - Oxidation number of calcium decreases from 2 to 0 - Oxidation number of iodine increases from -1 to 0 2 points = 1 mark 3 points = 2 marks	2												
3(b)(iii)	- metal / grey / silvery - purple AND vapour / gas / fumes - amount of melt decreases any 2 points for 1 mark	1												
3(c)	2 × 60 × 60 × 0.8 = 5760 C AND 5760 / 96500 = 0.060 (0.0597) F	1												
	1.11 / 55.8 = 0.020 (0.0199) mol of Fe	1												
	0.06 / 0.02 = 3 ∴ Fe ³⁺ or +3 or 3	1												


Question	Answer	Marks
4(a)	calcium – red flame	1
	barium – green flame	1
4(b)	- the temperature increases down the group - ionic radius increases / charge density decreases down the group - decreasing distortion / polarisation or decreasing weakening of bonds - of the anion / the CO_3^{2-} ion 2 points = 1 mark 3 points = 2 marks 4 points = 3 marks	3
4(c)	the gas is colourless / looks like air / cannot be seen	1
	appearance of solid doesn't change / white solid becomes white solid	1



Question	Answer	Marks
5(a)	$3s^2 3p^6 3d^{10} 4s^1$	1
	$3s^2 3p^6 3d^{10}$	1
5(b)	Cl-Cu-Cl	1
	one minus charge	1
5(c)		1
	octahedral and 90° or 180° labelled correctly on diagram as appropriate	1

Question	Answer	Marks
5(d)	reaction 1: blue ppt / blue solid	1
	$[\text{Cu}(\text{H}_2\text{O})_6]^{2+} + 2\text{OH}^- \rightarrow \text{Cu}(\text{OH})_2 + 6\text{H}_2\text{O}$ or $[\text{Cu}(\text{H}_2\text{O})_6]^{2+} + 2\text{OH}^- \rightarrow \text{Cu}(\text{OH})_2(\text{H}_2\text{O})_4 + 2\text{H}_2\text{O}$ or $[\text{Cu}(\text{H}_2\text{O})_6]^{2+} + 2\text{NH}_3 \rightarrow \text{Cu}(\text{OH})_2(\text{H}_2\text{O})_4 + 2\text{NH}_4^+$	1
	reaction 2: deep / dark / royal blue solution	1
	$\text{Cu}(\text{OH})_2 + 4\text{NH}_3 + 2\text{H}_2\text{O} \rightarrow [\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+} + 2\text{OH}^-$ or $\text{Cu}(\text{OH})_2 + 4\text{NH}_3 \rightarrow [\text{Cu}(\text{NH}_3)_4]^{2+} + 2\text{OH}^-$ or $\text{Cu}(\text{OH})_2(\text{H}_2\text{O})_4 + 4\text{NH}_3 \rightarrow [\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+} + 2\text{OH}^- + 2\text{H}_2\text{O}$	1
5(e)(i)	ligand exchange / displacement / replacement / substitution	1
5(e)(ii)	$K_{\text{stab}} = \frac{[\text{CuEDTA}^{2-}]}{[\text{Cu}(\text{H}_2\text{O})_6]^{2+}][\text{EDTA}^{4-}]}$	1
5(e)(iii)	stable / more stable than $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$	1
5(f)(i)	donates lone pairs / forms dative / co-ordinate bonds to (central) metal atom / metal ion	1
	donates two lone pairs / forms two (dative or coordinate) bonds	1
5(f)(ii)	$[\text{Zr}(\text{C}_2\text{O}_4)_4]^{4-}$	1
	not octahedral because 8 dative bonds to Zr or not octahedral because not 6 dative bonds to Zr or not octahedral because co-ordination number is 8 / is not 6	1


Question	Answer	Marks
6(a)(i)	D 2-chloropropane	1
	E hydrogen chloride	1
6(a)(ii)	(Friedel-Crafts) alkylation	1
6(b)(i)	$AlCl_3$ or $FeCl_3$	1
6(b)(ii)		1
6(b)(iii)	sunlight or UV OR $T > 100\text{ }^{\circ}\text{C}$	1
6(b)(iv)		1

Question	Answer	Marks
6(c)	reaction with hot $\text{KMnO}_4(\text{aq})$ 	1
	reaction with $\text{H}_2 + \text{Ni}$, high pressure 	1
6(d)		
	attacking species is NO_2^+	1
	curly arrow starting within hexagon and going to NO_2^+	1
	correct intermediate	1
	2nd curly arrow from C-H bond into ring	1

Question	Answer	Marks
7(a)	$2\text{C}_3\text{H}_7\text{OH} + 2\text{Na} \rightarrow 2\text{C}_3\text{H}_7\text{ONa} + \text{H}_2$	1
7(b)(i)	propanoic acid, phenol, propan-1-ol	1
7(b)(ii)	- propan-1-ol: O-H bond strengthened by positive inductive effect of alkyl group OR propoxide ion is destabilised by positive inductive effect of alkyl group - phenol: O-H bond weakened by negative inductive effect of ring OR phenoxide ion is stabilised by delocalisation of oxygen lone pair into ring - propanoic acid: O-H bond weakened by negative inductive effect of C=O OR propanoate ion is stabilised by delocalisation of minus charge by C=O 1 mark for a correct explanation, max 2 marks	2
7(c)	Tollens' reagent or Fehling's reagent	1
	methanoic acid gives a silver mirror/solid with Tollen's reagent OR red / orange ppt / solid with Fehlings' reagent	1
7(d)	PCl_5 or PCl_3 (+heat) or SOCl_2 (added to propanoic acid)	1
	product of first step: 	1
	add product of first step to phenol in NaOH	1
7(e)(i)	propanoic acid	1
7(e)(ii)	propan-1-ol would have peak at 0.5–6.0 because of OH group	1
	propanal would have peak at 9.3–10.5 because of CHO / aldehyde	1

Question	Answer	Marks
8(a)	correct chiral centre labelled only  $C_{15}H_{22}O_4$	1
8(b)	$C_{15}H_{22}O_4$	1
8(c)(i)		1
8(c)(ii)	CO_2	1
	oxidation / oxidative cleavage	1
8(c)(iii)	CH_3COCO_2H	1

PUBLISHED

Question	Answer	Marks
9(a)	$\text{C}_8\text{H}_{11}\text{O}_3\text{N}$	1
9(b)	yes, as it has a chiral C atom	1
9(c)(i)	(phenyldiazonium ion is stabilised because) positive charge is delocalised by ring / positive charge is spread over ring	1
9(c)(ii)		1
	N_2	1