



Cambridge International AS & A Level

CHEMISTRY

9701/51

Paper 5 Planning, Analysis and Evaluation

May/June 2023

MARK SCHEME

Maximum Mark: 30

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2023 series for most Cambridge IGCSE, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

This document consists of **8** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Science-Specific Marking Principles

1 Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.

2 The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.

3 Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).

4 The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.

5 'List rule' guidance

For questions that require ***n*** responses (e.g. State **two** reasons ...):

- The response should be read as continuous prose, even when numbered answer spaces are provided.
- Any response marked *ignore* in the mark scheme should not count towards ***n***.
- Incorrect responses should not be awarded credit but will still count towards ***n***.
- Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should **not** be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response.
- Non-contradictory responses after the first ***n*** responses may be ignored even if they include incorrect science.

6 Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form (e.g. $a \times 10^n$) in which the convention of restricting the value of the coefficient (a) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

7 Guidance for chemical equations

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.

Question	Answer	Marks
1(a)(i)	aqueous AND water has greater density (than ethoxyethane so it forms the lower layer)	1
1(a)(ii)	(10 cm ³) (volumetric) pipette	1
1(a)(iii)	the acid transfers / dissolves into the water (before titration / reaction with sodium hydroxide)	1
1(b)(i)	$n(\text{NaOH}) \text{ in titre} = \frac{27.25}{1000} \times 0.5 (= 0.0136250)$ $n(\text{HOOCCH}_2\text{CH}_2\text{COOH(aq)}) = 0.0136250 / 2 = 0.0068125$ $[\text{HOOCCH}_2\text{CH}_2\text{COOH(aq)}] = \left(\left(\frac{27.25}{1000} \times 0.5 \right) \div 2 \right) \times \frac{1000}{10} = 0.6813 \text{ mol dm}^{-3} (0.68125 \text{ mol dm}^{-3})$ at least 2 significant figures	1
1(b)(ii)	M1 $[\text{HOOCCH}_2\text{CH}_2\text{COOH(ethoxyethane)}] = 0.1125 \text{ mol dm}^{-3}$ $(22.50 \times 0.1 / 1000 \div 10 / 1000) / 2$ M2 $K_{\text{pc}} = M1 / (b)(i)$ Expected answer = $0.1125 / 0.68125 = 0.165(1376)$ $(0.1614\text{--}0.1662)$ at least 2 significant figures	2
1(b)(iii)	it is not possible to pipette all (30 cm ³) of the aqueous layer from the beaker (as some will always be left behind)	1
1(b)(iv)	use a larger volume (>30cm ³) of water (in step 1)	1
1(b)(v)	K_{pc} value would be smaller / decrease AND because (equilibrium had not been attained and) more butanedioic acid remains in aqueous layer	1

PUBLISHED

Question	Answer	Marks
2(a)	the application of the hydrolysed samples (to the paper) is more controlled	1
2(b)	to make the amino acids / spots visible	1
2(c)	lysine, tryptophan and leucine two correct for one mark, three correct for two marks	2
2(d)	two (of the three) amino acids (that compose the tripeptide) are the same	1
2(e)(i)	(the spots of two amino acids) have similar (not the same) R_f values	1
2(e)(ii)	use a longer stationary phase / (chromatography) paper OR use a different solvent / mobile phase	1

Question	Answer	Marks																											
3(a)	M1 add (a small volume of) distilled water (to the 50 cm³ beaker) AND dissolve the (aromatic diazonium) compound / solid M2 transfer the solution and washings (using a funnel) AND into a 100 cm³ volumetric flask M3 make up to the (calibration) mark / line with (distilled) water AND then mix the solution (by inverting the flask)	3																											
3(b)(i)	thermostatically controlled water bath	1																											
3(b)(ii)	volume of (nitrogen) gas (produced)	1																											
3(b)(iii)	not possible to weigh apparatus (while temperature is maintained) in water bath	1																											
3(c)(i)	<table border="1"> <thead> <tr> <th>time / min</th><th>volume, V_t / cm³</th><th>$V_{\text{final}} - V_t$ / cm³</th></tr> </thead> <tbody> <tr> <td>0</td><td>0.0</td><td>57.2</td></tr> <tr> <td>5</td><td>17.3</td><td>39.9</td></tr> <tr> <td>9</td><td>27.0</td><td>30.2</td></tr> <tr> <td>16</td><td>39.5</td><td>17.7</td></tr> <tr> <td>21</td><td>42.6</td><td>14.6</td></tr> <tr> <td>28</td><td>49.0</td><td>8.2</td></tr> <tr> <td>36</td><td>52.8</td><td>4.4</td></tr> <tr> <td>final</td><td>57.2</td><td>0.0</td></tr> </tbody> </table>	time / min	volume, V_t / cm ³	$V_{\text{final}} - V_t$ / cm ³	0	0.0	57.2	5	17.3	39.9	9	27.0	30.2	16	39.5	17.7	21	42.6	14.6	28	49.0	8.2	36	52.8	4.4	final	57.2	0.0	1
time / min	volume, V_t / cm ³	$V_{\text{final}} - V_t$ / cm ³																											
0	0.0	57.2																											
5	17.3	39.9																											
9	27.0	30.2																											
16	39.5	17.7																											
21	42.6	14.6																											
28	49.0	8.2																											
36	52.8	4.4																											
final	57.2	0.0																											

PUBLISHED

Question	Answer	Marks
3(c)(ii)	M1 all points plotted correctly M2 smooth curve of best fit line drawn passing close to all points except marked anomaly	2
3(c)(iii)	point most anomalous to the plotted line of best fit circled.	1
3(c)(iv)	(for anomalous point above the line of best fit) measurement (of volume) was made before the time recorded in the results table (for anomalous point below the line of best fit) measurement (of volume) was made after the time recorded in the results table	1
3(c)(v)	M1 two sets of co-ordinates from the line of best fit correctly recorded for first $t_{1/2}$ AND second $t_{1/2}$ in the form (x, y) M2 both half-lives correctly calculated from co-ordinates	2
3(c)(vi)	first order AND half-lives are constant (within experimental error)	1



Cambridge International AS & A Level

CHEMISTRY

9701/52

Paper 5 Planning, Analysis and Evaluation

May/June 2023

MARK SCHEME

Maximum Mark: 30

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2023 series for most Cambridge IGCSE, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

This document consists of 7 printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

PUBLISHED**GENERIC MARKING PRINCIPLE 5:**

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Science-Specific Marking Principles

1	Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.
2	The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.
3	Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).
4	The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.
5	<u>'List rule' guidance</u> For questions that require <i>n</i> responses (e.g. State two reasons ...): • The response should be read as continuous prose, even when numbered answer spaces are provided. • Any response marked <i>ignore</i> in the mark scheme should not count towards <i>n</i>. • Incorrect responses should not be awarded credit but will still count towards <i>n</i>. • Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should not be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response. • Non-contradictory responses after the first <i>n</i> responses may be ignored even if they include incorrect science.

6 Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form (e.g. $a \times 10^n$) in which the convention of restricting the value of the coefficient (a) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

7 Guidance for chemical equations

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.

Question	Answer	Marks
1(a)(i)	$1.242 \text{ g} / 1.08 = 1.15 \text{ cm}^3$	1
1(a)(ii)	burette	1
1(a)(iii)	$1 \times 100 / 1000 \times 40 = 4.0(0) \text{ g}$	1
1(b)	M1 dissolve the solid / sodium hydroxide (in the beaker) (using a small volume of) (distilled) water M2 transfer / add to a (100 cm ³ volumetric) flask AND rinse (with distilled water) M3 top the 100 cm ³ volumetric flask up to mark (with distilled water)	3
1(c)	M1 mol C ₆ H ₅ COOCH ₃ = $1.242 / 136 = 0.00913$ AND mol of NaOH = $0.100 \times (10 / 1000) = 0.001(00)$ M2 mass of benzoic acid decreases AND NaOH no longer in excess / NaOH (now) the limiting (reagent) / not all C ₆ H ₅ COOCH ₃ reacts / statement that mol NaOH < mol ester	2
1(d)(i)	allows heating to increase the rate of reaction without the loss of volatile substances	1
1(d)(ii)	(substances) are flammable	1
1(d)(iii)	to remove the anti-bumping granules (from the reaction mixture)	1
1(e)(i)	add (a little) (named) (acid–base) indicator AND the reaction mixture shows the colour for a (strong) acid in that indicator	1
1(e)(ii)	precipitate forms	1
1(f)	benzoic acid is less soluble in cold water	1
1(g)	dry the precipitate and reweigh until constant mass	1
1(h)(i)	$1.242 \times 122 / 136 = 1.11$	1
1(h)(ii)	74.0(%)	1

Question	Answer	Marks
2(a)(i)	$1.995 \times 10^{-7} \text{ (mol dm}^{-3}\text{) or } 2.00 \times 10^{-7} \text{ (mol dm}^{-3}\text{)}$	1

Question	Answer	Marks																																
2(a)(ii)	$3.98 \times 10^{-14} \text{ (mol}^2 \text{ dm}^{-6}\text{)}$	1																																
2(a)(iii)	as temperature increases K_w increases	1																																
2(b)(i)	M1 $1/T$ all correct to 3 significant figures AND standard form M2 $\log K_w$ all correct to 2 decimal places <table><tr><th>T</th><th>$1 / T$</th><th>K_w</th><th>$\log K_w$</th></tr><tr><td>283</td><td>3.53×10^{-3}</td><td>2.93×10^{-15}</td><td>– 14.53</td></tr><tr><td>293</td><td>3.41×10^{-3}</td><td>6.81×10^{-15}</td><td>– 14.17</td></tr><tr><td>303</td><td>3.30×10^{-3}</td><td>1.47×10^{-14}</td><td>– 13.83</td></tr><tr><td>313</td><td>3.19×10^{-3}</td><td>2.92×10^{-14}</td><td>– 13.53</td></tr><tr><td>323</td><td>3.10×10^{-3}</td><td>5.48×10^{-14}</td><td>– 13.26</td></tr><tr><td>333</td><td>3.00×10^{-3}</td><td>1.09×10^{-13}</td><td>– 12.96</td></tr><tr><td>343</td><td>2.92×10^{-3}</td><td>1.45×10^{-13}</td><td>– 12.84</td></tr></table>	T	$1 / T$	K_w	$\log K_w$	283	3.53×10^{-3}	2.93×10^{-15}	– 14.53	293	3.41×10^{-3}	6.81×10^{-15}	– 14.17	303	3.30×10^{-3}	1.47×10^{-14}	– 13.83	313	3.19×10^{-3}	2.92×10^{-14}	– 13.53	323	3.10×10^{-3}	5.48×10^{-14}	– 13.26	333	3.00×10^{-3}	1.09×10^{-13}	– 12.96	343	2.92×10^{-3}	1.45×10^{-13}	– 12.84	2
T	$1 / T$	K_w	$\log K_w$																															
283	3.53×10^{-3}	2.93×10^{-15}	– 14.53																															
293	3.41×10^{-3}	6.81×10^{-15}	– 14.17																															
303	3.30×10^{-3}	1.47×10^{-14}	– 13.83																															
313	3.19×10^{-3}	2.92×10^{-14}	– 13.53																															
323	3.10×10^{-3}	5.48×10^{-14}	– 13.26																															
333	3.00×10^{-3}	1.09×10^{-13}	– 12.96																															
343	2.92×10^{-3}	1.45×10^{-13}	– 12.84																															
2(b)(ii)	M1 seven points plotted correctly M2 line of best fit drawn	2																																
2(b)(iii)	plot at 2.92×10^{-3} , –12.84	1																																
2(b)(iv)	pH was recorded at a lower temperature than 343 K	1																																
2(b)(v)	M1 co-ordinates read and recorded correctly M2 gradient calculated to three significant figures	2																																
2(b)(vi)	M1 gradient = $-\Delta H / 2.303R$ OR $\Delta H = -\text{gradient} \times 2.303 \times 8.31$ OR any correct rearrangement of this M2 correct calculation of ΔH from 2(b)(v)	2																																



Cambridge International AS & A Level

CHEMISTRY

9701/53

Paper 5 Planning, Analysis and Evaluation

May/June 2023

MARK SCHEME

Maximum Mark: 30

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2023 series for most Cambridge IGCSE, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

This document consists of **8** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Science-Specific Marking Principles

1 Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.

2 The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.

3 Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).

4 The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.

5 'List rule' guidance

For questions that require ***n*** responses (e.g. State **two** reasons ...):

- The response should be read as continuous prose, even when numbered answer spaces are provided.
- Any response marked *ignore* in the mark scheme should not count towards ***n***.
- Incorrect responses should not be awarded credit but will still count towards ***n***.
- Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should **not** be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response.
- Non-contradictory responses after the first ***n*** responses may be ignored even if they include incorrect science.

6 Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form (e.g. $a \times 10^n$) in which the convention of restricting the value of the coefficient (a) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

7 Guidance for chemical equations

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.

Question	Answer	Marks
1(a)(i)	aqueous AND water has greater density (than ethoxyethane so it forms the lower layer)	1
1(a)(ii)	(10 cm ³) (volumetric) pipette	1
1(a)(iii)	the acid transfers / dissolves into the water (before titration / reaction with sodium hydroxide)	1
1(b)(i)	$n(\text{NaOH}) \text{ in titre} = \frac{27.25}{1000} \times 0.5 (= 0.0136250)$ $n(\text{HOOCCH}_2\text{CH}_2\text{COOH(aq)}) = 0.0136250 / 2 = 0.0068125$ $[\text{HOOCCH}_2\text{CH}_2\text{COOH(aq)}] = \left(\left(\frac{27.25}{1000} \times 0.5 \right) \div 2 \right) \times \frac{1000}{10} = 0.6813 \text{ mol dm}^{-3} (0.68125 \text{ mol dm}^{-3})$ at least 2 significant figures	1
1(b)(ii)	M1 $[\text{HOOCCH}_2\text{CH}_2\text{COOH(ethoxyethane)}] = 0.1125 \text{ mol dm}^{-3}$ $(22.50 \times 0.1 / 1000 \div 10 / 1000) / 2$ M2 $K_{\text{pc}} = M1 / (b)(i)$ Expected answer = $0.1125 / 0.68125 = 0.165(1376)$ $(0.1614\text{--}0.1662)$ at least 2 significant figures	2
1(b)(iii)	it is not possible to pipette all (30 cm ³) of the aqueous layer from the beaker (as some will always be left behind)	1
1(b)(iv)	use a larger volume (>30cm ³) of water (in step 1)	1
1(b)(v)	K_{pc} value would be smaller / decrease AND because (equilibrium had not been attained and) more butanedioic acid remains in aqueous layer	1

PUBLISHED

Question	Answer	Marks
2(a)	the application of the hydrolysed samples (to the paper) is more controlled	1
2(b)	to make the amino acids / spots visible	1
2(c)	lysine, tryptophan and leucine two correct for one mark, three correct for two marks	2
2(d)	two (of the three) amino acids (that compose the tripeptide) are the same	1
2(e)(i)	(the spots of two amino acids) have similar (not the same) R_f values	1
2(e)(ii)	use a longer stationary phase / (chromatography) paper OR use a different solvent / mobile phase	1

Question	Answer	Marks																											
3(a)	M1 add (a small volume of) distilled water (to the 50 cm³ beaker) AND dissolve the (aromatic diazonium) compound / solid M2 transfer the solution and washings (using a funnel) AND into a 100 cm³ volumetric flask M3 make up to the (calibration) mark / line with (distilled) water AND then mix the solution (by inverting the flask)	3																											
3(b)(i)	thermostatically controlled water bath	1																											
3(b)(ii)	volume of (nitrogen) gas (produced)	1																											
3(b)(iii)	not possible to weigh apparatus (while temperature is maintained) in water bath	1																											
3(c)(i)	<table border="1"> <thead> <tr> <th>time / min</th><th>volume, V_t / cm³</th><th>$V_{\text{final}} - V_t$ / cm³</th></tr> </thead> <tbody> <tr> <td>0</td><td>0.0</td><td>57.2</td></tr> <tr> <td>5</td><td>17.3</td><td>39.9</td></tr> <tr> <td>9</td><td>27.0</td><td>30.2</td></tr> <tr> <td>16</td><td>39.5</td><td>17.7</td></tr> <tr> <td>21</td><td>42.6</td><td>14.6</td></tr> <tr> <td>28</td><td>49.0</td><td>8.2</td></tr> <tr> <td>36</td><td>52.8</td><td>4.4</td></tr> <tr> <td>final</td><td>57.2</td><td>0.0</td></tr> </tbody> </table>	time / min	volume, V_t / cm ³	$V_{\text{final}} - V_t$ / cm ³	0	0.0	57.2	5	17.3	39.9	9	27.0	30.2	16	39.5	17.7	21	42.6	14.6	28	49.0	8.2	36	52.8	4.4	final	57.2	0.0	1
time / min	volume, V_t / cm ³	$V_{\text{final}} - V_t$ / cm ³																											
0	0.0	57.2																											
5	17.3	39.9																											
9	27.0	30.2																											
16	39.5	17.7																											
21	42.6	14.6																											
28	49.0	8.2																											
36	52.8	4.4																											
final	57.2	0.0																											

PUBLISHED

Question	Answer	Marks
3(c)(ii)	M1 all points plotted correctly M2 smooth curve of best fit line drawn passing close to all points except marked anomaly	2
3(c)(iii)	point most anomalous to the plotted line of best fit circled.	1
3(c)(iv)	(for anomalous point above the line of best fit) measurement (of volume) was made before the time recorded in the results table (for anomalous point below the line of best fit) measurement (of volume) was made after the time recorded in the results table	1
3(c)(v)	M1 two sets of co-ordinates from the line of best fit correctly recorded for first $t_{1/2}$ AND second $t_{1/2}$ in the form (x, y) M2 both half-lives correctly calculated from co-ordinates	2
3(c)(vi)	first order AND half-lives are constant (within experimental error)	1