
CHEMISTRY

9701/41

Paper 4 A Level Structured Questions

October/November 2018

MARK SCHEME

Maximum Mark: 100

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2018 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

This document consists of **14** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:


Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

PUBLISHED**GENERIC MARKING PRINCIPLE 5:**


Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Question	Answer	Marks												
1(a)(i)	<table border="1"> <thead> <tr> <th>peak</th><th>organic compound</th><th>explanation</th></tr> </thead> <tbody> <tr> <td>X</td><td>alkane</td><td>London forces only OR no hydrogen bonding</td></tr> <tr> <td>Y</td><td>aldehyde</td><td>(Permanent dipole-dipole and London forces)</td></tr> <tr> <td>Z</td><td>carboxylic acid</td><td>(contains) hydrogen bonding</td></tr> </tbody> </table> M1 peak assignments [1] M2 explanation of Z OR X [1]	peak	organic compound	explanation	X	alkane	London forces only OR no hydrogen bonding	Y	aldehyde	(Permanent dipole-dipole and London forces)	Z	carboxylic acid	(contains) hydrogen bonding	2
peak	organic compound	explanation												
X	alkane	London forces only OR no hydrogen bonding												
Y	aldehyde	(Permanent dipole-dipole and London forces)												
Z	carboxylic acid	(contains) hydrogen bonding												
1(a)(ii)	% of Z = $47/98 = 48\%$	1												
1(b)(i)	^{37}Cl and ^{81}Br	1												
1(b)(ii)	M peak $\text{CH}_2^{35}\text{Cl}^{79}\text{Br}$ M+2 peak $\text{CH}_2^{37}\text{Cl}^{79}\text{Br}$ OR $\text{CH}_2^{35}\text{Cl}^{81}\text{Br}$ M+4 peak $\text{CH}_2^{37}\text{Cl}^{81}\text{Br}$ two correct scores 1 mark all 3 correct scores 2 marks	2												
1(c)(i)	 M1 correct dipole on HBr AND any correct curly arrow [1] M2 two other correct curly arrows AND lone pair required on Br^- [1] M3 intermediate [1]	3												



Question	Answer	Marks
1(c)(ii)	(major product is) formed via the most stable tertiary carbocation / intermediate OR tertiary halogenoalkane formed via more stable carbocation / intermediate	1
1(d)(i)	M1 ratio of the concentrations of solute in two (immiscible) solvents [1] M2 at equilibrium [1]	2
1(d)(ii)	$K_{\text{partition}} = (x/10)/(1.25-x/50)$ [1] $4.75(1.25-x) = 5x$ $x = 5.9375/9.75 = \mathbf{0.61\text{ g}}$ [1] correct answer scores [2]	2

Question	Answer	Marks
2(a)	species that forms dative bond(s) to a (central) metal atom / ion	1
2(b)	 any two structures [1] × 2	2
2(c)(i)	$K_{\text{sp}} = [\text{Ca}^{2+}][\text{C}_2\text{O}_4^{2-}]$ [1] units $\text{mol}^2 \text{dm}^{-6}$ [1]	2
2(c)(ii)	$[\text{Ca}^{2+}] = [\text{C}_2\text{O}_4^{2-}] = 6.65 \times 10^{-3}/128.1 = 5.19 \times 10^{-5} \text{ mol dm}^{-3}$ [1] $K_{\text{sp}} = (5.19 \times 10^{-5})^2 = \mathbf{2.7 \times 10^{-9} \text{ mol}^2 \text{dm}^{-6}}$ [1]	2

PUBLISHED

Question	Answer	Marks															
3(a)	[1] for each column <table border="1"> <tr> <td></td><td colspan="2">number of unpaired electrons in</td></tr> <tr> <td>element</td><td>3d</td><td>4s</td></tr> <tr> <td>Cr</td><td>5</td><td>1</td></tr> <tr> <td>Mn</td><td>5</td><td>0</td></tr> <tr> <td>Fe</td><td>4</td><td>0</td></tr> </table>		number of unpaired electrons in		element	3d	4s	Cr	5	1	Mn	5	0	Fe	4	0	2
	number of unpaired electrons in																
element	3d	4s															
Cr	5	1															
Mn	5	0															
Fe	4	0															
3(b)	$2\text{KMnO}_4 \rightarrow \text{K}_2\text{MnO}_4 + \text{O}_2 + \text{MnO}_2$ formulae of K_2MnO_4 and O_2 [1] rest of the equation [1]	2															
3(c)	M1 d orbitals split into two levels / lower and upper orbitals [1] M2 visible light is absorbed and the complementary colour observed [1] M3 electron(s) promoted / excited [1]	3															
3(d)(i)	precipitate A $[\text{Cu}(\text{H}_2\text{O})_4(\text{OH})_2]$ OR $\text{Cu}(\text{OH})_2$ [1] solution B $[\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}$ [1] solution C $\text{Cu}(\text{CH}_3\text{CO}_2)_2$ [1]	3															
3(d)(ii)	Na_2CO_3 or CO_3^{2-}	1															
3(d)(iii)	$\text{CuCO}_3 + 2\text{CH}_3\text{CO}_2\text{H} \rightarrow \text{Cu}(\text{CH}_3\text{CO}_2)_2 + \text{CO}_2 + \text{H}_2\text{O}$	1															
3(d)(iv)	any two for one mark - fizzing / bubbles / effervescence - solid disappears - green / blue solution (formed)	1															


PUBLISHED


Question	Answer	Marks
3(e)	sum of the charges of the (four) ligands equals the oxidation number / charge of Pt OR a calculation Pt +2, NH ₃ neutral / no charge, both Cl ⁻ 's -1 (so no overall charge)	1
3(f)(i)	 square planar and 180° [1]	2
3(f)(ii)	M1 this can bond / bind with DNA [1] M2 which prevents replication of the DNA / strand OR prevents cell division [1]	2
3(g)		1


Question	Answer	Marks
4(a)	M1 solubility decreases (down the Group) [1] M2 because lattice energy and hydration energy decreases OR lattice energy and hydration energy become less exothermic / more endothermic [1] M3 because hydration energy decreases to a greater extent (than does ΔH_{Latt}) [1]	3
4(b)(i)	$(K_w =) [\text{H}^+][\text{OH}^-]$	1


PUBLISHED

Question	Answer	Marks												
4(b)(ii)	[1] or each correct tick <table><tr><td>effect of increasing temperature</td><td>decreases</td><td>stay the same</td><td>increase</td></tr><tr><td>pH</td><td>✓</td><td></td><td></td></tr><tr><td>ratio of [H⁺]:[OH⁻]</td><td></td><td>✓</td><td></td></tr></table>	effect of increasing temperature	decreases	stay the same	increase	pH	✓			ratio of [H ⁺]:[OH ⁻]		✓		2
effect of increasing temperature	decreases	stay the same	increase											
pH	✓													
ratio of [H ⁺]:[OH ⁻]		✓												
4(c)	[H⁺] = 10^{-13.25} = 5.62 × 10⁻¹⁴ [1] [OH⁻] = K_w/[H⁺] = 1.0 × 10⁻¹⁴/5.62 × 10⁻¹⁴ [OH⁻] = 0.18 (0.178) (mol dm⁻³) [1] ecf correct answer scores [2]	2												
4(d)	HCO₃⁻ + H⁺ → H₂CO₃ OR HCO₃⁻ + H⁺ → CO₂ + H₂O [1] H₂CO₃ + OH⁻ → HCO₃⁻ + H₂O [1]	2												
4(e)(i)	CH₃COOH + H₂O ⇌ CH₃COO⁻ + H₃O⁺ [1] acid + base ⇌ base + acid [1]	2												
4(e)(ii)	M1 moles NaOH = 0.15 × 20/1000 = 0.0030 AND initial moles CH₃COOH = 0.25 × 30/1000 OR 0.0075 [1] M2 equilibrium moles CH₃COOH = 0.0045 AND equilibrium moles CH₃COONa = 0.0030 [1] M3 [CH₃COOH] = 0.0045/0.05 = 0.090 AND [CH₃COONa] = 0.003/0.05 = 0.060 [H⁺] = K_a × [CH₃COOH]/[CH₃COONa] = 2.625 × 10⁻⁵ [1] M4 pH = -log[H⁺] = 4.6 [1] correct answer scores [4]	4												
4(f)(i)	end point = 28 cm³	1												
4(f)(ii)	M1 reaction M bromothymol (blue) / bromocresol (green) AND reaction N bromothymol (blue) / thymolphthalein [1] M2 (both indicators have) a pH range / colour change within / in end-point / vertical region / sharp fall of the graph [1]	2												





Question	Answer	Marks												
5(a)(i)	[1] for each correct answer <table><tr><td></td><td>number of peaks</td></tr><tr><td>F</td><td>3</td></tr><tr><td>G</td><td>6</td></tr></table>		number of peaks	F	3	G	6	2						
	number of peaks													
F	3													
G	6													
5(a)(ii)	 one amide bond displayed in full [1] rest of the structure – one repeat unit only [1]	2												
5(b)	[1] for each correct tick <table><tr><td></td><td>σ-bonds only</td><td>π-bonds only</td><td>both σ- and π-bonds</td></tr><tr><td>bonds broken</td><td></td><td>✓</td><td></td></tr><tr><td>bonds formed</td><td>✓</td><td></td><td></td></tr></table>		σ -bonds only	π -bonds only	both σ - and π -bonds	bonds broken		✓		bonds formed	✓			2
	σ -bonds only	π -bonds only	both σ - and π -bonds											
bonds broken		✓												
bonds formed	✓													

Question	Answer	Marks
5(c)	 M1 length of chain with both monomers [1] M2 continuation bonds [1]	2
5(d)(i)	C-C bonds are non-polar / have no dipole so cannot be hydrolysed [1]	1
5(d)(ii)	M1 <u>Hydrolysis</u> using acid / base / alkali / enzymes [1] M2 action of UV light [1]	2


Question	Answer	Marks
6(a)	 M1 amide bond displayed [1] M2 rest of the structure [1]	2

Question	Answer	Marks
6(b)	 M1 valine on the cross [1] M2 Val-Lys and Lys on the right of the cross (in any order) [1] M3 relative order of Val-Lys and Lys (on the same side of the cross) [1] Explanation - Val does not move as it is a zwitterion / neutral (at pH6) OR Lys / Val-Lys move towards negative (pole) as they are positively charged - Lys moves the furthest as it has the lowest M_r (with the same positive charge) OR Val-Lys moves the least as it has the largest M_r (with the same positive charge) [1] × 2	5

Question	Answer	Marks
7(a)	M1 C-X / C-Cl / C-O bond is stronger (in chlorobenzene / phenol) [1] M2 p-orbital / lone pair on Cl / O(H) / X (in chlorobenzene / phenol) [1] M3 electrons of the (Cl / O / electronegative atom) AND overlap / delocalise with π-electron cloud / delocalise into ring [1]	3

Question	Answer	Marks
7(b)	 2-bromophenol  4-bromophenol structure and name correct [1]	2
7(c)(i)	step 1 conc. HNO_3 + H_2SO_4 (and temperare 50–55 °C) [1] step 2 Sn + HCl AND one of conc.HCl + heat [1] step 4 H_2O warm / heat [1]	3
7(c)(ii)	 OR 	1
7(c)(iii)	step 1 electrophilic substitution	1
7(c)(iv)	$\text{C}_6\text{H}_5\text{NO}_2$ + $6[\text{H}] \rightarrow \text{C}_6\text{H}_5\text{NH}_2$ + $2\text{H}_2\text{O}$	1

PUBLISHED

Question	Answer	Marks									
8(a)	 M1 continuous increase in S from 0–300 K (excluding m.p.) [1] M2 steep vertical increase in S ONLY at the m.p. AND continuous increase in S after m.p. [1]	2									
8(b)	[1] for each correct tick <table border="1"> <thead> <tr> <th></th><th>negative $\Delta S^\ominus$</th><th>positive $\Delta S^\ominus$</th></tr> </thead> <tbody> <tr> <td>solid dissolving in water</td><td></td><td>✓</td></tr> <tr> <td>water boiling to steam</td><td></td><td>✓</td></tr> </tbody> </table>		negative $\Delta S^\ominus$	positive $\Delta S^\ominus$	solid dissolving in water		✓	water boiling to steam		✓	1
	negative $\Delta S^\ominus$	positive $\Delta S^\ominus$									
solid dissolving in water		✓									
water boiling to steam		✓									
8(c)	$\Delta H^\ominus = (2 \times \text{C=O}) + (3 \times \text{H-H}) - (3 \times \text{C-H}) - (\text{C-O}) - (3 \times \text{O-H})$ $\Delta H^\ominus = (2 \times 805) + (3 \times 436) - (3 \times 410) - (1 \times 360) - (3 \times 460)$ [1] $\Delta H^\ominus = 1610 + 1308 - 1230 - 360 - 1380 = -52 \text{ (kJ mol}^{-1}\text{)}$ [1] ecf correct answer scores [2]	2									
8(d)(i)	$\Delta S^\ominus = 127 + 70 - (214 + 3 \times 131)$ [1] $= -410 \text{ (J K}^{-1} \text{ mol}^{-1}\text{)}$ [1] ecf correct answer scores [2]	2									
8(d)(ii)	$\Delta G^\ominus = \Delta H^\ominus - T\Delta S^\ominus$ [1] $\Delta G^\ominus = -131 - (298 \times -0.41) = -8.8(2) \text{ (kJ mol}^{-1}\text{)}$ [1] correct answer scores [2]	2									

PUBLISHED

Question	Answer	Marks
8(d)(iii)	(as temperature increases) feasibility decreases	1
8(e)(i)	$2\text{CH}_3\text{OH} + 3\text{O}_2 \rightleftharpoons 2\text{CO}_2 + 4\text{H}_2\text{O}$ OR $2\text{CH}_3\text{OH} + 3\text{O}_2 \rightleftharpoons 2\text{CO}_2 + 4\text{H}^+ + 4\text{OH}^-$	1
8(e)(ii)	$E^\ominus_{\text{cell}} = 1.23 - 0.02 = \mathbf{1.21\text{ V}}$	1

CHEMISTRY

9701/42

Paper 4 A Level Structured Questions

October/November 2018

MARK SCHEME

Maximum Mark: 100

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2018 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

This document consists of **13** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.


GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).




GENERIC MARKING PRINCIPLE 6:


Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Question	Answer	Marks												
1(a)(i)	<table border="1"> <thead> <tr> <th>peak</th><th>organic compound</th><th>explanation</th></tr> </thead> <tbody> <tr> <td>J</td><td>carboxylic acid</td><td>most polar</td></tr> <tr> <td>K</td><td>ketone</td><td>polarity between J and L</td></tr> <tr> <td>L</td><td>alkene</td><td>most non-polar</td></tr> </tbody> </table> peak assignments [1] explanation of J or L [1]	peak	organic compound	explanation	J	carboxylic acid	most polar	K	ketone	polarity between J and L	L	alkene	most non-polar	2
peak	organic compound	explanation												
J	carboxylic acid	most polar												
K	ketone	polarity between J and L												
L	alkene	most non-polar												
1(a)(ii)	% of K = $18 / 92 = 19.6\%$	1												
1(b)(i)	$Cl_2 + AlCl_3 \rightarrow Cl^+ + AlCl_4^-$	1												
1(b)(ii)	intermediate [1] curly arrow [1]	4												
1(b)(iii)	$H^+ + AlCl_4^- \rightarrow AlCl_3 + HCl$	1												
1(c)(i)	catalyst and the reactants are in the same phase / state	1												
1(c)(ii)	(Rh) heterogeneous AND (Fe ³⁺) homogeneous	1												



Question	Answer	Marks
2(a)	3d Cu [Ar] ↑↓ ↑↓ ↑↓ ↑↓ ↑↓ Cu²⁺ [Ar] ↑↓ ↑↓ ↑↓ ↑↓ ↑ 4s ↑ [1] × 2	2
2(b)(i)	orbitals have the same energy	1
2(b)(ii)	d-d splitting seen, leading to 2 upper and 3 lower orbitals	1
2(c)	an ion / molecule that donates two pairs of electrons	1
2(d)	 one correct [1] two correct and mirror images of each other [1]	2



Question	Answer	Marks
3(a)(i)	solution of A $[\text{Co}(\text{NH}_3)_6]^{2+}$ [1] precipitate B CoCO_3 [1]	2
3(a)(ii)	$\text{NaOH}(\text{aq}) / \text{OH}^-(\text{aq})$	1
3(a)(iii)	$[\text{Co}(\text{H}_2\text{O})_4(\text{OH})_2] + 6\text{NH}_3 \rightarrow [\text{Co}(\text{NH}_3)_6]^{2+} + 4\text{H}_2\text{O} + 2\text{OH}^-$	1

Question	Answer	Marks
3(a)(iv)	$[\text{Co}(\text{H}_2\text{O})_6]^{2+} + \text{CO}_3^{2-} \rightarrow \text{CoCO}_3 + 6\text{H}_2\text{O}$	1
3(b)(i)	variable oxidation states	1
3(b)(ii)		1
3(b)(iii)	$\text{C}_4\text{H}_4\text{O}_6^{2-} + 3[\text{O}] \rightarrow 2\text{HCO}_2^- + 2\text{CO}_2 + \text{H}_2\text{O}$	1
3(c)(i)	 cis trans square planar shape of one isomer [1] both isomers drawn and assigned as cis and trans correctly [1]	2
3(c)(ii)	this can react / bond / bind with <u>DNA</u> [1] which prevents replication of the strand / prevents cell division / prevents mitosis [1]	2
3(d)		1

Question	Answer	Marks
4(a)	$\text{Ca}_3\text{N}_2 + 6\text{H}_2\text{O} \rightarrow 3\text{Ca}(\text{OH})_2 + 2\text{NH}_3$ products are $\text{Ca}(\text{OH})_2$ and NH_3 [1] rest of the equation, balanced [1]	2
4(b)	M1: solubility increases (down the Group) [1] M2: because lattice energy and hydration energy decreases or lattice energy and hydration energy become less exothermic / (more) endothermic [1] M3: because lattice energy decreases to a greater extent (than does ΔH_{hyd}) [1]	3
4(c)	 aqueous ions gaseous ions ΔH_{hyd} ΔH_{sol} ΔH_{latt} ionic solid arrow label and direction correct [1] x 3	3
4(d)(i)	$K_{\text{sp}} = [\text{Ca}^{2+}][\text{F}^-]^2$ [1] units = $\text{mol}^3 \text{dm}^{-9}$ [1]	2
4(d)(ii)	$K_{\text{sp}} = 4x^3 = 3.45 \times 10^{-11}$ $x = 2.05 \times 10^{-4} (\text{mol dm}^{-3})$	1


Question	Answer	Marks
5(a)	ionic radius / ion size increases OR charge density decreases (down the group) [1] less polarisation / distortion of anion / nitrate ion / NO_3^- / nitrate group OR N-O / N=O bond is less weakened / distorted / polarised OR more energy to break N-O / N=O bond [1]	2
5(b)	- moles of $\text{Ce}^{4+} = 0.0400 \times 21.8 / 1000 = 8.72 \times 10^{-4}$ (moles of Ce^{4+}) - moles of $\text{NO}_2^- = 8.72 \times 10^{-4} / 2 = 4.36 \times 10^{-4}$ in 25 cm^3 (use of 2:1 ratio correctly) - moles of $\text{NO}_2^- = 4.36 \times 10^{-4} \times 4 = 1.74(4) \times 10^{-3}$ in 100 cm^3 (use of 4:1 ratio correctly) - mass $\text{NaNO}_2 = 1.74(4) \times 10^{-3} \times (23.0 + 14.0 + 32.0) = 0.120 \text{ g}$ (use of M_r correctly) % purity = $0.120 / 0.138 = 86.96\%$ (use of 0.0138 correctly) two points = [1] four points = [2] all five points = [3]	3
5(c)(i)	$5\text{NO}_2^- + 2\text{MnO}_4^- + 6\text{H}^+ \rightarrow 2\text{Mn}^{2+} + 5\text{NO}_3^- + 3\text{H}_2\text{O}$ OR $5\text{HNO}_2 + 2\text{MnO}_4^- + \text{H}^+ \rightarrow 2\text{Mn}^{2+} + 5\text{NO}_3^- + 3\text{H}_2\text{O}$ all species correct [1] balanced [1]	2
5(c)(ii)	$E^\ominus_{\text{cell}} = 1.52 - 0.94 = \mathbf{0.58 \text{ (V)}}$	1
5(d)(i)	weak acid is partly ionised and strong acid is completely ionised	1
5(d)(ii)	$K_a = \frac{[\text{H}^+][\text{NO}_2^-]}{[\text{HNO}_2]}$	1
5(d)(iii)	$K_a = [\text{H}^+]^2 / [\text{HNO}_2]$ $[\text{H}^+] = \sqrt{0.00069 \times 0.15} = 1.02 \times 10^{-2}$ [1] $\text{pH} = -\log[\text{H}^+] = \mathbf{2.0 \text{ (1.99)}}$ [1] minimum 2 significant figures	2
5(d)(iv)	% ionisation = $100 \times 1.02 \times 10^{-2} / 0.15 = \mathbf{6.7-6.8 \%}$	1

Question	Answer	Marks						
5(e)(i)	M1 A solution that resists changes in pH [1] M2 when small amounts of acid or alkali are added to it [1]	2						
5(e)(ii)	M1 $\text{HNO}_2 + \text{OH}^- \rightarrow \text{NO}_2^- + \text{H}_2\text{O}$ [1] M2 $\text{NO}_2^- + \text{H}^+ \rightarrow \text{HNO}_2$ [1]	2						
5(f)(i)	CuCN / copper(I) cyanide	1						
5(f)(ii)	 Y  Z [1] × 2	2						
5(g)	<table border="1"><thead><tr><th></th><th>number of peaks</th></tr></thead><tbody><tr><td>W</td><td>6</td></tr><tr><td>Z</td><td>3</td></tr></tbody></table> [1] × 2		number of peaks	W	6	Z	3	2
	number of peaks							
W	6							
Z	3							

Question	Answer	Marks
6(a)(i)	KCN / NaCN / CN^-	1
6(a)(ii)	step 1 PCl_3 + heat / PCl_5 / SOCl_2 [1] step 4 NaBH_4 [1]	2
6(b)(i)		2
6(b)(ii)	step I condensation [1] step II reduction [1]	2
6(c)	 • amide bond (CO-NH) • structure of polymer with exactly two repeat units • continuation bonds • hydrocarbon portions correct two points = [1] four points = [1]	2

Question	Answer	Marks										
7(a)(i)	$C_{15}H_{10}N_2O_2$	1										
7(a)(ii)	-NH-CO-O- linkage [1] whole molecule correct [1]	2										
7(a)(iii)	<table><tr><td>intermolecular force</td><td>group(s) involved</td></tr><tr><td>hydrogen bonding</td><td>NH</td></tr><tr><td>VDW forces / Induced dipole-dipole forces / polar forces</td><td>-C₆H₄CH₂- allow benzene / aromatic rings</td></tr></table> M1 hydrogen bonding [1] M2 NH group for hydrogen AND second correct IMF [1]	intermolecular force	group(s) involved	hydrogen bonding	NH	VDW forces / Induced dipole-dipole forces / polar forces	-C ₆ H ₄ CH ₂ - allow benzene / aromatic rings	2				
intermolecular force	group(s) involved											
hydrogen bonding	NH											
VDW forces / Induced dipole-dipole forces / polar forces	-C ₆ H ₄ CH ₂ - allow benzene / aromatic rings											
7(b)	<table><tr><td>type of polymer</td><td>example</td></tr><tr><td>synthetic polyamide</td><td>nylon / Kevlar</td></tr><tr><td>synthetic polyester</td><td>Terylene</td></tr><tr><td>conducting polymer</td><td>polyacetylene / polyethyne</td></tr><tr><td>non-solvent based adhesive</td><td>epoxyresins / superglue</td></tr></table> one mark [1] for each correct answer up to a maximum of [3]	type of polymer	example	synthetic polyamide	nylon / Kevlar	synthetic polyester	Terylene	conducting polymer	polyacetylene / polyethyne	non-solvent based adhesive	epoxyresins / superglue	3
type of polymer	example											
synthetic polyamide	nylon / Kevlar											
synthetic polyester	Terylene											
conducting polymer	polyacetylene / polyethyne											
non-solvent based adhesive	epoxyresins / superglue											

PUBLISHED

Question	Answer	Marks
8(a)(i)	species with an unpaired electron	1
8(a)(ii)	$\text{NH}_2 + \text{Cl} \rightarrow \text{NH}_2\text{Cl}$	1
8(b)(i)		1
8(b)(ii)	sp^3 AND $100\text{--}107^\circ$	1
8(c)(i)	(entropy) is a measure of the disorder/randomness of a system	1
8(c)(ii)	$\Delta S^\circ = 237 + 187 - (241 + 198) = -15.0 \text{ (J K}^{-1} \text{ mol}^{-1}\text{)}$	1
8(c)(iii)	$\Delta H^\circ = 95.4 - 92.3 - (80.1 - 45.9) = -31.1 \text{ (kJ mol}^{-1}\text{)}$	1
8(c)(iv)	$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$ [1] $\Delta G^\circ = -31.1 - (298 \times -0.015) = -26.6 \text{ (kJ mol}^{-1}\text{)}$ [1]	2
8(c)(v)	(at higher temperatures) $T\Delta S^\circ$ becomes more negative so ΔG° becomes more positive OR (at high temperatures) $-T\Delta S^\circ$ is becomes more positive so ΔG° becomes more positive	1
8(d)	ethylamine > ammonia > phenylamine [1] ethyl group is electron donating group [1] p-orbital from N in phenylamine overlaps with π -ring system OR lone pair on N is delocalised into benzene ring [1] basicity linked to ability of N to accept a proton [1]	4

CHEMISTRY

9701/43

Paper 4 A Level Structured Questions

October/November 2018

MARK SCHEME

Maximum Mark: 100

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2018 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

This document consists of **14** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.


PUBLISHED**GENERIC MARKING PRINCIPLE 5:**

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).


GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

PUBLISHED

Question	Answer	Marks												
1(a)(i)	<table border="1"> <thead> <tr> <th>peak</th><th>organic compound</th><th>explanation</th></tr> </thead> <tbody> <tr> <td>X</td><td>alkane</td><td>London forces only OR no hydrogen bonding</td></tr> <tr> <td>Y</td><td>aldehyde</td><td>(Permanent dipole-dipole and London forces)</td></tr> <tr> <td>Z</td><td>carboxylic acid</td><td>(contains) hydrogen bonding</td></tr> </tbody> </table> M1 peak assignments [1] M2 explanation of Z OR X [1]	peak	organic compound	explanation	X	alkane	London forces only OR no hydrogen bonding	Y	aldehyde	(Permanent dipole-dipole and London forces)	Z	carboxylic acid	(contains) hydrogen bonding	2
peak	organic compound	explanation												
X	alkane	London forces only OR no hydrogen bonding												
Y	aldehyde	(Permanent dipole-dipole and London forces)												
Z	carboxylic acid	(contains) hydrogen bonding												
1(a)(ii)	% of Z = $47/98 = 48\%$	1												
1(b)(i)	^{37}Cl and ^{81}Br	1												
1(b)(ii)	M peak $\text{CH}_2^{35}\text{Cl}^{79}\text{Br}$ M+2 peak $\text{CH}_2^{37}\text{Cl}^{79}\text{Br}$ OR $\text{CH}_2^{35}\text{Cl}^{81}\text{Br}$ M+4 peak $\text{CH}_2^{37}\text{Cl}^{81}\text{Br}$ two correct scores 1 mark all 3 correct scores 2 marks	2												
1(c)(i)	 M1 correct dipole on HBr AND any correct curly arrow [1] M2 two other correct curly arrows AND lone pair required on Br^- [1] M3 intermediate [1]	3												



Question	Answer	Marks
1(c)(ii)	(major product is) formed via the most stable tertiary carbocation / intermediate OR tertiary halogenoalkane formed via more stable carbocation / intermediate	1
1(d)(i)	M1 ratio of the concentrations of solute in two (immiscible) solvents [1] M2 at equilibrium [1]	2
1(d)(ii)	$K_{\text{partition}} = (x/10)/(1.25-x/50)$ [1] $4.75(1.25-x) = 5x$ $x = 5.9375/9.75 = \mathbf{0.61\text{ g}}$ [1] correct answer scores [2]	2

Question	Answer	Marks
2(a)	species that forms dative bond(s) to a (central) metal atom / ion	1
2(b)	 any two structures [1] × 2	2
2(c)(i)	$K_{\text{sp}} = [\text{Ca}^{2+}][\text{C}_2\text{O}_4^{2-}]$ [1] units $\text{mol}^2 \text{dm}^{-6}$ [1]	2
2(c)(ii)	$[\text{Ca}^{2+}] = [\text{C}_2\text{O}_4^{2-}] = 6.65 \times 10^{-3}/128.1 = 5.19 \times 10^{-5} \text{ mol dm}^{-3}$ [1] $K_{\text{sp}} = (5.19 \times 10^{-5})^2 = \mathbf{2.7 \times 10^{-9} \text{ mol}^2 \text{dm}^{-6}}$ [1]	2

PUBLISHED

Question	Answer	Marks															
3(a)	[1] for each column <table border="1"> <tr> <td></td><td colspan="2">number of unpaired electrons in</td></tr> <tr> <td>element</td><td>3d</td><td>4s</td></tr> <tr> <td>Cr</td><td>5</td><td>1</td></tr> <tr> <td>Mn</td><td>5</td><td>0</td></tr> <tr> <td>Fe</td><td>4</td><td>0</td></tr> </table>		number of unpaired electrons in		element	3d	4s	Cr	5	1	Mn	5	0	Fe	4	0	2
	number of unpaired electrons in																
element	3d	4s															
Cr	5	1															
Mn	5	0															
Fe	4	0															
3(b)	$2\text{KMnO}_4 \rightarrow \text{K}_2\text{MnO}_4 + \text{O}_2 + \text{MnO}_2$ formulae of K_2MnO_4 and O_2 [1] rest of the equation [1]	2															
3(c)	M1 d orbitals split into two levels / lower and upper orbitals [1] M2 visible light is absorbed and the complementary colour observed [1] M3 electron(s) promoted / excited [1]	3															
3(d)(i)	precipitate A $[\text{Cu}(\text{H}_2\text{O})_4(\text{OH})_2]$ OR $\text{Cu}(\text{OH})_2$ [1] solution B $[\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}$ [1] solution C $\text{Cu}(\text{CH}_3\text{CO}_2)_2$ [1]	3															
3(d)(ii)	Na_2CO_3 or CO_3^{2-}	1															
3(d)(iii)	$\text{CuCO}_3 + 2\text{CH}_3\text{CO}_2\text{H} \rightarrow \text{Cu}(\text{CH}_3\text{CO}_2)_2 + \text{CO}_2 + \text{H}_2\text{O}$	1															
3(d)(iv)	any two for one mark - fizzing / bubbles / effervescence - solid disappears - green / blue solution (formed)	1															


PUBLISHED


Question	Answer	Marks
3(e)	sum of the charges of the (four) ligands equals the oxidation number / charge of Pt OR a calculation Pt +2, NH ₃ neutral / no charge, both Cl ⁻ 's -1 (so no overall charge)	1
3(f)(i)	 square planar and 180° [1]	2
3(f)(ii)	M1 this can bond / bind with DNA [1] M2 which prevents replication of the DNA / strand OR prevents cell division [1]	2
3(g)		1


Question	Answer	Marks
4(a)	M1 solubility decreases (down the Group) [1] M2 because lattice energy and hydration energy decreases OR lattice energy and hydration energy become less exothermic / more endothermic [1] M3 because hydration energy decreases to a greater extent (than does ΔH_{Latt}) [1]	3
4(b)(i)	$(K_w =) [\text{H}^+][\text{OH}^-]$	1


PUBLISHED

Question	Answer	Marks												
4(b)(ii)	[1] or each correct tick <table><tr><td>effect of increasing temperature</td><td>decreases</td><td>stay the same</td><td>increase</td></tr><tr><td>pH</td><td>✓</td><td></td><td></td></tr><tr><td>ratio of [H⁺]:[OH⁻]</td><td></td><td>✓</td><td></td></tr></table>	effect of increasing temperature	decreases	stay the same	increase	pH	✓			ratio of [H ⁺]:[OH ⁻]		✓		2
effect of increasing temperature	decreases	stay the same	increase											
pH	✓													
ratio of [H ⁺]:[OH ⁻]		✓												
4(c)	[H⁺] = 10^{-13.25} = 5.62 × 10⁻¹⁴ [1] [OH⁻] = K_w/[H⁺] = 1.0 × 10⁻¹⁴/5.62 × 10⁻¹⁴ [OH⁻] = 0.18 (0.178) (mol dm⁻³) [1] ecf correct answer scores [2]	2												
4(d)	HCO₃⁻ + H⁺ → H₂CO₃ OR HCO₃⁻ + H⁺ → CO₂ + H₂O [1] H₂CO₃ + OH⁻ → HCO₃⁻ + H₂O [1]	2												
4(e)(i)	CH₃COOH + H₂O ⇌ CH₃COO⁻ + H₃O⁺ [1] acid + base ⇌ base + acid [1]	2												
4(e)(ii)	M1 moles NaOH = 0.15 × 20/1000 = 0.0030 AND initial moles CH₃COOH = 0.25 × 30/1000 OR 0.0075 [1] M2 equilibrium moles CH₃COOH = 0.0045 AND equilibrium moles CH₃COONa = 0.0030 [1] M3 [CH₃COOH] = 0.0045/0.05 = 0.090 AND [CH₃COONa] = 0.003/0.05 = 0.060 [H⁺] = K_a × [CH₃COOH]/[CH₃COONa] = 2.625 × 10⁻⁵ [1] M4 pH = -log[H⁺] = 4.6 [1] correct answer scores [4]	4												
4(f)(i)	end point = 28 cm³	1												
4(f)(ii)	M1 reaction M bromothymol (blue) / bromocresol (green) AND reaction N bromothymol (blue) / thymolphthalein [1] M2 (both indicators have) a pH range / colour change within / in end-point / vertical region / sharp fall of the graph [1]	2												





Question	Answer	Marks												
5(a)(i)	[1] for each correct answer <table><tr><td></td><td>number of peaks</td></tr><tr><td>F</td><td>3</td></tr><tr><td>G</td><td>6</td></tr></table>		number of peaks	F	3	G	6	2						
	number of peaks													
F	3													
G	6													
5(a)(ii)	 one amide bond displayed in full [1] rest of the structure – one repeat unit only [1]	2												
5(b)	[1] for each correct tick <table><tr><td></td><td>σ-bonds only</td><td>π-bonds only</td><td>both σ- and π-bonds</td></tr><tr><td>bonds broken</td><td></td><td>✓</td><td></td></tr><tr><td>bonds formed</td><td>✓</td><td></td><td></td></tr></table>		σ -bonds only	π -bonds only	both σ - and π -bonds	bonds broken		✓		bonds formed	✓			2
	σ -bonds only	π -bonds only	both σ - and π -bonds											
bonds broken		✓												
bonds formed	✓													

Question	Answer	Marks
5(c)	 M1 length of chain with both monomers [1] M2 continuation bonds [1]	2
5(d)(i)	C-C bonds are non-polar / have no dipole so cannot be hydrolysed [1]	1
5(d)(ii)	M1 <u>Hydrolysis</u> using acid / base / alkali / enzymes [1] M2 action of UV light [1]	2


Question	Answer	Marks
6(a)	 M1 amide bond displayed [1] M2 rest of the structure [1]	2

Question	Answer	Marks
6(b)	 M1 valine on the cross [1] M2 Val-Lys and Lys on the right of the cross (in any order) [1] M3 relative order of Val-Lys and Lys (on the same side of the cross) [1] Explanation - Val does not move as it is a zwitterion / neutral (at pH6) OR Lys / Val-Lys move towards negative (pole) as they are positively charged - Lys moves the furthest as it has the lowest M_r (with the same positive charge) OR Val-Lys moves the least as it has the largest M_r (with the same positive charge) [1] × 2	5

Question	Answer	Marks
7(a)	M1 C-X / C-Cl / C-O bond is stronger (in chlorobenzene / phenol) [1] M2 p-orbital / lone pair on Cl / O(H) / X (in chlorobenzene / phenol) [1] M3 electrons of the (Cl / O / electronegative atom) AND overlap / delocalise with π-electron cloud / delocalise into ring [1]	3

Question	Answer	Marks
7(b)	 2-bromophenol  4-bromophenol structure and name correct [1]	2
7(c)(i)	step 1 conc. HNO_3 + H_2SO_4 (and temperare 50–55 °C) [1] step 2 Sn + HCl AND one of conc.HCl + heat [1] step 4 H_2O warm / heat [1]	3
7(c)(ii)	 OR 	1
7(c)(iii)	step 1 electrophilic substitution	1
7(c)(iv)	$\text{C}_6\text{H}_5\text{NO}_2$ + $6[\text{H}] \rightarrow \text{C}_6\text{H}_5\text{NH}_2$ + $2\text{H}_2\text{O}$	1

PUBLISHED

Question	Answer	Marks									
8(a)	 M1 continuous increase in S from 0–300 K (excluding m.p.) [1] M2 steep vertical increase in S ONLY at the m.p. AND continuous increase in S after m.p. [1]	2									
8(b)	[1] for each correct tick <table border="1"> <thead> <tr> <th></th><th>negative $\Delta S^\ominus$</th><th>positive $\Delta S^\ominus$</th></tr> </thead> <tbody> <tr> <td>solid dissolving in water</td><td></td><td>✓</td></tr> <tr> <td>water boiling to steam</td><td></td><td>✓</td></tr> </tbody> </table>		negative $\Delta S^\ominus$	positive $\Delta S^\ominus$	solid dissolving in water		✓	water boiling to steam		✓	1
	negative $\Delta S^\ominus$	positive $\Delta S^\ominus$									
solid dissolving in water		✓									
water boiling to steam		✓									
8(c)	$\Delta H^\ominus = (2 \times \text{C=O}) + (3 \times \text{H-H}) - (3 \times \text{C-H}) - (\text{C-O}) - (3 \times \text{O-H})$ $\Delta H^\ominus = (2 \times 805) + (3 \times 436) - (3 \times 410) - (1 \times 360) - (3 \times 460)$ [1] $\Delta H^\ominus = 1610 + 1308 - 1230 - 360 - 1380 = -52 \text{ (kJ mol}^{-1}\text{)}$ [1] ecf correct answer scores [2]	2									
8(d)(i)	$\Delta S^\ominus = 127 + 70 - (214 + 3 \times 131)$ [1] $= -410 \text{ (J K}^{-1} \text{ mol}^{-1}\text{)}$ [1] ecf correct answer scores [2]	2									
8(d)(ii)	$\Delta G^\ominus = \Delta H^\ominus - T\Delta S^\ominus$ [1] $\Delta G^\ominus = -131 - (298 \times -0.41) = -8.8(2) \text{ (kJ mol}^{-1}\text{)}$ [1] correct answer scores [2]	2									

PUBLISHED

Question	Answer	Marks
8(d)(iii)	(as temperature increases) feasibility decreases	1
8(e)(i)	$2\text{CH}_3\text{OH} + 3\text{O}_2 \rightleftharpoons 2\text{CO}_2 + 4\text{H}_2\text{O}$ OR $2\text{CH}_3\text{OH} + 3\text{O}_2 \rightleftharpoons 2\text{CO}_2 + 4\text{H}^+ + 4\text{OH}^-$	1
8(e)(ii)	$E^\ominus_{\text{cell}} = 1.23 - 0.02 = \mathbf{1.21\text{ V}}$	1