

MATHEMATICS

9709/31

Paper 3 Pure Mathematics 3 (P3)

October/November 2014

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
 Graph Paper
 List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 75.

Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

This document consists of **3** printed pages and **1** blank page.

- 1 Use logarithms to solve the equation $e^x = 3^{x-2}$, giving your answer correct to 3 decimal places. [3]

- 2 (i) Use the trapezium rule with 3 intervals to estimate the value of

$$\int_{\frac{1}{6}\pi}^{\frac{2}{3}\pi} \operatorname{cosec} x \, dx,$$

giving your answer correct to 2 decimal places. [3]

- (ii) Using a sketch of the graph of $y = \operatorname{cosec} x$, explain whether the trapezium rule gives an overestimate or an underestimate of the true value of the integral in part (i). [2]

- 3 The polynomial $ax^3 + bx^2 + x + 3$, where a and b are constants, is denoted by $p(x)$. It is given that $(3x + 1)$ is a factor of $p(x)$, and that when $p(x)$ is divided by $(x - 2)$ the remainder is 21. Find the values of a and b . [5]

- 4 The parametric equations of a curve are

$$x = \frac{1}{\cos^3 t}, \quad y = \tan^3 t,$$

where $0 \leq t < \frac{1}{2}\pi$.

- (i) Show that $\frac{dy}{dx} = \sin t$. [4]

- (ii) Hence show that the equation of the tangent to the curve at the point with parameter t is $y = x \sin t - \tan t$. [3]

- 5 **Throughout this question the use of a calculator is not permitted.**

The complex numbers w and z satisfy the relation

$$w = \frac{z + i}{iz + 2}.$$

- (i) Given that $z = 1 + i$, find w , giving your answer in the form $x + iy$, where x and y are real. [4]
- (ii) Given instead that $w = z$ and the real part of z is negative, find z , giving your answer in the form $x + iy$, where x and y are real. [4]

6 It is given that $\int_1^a \ln(2x) \, dx = 1$, where $a > 1$.

(i) Show that $a = \frac{1}{2} \exp\left(1 + \frac{\ln 2}{a}\right)$, where $\exp(x)$ denotes e^x . [6]

(ii) Use the iterative formula

$$a_{n+1} = \frac{1}{2} \exp\left(1 + \frac{\ln 2}{a_n}\right)$$

to determine the value of a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

7 In a certain country the government charges tax on each litre of petrol sold to motorists. The revenue per year is R million dollars when the rate of tax is x dollars per litre. The variation of R with x is modelled by the differential equation

$$\frac{dR}{dx} = R\left(\frac{1}{x} - 0.57\right),$$

where R and x are taken to be continuous variables. When $x = 0.5$, $R = 16.8$.

(i) Solve the differential equation and obtain an expression for R in terms of x . [6]

(ii) This model predicts that R cannot exceed a certain amount. Find this maximum value of R . [3]

8 (i) By first expanding $\sin(2\theta + \theta)$, show that

$$\sin 3\theta = 3 \sin \theta - 4 \sin^3 \theta. \quad [4]$$

(ii) Show that, after making the substitution $x = \frac{2 \sin \theta}{\sqrt{3}}$, the equation $x^3 - x + \frac{1}{6}\sqrt{3} = 0$ can be written in the form $\sin 3\theta = \frac{3}{4}$. [1]

(iii) Hence solve the equation

$$x^3 - x + \frac{1}{6}\sqrt{3} = 0,$$

giving your answers correct to 3 significant figures. [4]

9 Let $f(x) = \frac{x^2 - 8x + 9}{(1-x)(2-x)^2}$.

(i) Express $f(x)$ in partial fractions. [5]

(ii) Hence obtain the expansion of $f(x)$ in ascending powers of x , up to and including the term in x^2 . [5]

10 The line l has equation $\mathbf{r} = 4\mathbf{i} - 9\mathbf{j} + 9\mathbf{k} + \lambda(-2\mathbf{i} + \mathbf{j} - 2\mathbf{k})$. The point A has position vector $3\mathbf{i} + 8\mathbf{j} + 5\mathbf{k}$.

(i) Show that the length of the perpendicular from A to l is 15. [5]

(ii) The line l lies in the plane with equation $ax + by - 3z + 1 = 0$, where a and b are constants. Find the values of a and b . [5]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

MATHEMATICS

9709/32

Paper 3 Pure Mathematics 3 (P3)

October/November 2014

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 75.

Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

This document consists of **3** printed pages and **1** blank page.

- 1 Use logarithms to solve the equation $e^x = 3^{x-2}$, giving your answer correct to 3 decimal places. [3]

- 2 (i) Use the trapezium rule with 3 intervals to estimate the value of

$$\int_{\frac{1}{6}\pi}^{\frac{2}{3}\pi} \operatorname{cosec} x \, dx,$$

giving your answer correct to 2 decimal places. [3]

- (ii) Using a sketch of the graph of $y = \operatorname{cosec} x$, explain whether the trapezium rule gives an overestimate or an underestimate of the true value of the integral in part (i). [2]

- 3 The polynomial $ax^3 + bx^2 + x + 3$, where a and b are constants, is denoted by $p(x)$. It is given that $(3x + 1)$ is a factor of $p(x)$, and that when $p(x)$ is divided by $(x - 2)$ the remainder is 21. Find the values of a and b . [5]

- 4 The parametric equations of a curve are

$$x = \frac{1}{\cos^3 t}, \quad y = \tan^3 t,$$

where $0 \leq t < \frac{1}{2}\pi$.

- (i) Show that $\frac{dy}{dx} = \sin t$. [4]

- (ii) Hence show that the equation of the tangent to the curve at the point with parameter t is $y = x \sin t - \tan t$. [3]

- 5 **Throughout this question the use of a calculator is not permitted.**

The complex numbers w and z satisfy the relation

$$w = \frac{z + i}{iz + 2}.$$

- (i) Given that $z = 1 + i$, find w , giving your answer in the form $x + iy$, where x and y are real. [4]
- (ii) Given instead that $w = z$ and the real part of z is negative, find z , giving your answer in the form $x + iy$, where x and y are real. [4]

6 It is given that $\int_1^a \ln(2x) \, dx = 1$, where $a > 1$.

(i) Show that $a = \frac{1}{2} \exp\left(1 + \frac{\ln 2}{a}\right)$, where $\exp(x)$ denotes e^x . [6]

(ii) Use the iterative formula

$$a_{n+1} = \frac{1}{2} \exp\left(1 + \frac{\ln 2}{a_n}\right)$$

to determine the value of a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

7 In a certain country the government charges tax on each litre of petrol sold to motorists. The revenue per year is R million dollars when the rate of tax is x dollars per litre. The variation of R with x is modelled by the differential equation

$$\frac{dR}{dx} = R\left(\frac{1}{x} - 0.57\right),$$

where R and x are taken to be continuous variables. When $x = 0.5$, $R = 16.8$.

(i) Solve the differential equation and obtain an expression for R in terms of x . [6]

(ii) This model predicts that R cannot exceed a certain amount. Find this maximum value of R . [3]

8 (i) By first expanding $\sin(2\theta + \theta)$, show that

$$\sin 3\theta = 3 \sin \theta - 4 \sin^3 \theta. \quad [4]$$

(ii) Show that, after making the substitution $x = \frac{2 \sin \theta}{\sqrt{3}}$, the equation $x^3 - x + \frac{1}{6}\sqrt{3} = 0$ can be written in the form $\sin 3\theta = \frac{3}{4}$. [1]

(iii) Hence solve the equation

$$x^3 - x + \frac{1}{6}\sqrt{3} = 0,$$

giving your answers correct to 3 significant figures. [4]

9 Let $f(x) = \frac{x^2 - 8x + 9}{(1-x)(2-x)^2}$.

(i) Express $f(x)$ in partial fractions. [5]

(ii) Hence obtain the expansion of $f(x)$ in ascending powers of x , up to and including the term in x^2 . [5]

10 The line l has equation $\mathbf{r} = 4\mathbf{i} - 9\mathbf{j} + 9\mathbf{k} + \lambda(-2\mathbf{i} + \mathbf{j} - 2\mathbf{k})$. The point A has position vector $3\mathbf{i} + 8\mathbf{j} + 5\mathbf{k}$.

(i) Show that the length of the perpendicular from A to l is 15. [5]

(ii) The line l lies in the plane with equation $ax + by - 3z + 1 = 0$, where a and b are constants. Find the values of a and b . [5]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

MATHEMATICS

9709/33

Paper 3 Pure Mathematics 3 (P3)

October/November 2014

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
 Graph Paper
 List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 75.

Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

This document consists of **3** printed pages and **1** blank page.

- 1 Solve the inequality $|3x - 1| < |2x + 5|$. [4]

- 2 A curve is defined for $0 < \theta < \frac{1}{2}\pi$ by the parametric equations

$$x = \tan \theta, \quad y = 2 \cos^2 \theta \sin \theta.$$

Show that $\frac{dy}{dx} = 6 \cos^5 \theta - 4 \cos^3 \theta$. [5]

- 3 The polynomial $4x^3 + ax^2 + bx - 2$, where a and b are constants, is denoted by $p(x)$. It is given that $(x + 1)$ and $(x + 2)$ are factors of $p(x)$.

(i) Find the values of a and b . [4]

(ii) When a and b have these values, find the remainder when $p(x)$ is divided by $(x^2 + 1)$. [3]

- 4 (i) Show that $\cos(\theta - 60^\circ) + \cos(\theta + 60^\circ) \equiv \cos \theta$. [3]

(ii) Given that $\frac{\cos(2x - 60^\circ) + \cos(2x + 60^\circ)}{\cos(x - 60^\circ) + \cos(x + 60^\circ)} = 3$, find the exact value of $\cos x$. [4]

- 5 The complex numbers w and z are defined by $w = 5 + 3i$ and $z = 4 + i$.

(i) Express $\frac{iw}{z}$ in the form $x + iy$, showing all your working and giving the exact values of x and y . [3]

(ii) Find wz and hence, by considering arguments, show that

$$\tan^{-1}\left(\frac{3}{5}\right) + \tan^{-1}\left(\frac{1}{4}\right) = \frac{1}{4}\pi. \quad [4]$$

- 6 It is given that $I = \int_0^{0.3} (1 + 3x^2)^{-2} dx$.

(i) Use the trapezium rule with 3 intervals to find an approximation to I , giving the answer correct to 3 decimal places. [3]

(ii) For small values of x , $(1 + 3x^2)^{-2} \approx 1 + ax^2 + bx^4$. Find the values of the constants a and b .

Hence, by evaluating $\int_0^{0.3} (1 + ax^2 + bx^4) dx$, find a second approximation to I , giving the answer correct to 3 decimal places. [5]

- 7 The equations of two straight lines are

$$\mathbf{r} = \mathbf{i} + 4\mathbf{j} - 2\mathbf{k} + \lambda(\mathbf{i} + 3\mathbf{k}) \quad \text{and} \quad \mathbf{r} = a\mathbf{i} + 2\mathbf{j} - 2\mathbf{k} + \mu(\mathbf{i} + 2\mathbf{j} + 3a\mathbf{k}),$$

where a is a constant.

- (i) Show that the lines intersect for all values of a . [4]
- (ii) Given that the point of intersection is at a distance of 9 units from the origin, find the possible values of a . [4]

- 8 The variables x and y are related by the differential equation

$$\frac{dy}{dx} = \frac{1}{5}xy^{\frac{1}{2}} \sin\left(\frac{1}{3}x\right).$$

- (i) Find the general solution, giving y in terms of x . [6]
- (ii) Given that $y = 100$ when $x = 0$, find the value of y when $x = 25$. [3]

- 9 (i) Sketch the curve $y = \ln(x + 1)$ and hence, by sketching a second curve, show that the equation

$$x^3 + \ln(x + 1) = 40$$

has exactly one real root. State the equation of the second curve. [3]

- (ii) Verify by calculation that the root lies between 3 and 4. [2]

- (iii) Use the iterative formula

$$x_{n+1} = \sqrt[3]{(40 - \ln(x_n + 1))},$$

with a suitable starting value, to find the root correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

- (iv) Deduce the root of the equation

$$(e^y - 1)^3 + y = 40,$$

giving the answer correct to 2 decimal places. [2]

- 10 By first using the substitution $u = e^x$, show that

$$\int_0^{\ln 4} \frac{e^{2x}}{e^{2x} + 3e^x + 2} dx = \ln\left(\frac{8}{5}\right). \quad [10]$$

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.