



Cambridge International A Level

MATHEMATICS

9709/61

Paper 6 Probability & Statistics 2

October/November 2022

MARK SCHEME

Maximum Mark: 50

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2022 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

This document consists of **11** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

PUBLISHED

Mathematics Specific Marking Principles	
1	Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
2	Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
3	Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
4	Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
5	Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread.
6	Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

PUBLISHED**Mark Scheme Notes**

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B** Mark for a correct result or statement independent of method marks.
- DM or DB** When a part of a question has two or more ‘method’ steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- FT** Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.
- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
 - For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
 - The total number of marks available for each question is shown at the bottom of the Marks column.
 - Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
 - Square brackets [] around text or numbers show extra information not needed for the mark to be awarded.

Abbreviations

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no ‘follow through’ from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

PUBLISHED

Question	Answer	Marks	Guidance
1	$\frac{62.1}{10} = 6.21$	B1	OE
	$[\Sigma x^2 = 387.05]$ $\frac{10}{9} \left(\frac{\text{their '387.05'}}{10} - (\text{their '6.21'})^2 \right)$ or $\frac{1}{9} \left(\frac{\text{their '387.05'}}{10} - \frac{(\text{their '6.21'})^2}{10} \right)$	M1	Can be implied. Accept alternative methods (e.g. working mean of 6). Biased 0.1409 M0.
	$= 0.157 \text{ (3 sf) or } \frac{1409}{9000}$	A1	
		3	

Question	Answer	Marks	Guidance
2(a)	$H_0: P(\text{red}) = 0.2 \quad H_1: P(\text{red}) < 0.2$	B1	Allow $H_0: p = 0.2 \quad H_1: p < 0.2$.
	$P(X \leq 4) = 0.8^{40} + 40 \times 0.8^{39} \times 0.2 + {}^{40}C_2 \times 0.8^{38} \times 0.2^2$ $+ {}^{40}C_3 \times 0.8^{37} \times 0.2^3 + {}^{40}C_4 \times 0.8^{36} \times 0.2^4$	M1	For full expression seen. Allow one term omitted, incorrect or extra.
	0.0759	A1	SC 0.0759 without working B1.
	<i>their</i> '0.0759' > 0.05	M1	Valid comparison (from binomial probs) of their $P(X \leq 4)$ with 0.05.
	[Do not reject H_0]. Not enough evidence that it lands on red fewer times than if it were fair or not enough evidence to suggest that the spinner is biased	A1 FT	FT their 0.0759. In context, not definite, no contradictions.
		5	

PUBLISHED

Question	Answer	Marks	Guidance
4(a)	Mean = $[3(2500 + 3700)] = 18600$ (kg)	B1	
	Var(Total profit) = $3(120^2 + 130^2)$ or 93900	M1	or $\sqrt{\text{ of this stated.}}$
	sd = 306 (kg) (3 sf)	A1	
		3	
4(b)	$E(1.5X - 0.2Y) = 1.5 \times 2500 - 0.20 \times 3700 = [3010]$	B1	Give at early stage.
	$\text{Var}(1.5X - 0.2Y) = 1.5^2 \times 120^2 + 0.2^2 \times 130^2 \quad [= 33076]$	B1	Correct expression or result or sd = 182 (3 sf) seen.
	$\frac{3000 - 3010}{\sqrt{\text{their '33076'}}} \quad [= -0.055]$	M1	Ignore continuity correction attempts. E(X) and Var must come from a combination attempt. Can be implied.
	$\Phi(\text{their '-0.055'}) = 1 - \Phi(\text{their '0.055'})$	M1	For area consistent with their values. Can be implied.
	= 0.478 (3 sf)	A1	
		5	

Question	Answer	Marks	Guidance
5(a)	$45 \pm z \times \frac{6}{\sqrt{200}}$	M1	For expression of correct form, any z. Accept one side of interval for M1.
	$z = 1.96$	B1	Must be seen.
	44.2 to 45.8 (3 sf)	A1	Must be an interval.
		3	

PUBLISHED

Question	Answer	Marks	Guidance
5(b)	$z \times \frac{11}{\sqrt{200}} = 2$	M1	Or ... = 4 for M1
	$z = 2.571$	A1	Accept 3sf if nothing better seen.
	$\Phi(\text{their '2.571'}) = 0.9949$ and $\text{their '0.9949'} - (1 - \text{their '0.9949'}) [= 0.9898]$	M1	OE For area consistent with their values. Must be seen.
	$\alpha = 99.0$ (3 sf)	A1	Allow 99. cwo Final answer of 0.99 scores A0.
		4	

Question	Answer	Marks	Guidance
6(a)	Curve of similar shape, $x = 0$ to $x = 4$, with highest point (2, 0.375)	B1	Not straight lines, not bell shaped. Must be correct at $x = 0$ and $x = 4$, highest point must be at $x = 2$, y value $\pm \frac{1}{4}$ square. Must not go below the x-axis.
		1	
6(b)	Curve of similar shape, from $x = 0$ to $x = 2$, highest point at $x = 1$	B1	Not straight lines, not bell shaped. Must be correct at $x = 0$ and $x = 2$. Highest point must be at $x = 1$.
	Highest point (1, 0.75)	B1	
		2	

PUBLISHED

Question	Answer	Marks	Guidance
6(c)	$\frac{3}{32} \int_{1+a}^3 (3+2x-x^2)dx = \frac{1}{4}$ or $\frac{3}{32} \int_{1-a}^{1+a} (3+2x-x^2)dx = \frac{1}{2}$	M1	OE Attempt to integrate f(x) and correct limits with correct RHS.
	$\frac{3}{32} \left[3x + x^2 - \frac{x^3}{3} \right]_{1+a}^3 = \frac{1}{4}$ or $\frac{3}{32} \left[3x + x^2 - \frac{x^3}{3} \right]_{1-a}^{1+a} = \frac{1}{2}$	A1	Correct integration.
	$a^3 - 12a + 8 = 0$	A1	AG Substitute limits and correctly obtain equation. May see $3/32(6a+4a-6a/3-2a^3/3) = 0.5$ No errors seen..
		3	
6(d)	$0.69^3 - 12 \times 0.69 + 8 = 0.049$ (2 sf) > 0 $0.70^3 - 12 \times 0.70 + 8 = -0.057$ (2 sf) < 0 Hence $0.69 < a < 0.70$	B1	AG Must state either the correct expression and > 0 and < 0 or both answers to 2 sf. Both answers correct and conclusion. Accept equivalent expressions. OR: $a = 0.695$ (3 sf) which is between 0.69 & 0.70.
		1	

PUBLISHED

Question	Answer	Marks	Guidance
7(a)	H_0 : Population mean time (or μ) = 32.5 H_1 : Population mean time (or μ) < 32.5	B1	Not just “mean”.
	$\pm \frac{31.8 - 32.5}{3.1 \div \sqrt{50}}$	M1	Must have $\sqrt{50}$. Could be implied.
	= ± -1.597	A1	
	‘-1.597’ < -1.406 [or ‘1.597’ > 1.406]	M1	Valid comparison of their z_{calc} with ± 1.406 . or $0.0551 < 0.08$ (or $0.0552 < 0.08$).
	[reject H_0] There is evidence that [population] [mean] time has decreased	A1 FT	In context, not definite, no contradictions. Note: Accept critical value method 31.88 (31.9) M1 A1 and $31.8 < 31.88$ M1 conclusion A1 .
		5	
7(b)	$\frac{a - 32.5}{3.1 \div \sqrt{50}} = -1.406$	M1	Standardise with 32.5 and $\sqrt{50}$ and z value on RHS.
	$a = 31.88$ or 31.9	A1	May be seen in part (a). Can score M1A1 here as well using a similar approach to (a).
	$\frac{\text{their '31.88 - 31.5}}{3.1 \div \sqrt{50}} \quad [= 0.8668 \text{ to } 0.8760]$	M1	Standardise with <i>their</i> cv and mean = 31.5. Must have $\sqrt{50}$.
	$1 - \Phi(0.8668)$	M1	For area consistent with their working.
	= 0.190 to 0.193 (3 sf)	A1	
		5	



Cambridge International A Level

MATHEMATICS

9709/62

Paper 6 Probability & Statistics 2

October/November 2022

MARK SCHEME

Maximum Mark: 50

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2022 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

This document consists of **12** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

PUBLISHED

Mathematics Specific Marking Principles	
1	Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
2	Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
3	Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
4	Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
5	Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread.
6	Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

PUBLISHED**Mark Scheme Notes**

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B** Mark for a correct result or statement independent of method marks.
- DM or DB** When a part of a question has two or more ‘method’ steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- FT** Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.
- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
 - For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
 - The total number of marks available for each question is shown at the bottom of the Marks column.
 - Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
 - Square brackets [] around text or numbers show extra information not needed for the mark to be awarded.

Abbreviations

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no ‘follow through’ from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

PUBLISHED

Question	Answer	Marks	Guidance
1(a)	Est $\mu = 25.6$ or $\frac{2048}{80}$ or $\frac{128}{5}$	B1	
	Est $\sigma^2 = \frac{80}{79} \left(\frac{52760}{80} - \left(\frac{2048}{80} \right)^2 \right)$ or $\frac{1}{79} \left(52760 - \frac{2048^2}{80} \right)$	M1	Substitution into a correct formula. Biased 4.14 scores M0.
	= 4.19 (3 sf) or $\frac{1656}{395}$	A1	
		3	
1(b)	'25.6' + $z \sqrt{\frac{4.19}{80}}$ = 26.0	M1	Use of correct equation with their values.
	$z = 1.748$ or 1.747	A1	Accept 3sf. FT Biased $z = 1.758$.
	$(\Phi(1.748) = 0.960)$ '0.960' – (1 – '0.960')	M1	Correct area using their values.
	$\alpha = 92.0$ or 91.9	A1	Allow 92 . FT Biased 92.1. A final answer of 0.92 or 0.919 scores A0.
		4	

PUBLISHED

Question	Answer	Marks	Guidance
2	H ₀ : Population mean length = 10.3 cm H ₁ : Population mean length < 10.3 cm	B1	or $\mu = 10.3$ (not just ‘mean’). $\mu < 10.3$
	$\pm \frac{9.8 - 10.3}{2.6 / \sqrt{100}}$	M1	If ± 1.923 (or 0.0272) seen allow M1 implied.
	= -1.923	A1	Accept $\pm$. Accept 3sf.
	-1.923 > -2.054 or -2.055	M1	OE For a valid comparison. Or compare $1 - \phi(1.923)$ with 0.02 e.g. $0.0272 > 0.02$ Use of CV $9.8 > 9.766$ scores M1 A1 for 9.766 and M1 for comparison.
	[Not reject H ₀] No evidence that [mean] length has decreased	A1 FT	FT <i>their</i> z. No contradictions, not definite, in context.
		5	

Question	Answer	Marks	Guidance
3(a)	Use of Poisson. mean = 3.2	B1 B1	
	$1 - e^{-3.2} \left(1 + 3.2 + \frac{3.2^2}{2} + \frac{3.2^3}{3!} \right)$ or $1 - e^{-3.2} (1 + 3.2 + 5.12 + 5.46133)$ or $1 - (0.04076 + 0.1304 + 0.2087 + 0.2226)$	M1	Allow any λ . Allow one end error.
	= 0.397 or 0.398	A1	SC Use of binomial: B1 for answer 0.398 (3 sf). 0.397 or 0.398 with no working scores SC B1 .
		4	

PUBLISHED

Question	Answer	Marks	Guidance
3(b)	[Binomial with] [n =] 200 > 50	B1	
	[np =][200 × 0.016 =] 3.2 < 5 or [p =]0.016 < 0.1	B1	If B0 B0 SC n large (or $n > 50$), and p small or $p < 0.1$ or $np < 5$: B1.
		2	

Question	Answer	Marks	Guidance
4(a)	H_0 : Population mean = 7.2 or 2.4 H_1 : Population mean < 7.2 or 2.4	B1	or λ or μ = 7.2 or 2.4 (Not just ‘mean’). or λ or μ < 7.2 or 2.4
		1	
4(b)	$\lambda = 7.2$	B1	SOI
	$[P(X \leq 2)] = e^{-7.2} \left(1 + 7.2 + \frac{7.2^2}{2} \right)$ or $e^{-7.2}(1 + 7.2 + 25.92)$ or $0.0007465 + 0.0053754 + 0.01935 [= 0.0255]$ $[P(X \leq 3)] = '0.0255' + e^{-7.2} \times \frac{7.2^3}{3!}$ or $'0.0255' + e^{-7.2} (62.21)$ or $'0.0255' + 0.04644 [= 0.0719]$	M1	Both expressions needed, allow any λ If $\lambda \neq 7.2$ allow $P(X \leq n)$ for 2 consecutive values of n with $P(X \leq n) < 0.05$ and $P(X \leq n + 1) > 0.05$.
	$P(\text{Type I}) = 0.02547$ or 0.0255 (3 sf)	B1	
		3	

PUBLISHED

Question	Answer	Marks	Guidance
4(c)	$3 > 2$ or $P(X \leq 3) > 0.05$ or '0.0719' > 0.05	M1	For a valid comparison or 3 outside critical region. FT <i>their</i> CR in (b) .
	[Not reject H_0] No evidence that [mean] number of faults has decreased	A1 FT	No contradictions. In context, not definite.
		2	
4(d)	$1 - e^{-1.5}(1 + 1.5 + 1.5^2 / 2)$ or $1 - e^{-1.5}(1 + 1.5 + 1.125)$ or $1 - (0.2231 + 0.3347 + 0.2510)$	M1	Must see expression. FT <i>their</i> CR in (b) .
	= 0.191 (3 sf)	A1	
		2	

Question	Answer	Marks	Guidance
5(a)	For $X, \mu = 2 \sigma^2 = 1.6$		
	Mean = 2	B1	
	Variance = $\frac{1.6}{160}$ or $\frac{1}{100}$ or 0.01	B1	Accept Var = 0.1^2 (accept sd=0.1 if clearly identified).
	Normal	B1	
		3	

PUBLISHED

Question	Answer	Marks	Guidance
5(b)	$\pm \frac{1.8 - \frac{1}{320} - '2'}{\sqrt{0.01}}$ or $\pm \frac{1.8 - '2'}{\sqrt{0.01}}$ [= -2.03 or -2] or $\pm (287.5 - '320') / \sqrt{256}$ or $\pm (288 - '320') / \sqrt{256}$ [= -2.03 or -2]	M1	Allow with wrong continuity correction. M1 can be implied by correct final answer or for -2.03 / -2.0 or 0.9788 / 0.9772 seen.
	$\Phi(' -2.03') = 1 - \Phi('2.03')$	M1	Correct area consistent with their values. M1 can be implied by correct final answer.
	= 0.0212 or 0.0228 (3 sf)	A1	
		3	

Question	Answer	Marks	Guidance
6(a)	$E(T) = 4 \times 510 + 2 \times 1015$ [= 4070]	B1	
	$\text{Var}(T) = 4 \times 100 + 2 \times 324$ [= 1048]	B1	or $\sqrt{(4 \times 100 + 2 \times 324)} [= 32.4(3\text{sf})]$.
	$\frac{4130 - 4070}{\sqrt{1048}}$ [= 1.853]	M1	Standardising with their values. Variance must be from a combination attempt. M1 can be implied by correct final answer.
	$\Phi('1.853')$	M1	For area consistent with their values. M1 can be implied by correct final answer.
	= 0.968 (3 sf)	A1	As final answer.
		5	

PUBLISHED

Question	Answer	Marks	Guidance
6(b)	$E(D) = 4 \times 510 - 2 \times 1015 [= 10]$	B1	
	$\text{Var}(D) = 4 \times 100 + 2 \times 324 [= 1048]$	B1	Or $\sqrt{(4 \times 100 + 2 \times 324)} [= 32.4(3\text{sf})]$.
	$\frac{0 - '10'}{\sqrt{'1048'}} [= -0.309]$	M1	Standardising with their values. Variance must be from a combination attempt. M1 can be implied by correct final answer.
	$1 - \Phi(' -0.309') = \Phi('0.309')$	M1	For area consistent with their values. M1 can be implied by correct final answer.
	$= 0.621$	A1	As final answer.
		5	

Question	Answer	Marks	Guidance
7(a)	$1 - p$ or $p - 0.5$	M1	SOI, e.g. on diagram.
	$[P(-1 < X < 0) =] 2p - 1$	A1	Clearly as final answer.
		2	
7(b)(i)	$\int_{-3}^2 (a - b(x^2 + x))dx = 1$ or $\int_{-3}^2 (ax - b(x^3 + x^2))dx = -0.5$	M1	OE Attempt integral, with correct limits and RHS.
	$\left[ax - b\left(\frac{x^3}{3} + \frac{x^2}{2}\right) \right]_{-3}^2 (= 1)$ or $\left[a\frac{x^2}{2} - b\left(\frac{x^4}{4} + \frac{x^3}{3}\right) \right]_{-3}^2 (= -0.5)$	A1	OE Correct integration.
	$2a - 8b/3 - 2b + 3a - 9b + 9b/2 = 1$ or $2a - 4b - 8b/3 - 9a/2 + 81b/4 - 9b = -0.5$ leading to $30a - 55b = 6$ AG	A1	Correctly obtained. No errors seen.
		3	

PUBLISHED

Question	Answer	Marks	Guidance
7(b)(ii)	$a - b(9 - 3) = 0$ or $a - b(4 + 2) = 0$ [hence $a - 6b = 0$]	*M1	Use $f(-3) = 0$ or $f(2) = 0$. Further attempts at integration M0.
	Attempt to solve $30a - 55b = 6$ and their $a - 6b = 0$	DM1	
	$a = \frac{36}{125}$ or 0.288 $b = \frac{6}{125}$ or 0.048	A1	
		3	



Cambridge International A Level

MATHEMATICS

9709/63

Paper 6 Probability & Statistics 2

October/November 2022

MARK SCHEME

Maximum Mark: 50

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2022 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

This document consists of **11** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

PUBLISHED

Mathematics Specific Marking Principles	
1	Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
2	Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
3	Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
4	Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
5	Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread.
6	Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

PUBLISHED**Mark Scheme Notes**

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B** Mark for a correct result or statement independent of method marks.
- DM or DB** When a part of a question has two or more ‘method’ steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- FT** Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.
- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
 - For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
 - The total number of marks available for each question is shown at the bottom of the Marks column.
 - Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
 - Square brackets [] around text or numbers show extra information not needed for the mark to be awarded.

Abbreviations

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no ‘follow through’ from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

PUBLISHED

Question	Answer	Marks	Guidance
1	$\frac{62.1}{10} = 6.21$	B1	OE
	$[\Sigma x^2 = 387.05]$ $\frac{10}{9} \left(\frac{\text{their '387.05'}}{10} - (\text{their '6.21'})^2 \right)$ or $\frac{1}{9} \left(\frac{\text{their '387.05'}}{10} - \frac{(\text{their '6.21'})^2}{10} \right)$	M1	Can be implied. Accept alternative methods (e.g. working mean of 6). Biased 0.1409 M0.
	$= 0.157 \text{ (3 sf) or } \frac{1409}{9000}$	A1	
		3	

Question	Answer	Marks	Guidance
2(a)	$H_0: P(\text{red}) = 0.2 \quad H_1: P(\text{red}) < 0.2$	B1	Allow $H_0: p = 0.2 \quad H_1: p < 0.2$.
	$P(X \leq 4) = 0.8^{40} + 40 \times 0.8^{39} \times 0.2 + {}^{40}C_2 \times 0.8^{38} \times 0.2^2$ $+ {}^{40}C_3 \times 0.8^{37} \times 0.2^3 + {}^{40}C_4 \times 0.8^{36} \times 0.2^4$	M1	For full expression seen. Allow one term omitted, incorrect or extra.
	0.0759	A1	SC 0.0759 without working B1.
	<i>their</i> '0.0759' > 0.05	M1	Valid comparison (from binomial probs) of their $P(X \leq 4)$ with 0.05.
	[Do not reject H_0]. Not enough evidence that it lands on red fewer times than if it were fair or not enough evidence to suggest that the spinner is biased	A1 FT	FT their 0.0759. In context, not definite, no contradictions.
		5	

PUBLISHED

Question	Answer	Marks	Guidance
4(a)	Mean = $[3(2500 + 3700)] = 18600$ (kg)	B1	
	Var(Total profit) = $3(120^2 + 130^2)$ or 93900	M1	or $\sqrt{\text{ of this stated.}}$
	sd = 306 (kg) (3 sf)	A1	
		3	
4(b)	$E(1.5X - 0.2Y) = 1.5 \times 2500 - 0.20 \times 3700 = [3010]$	B1	Give at early stage.
	$\text{Var}(1.5X - 0.2Y) = 1.5^2 \times 120^2 + 0.2^2 \times 130^2$ [= 33076]	B1	Correct expression or result or sd = 182 (3 sf) seen.
	$\frac{3000 - 3010}{\sqrt{\text{their '33076'}}} \quad [= -0.055]$	M1	Ignore continuity correction attempts. E(X) and Var must come from a combination attempt. Can be implied.
	$\Phi(\text{their '-0.055'}) = 1 - \Phi(\text{their '0.055'})$	M1	For area consistent with their values. Can be implied.
	= 0.478 (3 sf)	A1	
		5	

Question	Answer	Marks	Guidance
5(a)	$45 \pm z \times \frac{6}{\sqrt{200}}$	M1	For expression of correct form, any z. Accept one side of interval for M1.
	$z = 1.96$	B1	Must be seen.
	44.2 to 45.8 (3 sf)	A1	Must be an interval.
		3	

PUBLISHED

Question	Answer	Marks	Guidance
5(b)	$z \times \frac{11}{\sqrt{200}} = 2$	M1	Or ... = 4 for M1
	$z = 2.571$	A1	Accept 3sf if nothing better seen.
	$\Phi(\text{their '2.571'}) = 0.9949$ and $\text{their '0.9949'} - (1 - \text{their '0.9949'}) [= 0.9898]$	M1	OE For area consistent with their values. Must be seen.
	$\alpha = 99.0$ (3 sf)	A1	Allow 99. cwo Final answer of 0.99 scores A0.
		4	

Question	Answer	Marks	Guidance
6(a)	Curve of similar shape, $x = 0$ to $x = 4$, with highest point (2, 0.375)	B1	Not straight lines, not bell shaped. Must be correct at $x = 0$ and $x = 4$, highest point must be at $x = 2$, y value $\pm \frac{1}{4}$ square. Must not go below the x-axis.
		1	
6(b)	Curve of similar shape, from $x = 0$ to $x = 2$, highest point at $x = 1$	B1	Not straight lines, not bell shaped. Must be correct at $x = 0$ and $x = 2$. Highest point must be at $x = 1$.
	Highest point (1, 0.75)	B1	
		2	

PUBLISHED

Question	Answer	Marks	Guidance
6(c)	$\frac{3}{32} \int_{1+a}^3 (3+2x-x^2)dx = \frac{1}{4}$ or $\frac{3}{32} \int_{1-a}^{1+a} (3+2x-x^2)dx = \frac{1}{2}$	M1	OE Attempt to integrate f(x) and correct limits with correct RHS.
	$\frac{3}{32} \left[3x + x^2 - \frac{x^3}{3} \right]_{1+a}^3 = \frac{1}{4}$ or $\frac{3}{32} \left[3x + x^2 - \frac{x^3}{3} \right]_{1-a}^{1+a} = \frac{1}{2}$	A1	Correct integration.
	$a^3 - 12a + 8 = 0$	A1	AG Substitute limits and correctly obtain equation. May see $3/32(6a+4a-6a/3-2a^3/3) = 0.5$ No errors seen..
		3	
6(d)	$0.69^3 - 12 \times 0.69 + 8 = 0.049$ (2 sf) > 0 $0.70^3 - 12 \times 0.70 + 8 = -0.057$ (2 sf) < 0 Hence $0.69 < a < 0.70$	B1	AG Must state either the correct expression and > 0 and < 0 or both answers to 2 sf. Both answers correct and conclusion. Accept equivalent expressions. OR: $a = 0.695$ (3 sf) which is between 0.69 & 0.70.
		1	

PUBLISHED

Question	Answer	Marks	Guidance
7(a)	H_0 : Population mean time (or μ) = 32.5 H_1 : Population mean time (or μ) < 32.5	B1	Not just “mean”.
	$\pm \frac{31.8 - 32.5}{3.1 \div \sqrt{50}}$	M1	Must have $\sqrt{50}$. Could be implied.
	= ± -1.597	A1	
	‘-1.597’ < -1.406 [or ‘1.597’ > 1.406]	M1	Valid comparison of their z_{calc} with ± 1.406 . or $0.0551 < 0.08$ (or $0.0552 < 0.08$).
	[reject H_0] There is evidence that [population] [mean] time has decreased	A1 FT	In context, not definite, no contradictions. Note: Accept critical value method 31.88 (31.9) M1 A1 and $31.8 < 31.88$ M1 conclusion A1 .
		5	
7(b)	$\frac{a - 32.5}{3.1 \div \sqrt{50}} = -1.406$	M1	Standardise with 32.5 and $\sqrt{50}$ and z value on RHS.
	$a = 31.88$ or 31.9	A1	May be seen in part (a). Can score M1A1 here as well using a similar approach to (a).
	$\frac{\text{their '31.88 - 31.5}}{3.1 \div \sqrt{50}}$ [= 0.8668 to 0.8760]	M1	Standardise with <i>their</i> cv and mean = 31.5. Must have $\sqrt{50}$.
	$1 - \Phi(0.8668)$	M1	For area consistent with their working.
	= 0.190 to 0.193 (3 sf)	A1	
		5	