CAMBRIDGE INTERNATIONAL AS & A LEVEL PHYSICS · 9702

PHYSICS



EXAM QUESTION PACK

12 MOTION IN A CIRCLE

SCORE	PERCENTAGE
/	%

STEMUP

These questions belong to CIE, not to us.

The examination questions, mark schemes and examiner comments in this pack were written and published by Cambridge International (CIE). They are past assessment materials, reproduced here so that students can revise from them.

What we do not own

STEMUP is not affiliated with CIE and does not own, and makes no claim to own, any of the question, mark scheme or examiner report material in this pack. CIE has not endorsed, approved or been involved in the production of it. CIE and the qualification titles used here belong to their respective owners and appear for identification only.

Where to find the originals

CIE publishes its past papers, mark schemes and examiner reports. Always check the current syllabus at cambridgeinternational.org before you revise.

If you hold the rights to any material in this pack and would like it amended or removed, contact admin@stemup.co.uk and we will act on it.

Chapter 1

Motion in a circle

STEMUP.CO.UK

1. 9702_w21_qp_41 Q: 1

(a) With reference to velocity and acceleration, describe uniform circular motion.

.....

.....

..... [2]

(b) Two cars are moving around a horizontal circular track. One car follows path X and the other follows path Y, as shown in Fig. 1.1.




Fig. 1.1 (not to scale)

The radius of path X is 318 m. Path Y is parallel to, and 27 m outside, path X. Both cars have mass 790 kg. The maximum lateral (sideways) friction force F that the cars can experience without sliding is the same for both cars.

(i) The maximum speed at which the car on path X can move around the track without sliding is 94 m s^{-1}.

Calculate F.

F = ... N [2]

(ii) Both cars move around the track. Each car has the maximum speed at which it can move without sliding.

Complete Table 1.1, by placing one tick in each row, to indicate how the quantities indicated for the car on path Y compare with the car on path X.

Table 1.1

	Y less than X	Y same as X	Y greater than X
centripetal acceleration			
maximum speed			
time taken for one lap of the track			

[3]

[Total: 7]

2. 9702_w21_qp_42 Q: 1

(a) State what is meant by *centripetal* acceleration.

.....

.....

..... [1]

(b) An unpowered toy car moves freely along a smooth track that is initially horizontal. The track contains a vertical circular loop around which the car travels, as shown in Fig. 1.1.




Fig. 1.1

The mass of the car is 230 g and the diameter of the loop is 62 cm. Assume that the resistive forces acting on the car are negligible.

(i) State what happens to the magnitude of the centripetal acceleration of the car as it moves around the loop from X to Y.

..... [1]

(ii) Explain, if the car remains in contact with the track, why the centripetal acceleration of the car at point Y must be greater than 9.8 m s^{-2}.

.....

.....

..... [2]

(c) The initial speed at which the car in **(b)** moves along the track is $3.8\,m\,s^{-1}$.

Determine whether the car is in contact with the track at point Y. Show your working.

[3]

(d) Suggest, with a reason but without calculation, whether your conclusion in **(c)** would be different for a car of mass 460 g moving with the same initial speed.

.....

.....

..... [1]

[Total: 8]

3. 9702_w21_qp_43 Q: 1

(a) With reference to velocity and acceleration, describe uniform circular motion.

.....

.....

..... [2]

(b) Two cars are moving around a horizontal circular track. One car follows path X and the other follows path Y, as shown in Fig. 1.1.




Fig. 1.1 (not to scale)

The radius of path X is 318 m. Path Y is parallel to, and 27 m outside, path X. Both cars have mass 790 kg. The maximum lateral (sideways) friction force F that the cars can experience without sliding is the same for both cars.

(i) The maximum speed at which the car on path X can move around the track without sliding is 94 m s^{-1}.

Calculate F.

F = ... N [2]

(ii) Both cars move around the track. Each car has the maximum speed at which it can move without sliding.

Complete Table 1.1, by placing one tick in each row, to indicate how the quantities indicated for the car on path Y compare with the car on path X.

Table 1.1

	Y less than X	Y same as X	Y greater than X
centripetal acceleration			
maximum speed			
time taken for one lap of the track			

[3]

[Total: 7]

Appendix A

Q1.

1 **(a)** **(i)** Define the *radian*.

.....

.....

..... [2]

(ii) A small mass is attached to a string. The mass is rotating about a fixed point P at constant speed, as shown in Fig. 1.1.




Fig. 1.1

Explain what is meant by the *angular* speed about point P of the mass.

.....

.....

..... [2]

(b) A horizontal flat plate is free to rotate about a vertical axis through its centre, as shown in Fig. 1.2.




Fig. 1.2

A small mass M is placed on the plate, a distance *d* from the axis of rotation.
The speed of rotation of the plate is gradually increased from zero until the mass is seen to slide off the plate.

The maximum frictional force *F* between the plate and the mass is given by the expression

$$F = 0.72W,$$

where *W* is the weight of the mass M.
The distance *d* is 35 cm.

Determine the maximum number of revolutions of the plate per minute for the mass M to remain on the plate. Explain your working.

number = ...[5]

(c) The plate in **(b)** is covered, when stationary, with mud.
Suggest and explain whether mud near the edge of the plate or near the centre will first leave the plate as the angular speed of the plate is slowly increased.

.....

.....

.....[2]

Q2.

4 A vertical peg is attached to the edge of a horizontal disc of radius r, as shown in Fig. 4.1.

For Examiner's Use




Fig. 4.1

The disc rotates at constant angular speed ω. A horizontal beam of parallel light produces a shadow of the peg on a screen, as shown in Fig. 4.2.




Fig. 4.2 (plan view)

At time zero, the peg is at P, producing a shadow on the screen at S.
At time t, the disc has rotated through angle θ. The peg is now at R, producing a shadow at Q.

(a) Determine,

(i) in terms of ω and t, the angle θ,

..... [1]

(ii) in terms of ω, t and r, the distance SQ.

..... [1]

(b) Use your answer to **(a)(ii)** to show that the shadow on the screen performs simple harmonic motion.

.....

.....

..... [2]

(c) The disc has radius r of 12 cm and is rotating with angular speed ω of 4.7 rad s^{-1}.

Determine, for the shadow on the screen,

(i) the frequency of oscillation,

frequency = ... Hz [2]

(ii) its maximum speed.

speed = ... cm s^{-1} [2]

Q3.

1 **(a)** Define the *radian*.

.....

.....

.....[2]

(b) A stone of weight 3.0 N is fixed, using glue, to one end P of a rigid rod CP, as shown in Fig. 1.1.




Fig. 1.1

The rod is rotated about end C so that the stone moves in a vertical circle of radius 85 cm.
The angular speed ω of the rod and stone is gradually increased from zero until the glue snaps. The glue fixing the stone snaps when the tension in it is 18 N.

For the position of the stone at which the glue snaps,

(i) on the dotted circle of Fig. 1.1, mark with the letter S the position of the stone, [1]

(ii) calculate the angular speed ω of the stone.

angular speed = ... rad s^{-1} [4]

Q4.

1 A particle is following a circular path and is observed to have an angular displacement of 10.3°.

(a) Express this angle in radians (rad). Show your working and give your answer to three significant figures.

angle = ...rad [2]

(b) **(i)** Determine tan10.3° to three significant figures.

tan10.3° = ...

(ii) Hence calculate the percentage error that is made when the angle 10.3°, as measured in radians, is assumed to be equal to tan10.3°.

percentage error = ...
[3]

Q5.

1 **(a)** Explain

(i) what is meant by a *radian*,

.....

.....

.....[2]

(ii) why one complete revolution is equivalent to an angular displacement of 2π rad.

.....

.....[1]

(b) An elastic cord has an unextended length of 13.0 cm. One end of the cord is attached to a fixed point C. A small mass of weight 5.0 N is hung from the free end of the cord. The cord extends to a length of 14.8 cm, as shown in Fig. 1.1.




Fig. 1.1

The cord and mass are now made to rotate at constant angular speed ω in a vertical plane about point C. When the cord is vertical and above C, its length is the unextended length of 13.0 cm, as shown in Fig. 1.2.




Fig. 1.2




Fig. 1.3

(i) Show that the angular speed ω of the cord and mass is 8.7 rad s^{-1}.

[2]

(ii) The cord and mass rotate so that the cord is vertically below C, as shown in Fig. 1.3.

Calculate the length L of the cord, assuming it obeys Hooke's law.

L = ... cm [4]

Q6.

7 **(a)** Define the *radian*.

.....

.....

..... [2]

(b) A telescope gives a clear view of a distant object when the angular displacement between the edges of the object is at least 9.7×10^{-6} rad.

(i) The Moon is approximately 3.8×10^{5} km from Earth.
Estimate the minimum diameter of a circular crater on the Moon's surface that can be seen using the telescope.

diameter = ... km [2]

(ii) Suggest why craters of the same diameter as that calculated in **(i)** but on the surface of Mars are not visible using this telescope.

.....

.....

..... [2]

Q7.

2 A large bowl is made from part of a hollow sphere.

A small spherical ball is placed inside the bowl and is given a horizontal speed. The ball follows a horizontal circular path of constant radius, as shown in Fig. 2.1.




Fig. 2.1

The forces acting on the ball are its weight W and the normal reaction force R of the bowl on the ball, as shown in Fig. 2.2.




Fig. 2.2

The normal reaction force R is at an angle θ to the horizontal.

(a) (i) By resolving the reaction force R into two perpendicular components, show that the resultant force F acting on the ball is given by the expression

$$W = F \tan\theta.$$

[2]

(ii) State the significance of the force F for the motion of the ball in the bowl.

.....

..... [1]

(b) The ball moves in a circular path of radius 14 cm. For this radius, the angle θ is 28°.

Calculate the speed of the ball.

speed = ... $m\,s^{-1}$ [3]

Circular Motion - 2021 A2

1. Nov/2021/Paper_41/No.1

(a) With reference to velocity and acceleration, describe uniform circular motion.

.....

.....

..... [2]

(b) Two cars are moving around a horizontal circular track. One car follows path X and the other follows path Y, as shown in Fig. 1.1.




Fig. 1.1 (not to scale)

The radius of path X is 318 m. Path Y is parallel to, and 27 m outside, path X. Both cars have mass 790 kg. The maximum lateral (sideways) friction force F that the cars can experience without sliding is the same for both cars.

(i) The maximum speed at which the car on path X can move around the track without sliding is 94 m s^{-1}.

Calculate F.

F = ... N [2]

STEMUP.CO.UK

(ii) Both cars move around the track. Each car has the maximum speed at which it can move without sliding.

Complete Table 1.1, by placing one tick in each row, to indicate how the quantities indicated for the car on path Y compare with the car on path X.

Table 1.1

	Y less than X	Y same as X	Y greater than X
centripetal acceleration			
maximum speed			
time taken for one lap of the track			

[3]

[Total: 7]

STEMUP.CO.UK

2. Nov/2021/Paper_42/No.1

(a) State what is meant by *centripetal* acceleration.

.....

.....

..... [1]

(b) An unpowered toy car moves freely along a smooth track that is initially horizontal. The track contains a vertical circular loop around which the car travels, as shown in Fig. 1.1.




Fig. 1.1

The mass of the car is 230 g and the diameter of the loop is 62 cm. Assume that the resistive forces acting on the car are negligible.

(i) State what happens to the magnitude of the centripetal acceleration of the car as it moves around the loop from X to Y.

..... [1]

(ii) Explain, if the car remains in contact with the track, why the centripetal acceleration of the car at point Y must be greater than 9.8 m s^{-2}.

.....

.....

..... [2]

STEMUP.CO.UK

(c) The initial speed at which the car in **(b)** moves along the track is $3.8\,m\,s^{-1}$.

Determine whether the car is in contact with the track at point Y. Show your working.

[3]

(d) Suggest, with a reason but without calculation, whether your conclusion in **(c)** would be different for a car of mass 460 g moving with the same initial speed.

.....

.....

..... [1]

[Total: 8]

STEMUP.CO.UK

1. Nov/2023/Paper_9702/42/No.1

(a) Define the radian.

.....

..... [1]

(b) The minute hand of a clock revolves at constant angular speed around the face of the clock, completing one revolution every hour. A small piece of modelling clay is attached to the hand with its centre of gravity at a distance L from the fixed end of the hand, as shown in Fig. 1.1.




Fig. 1.1

Calculate the angular speed ω of the minute hand.

ω = ... rad s^{-1} [2]

STEMUP.CO.UK

(c) During a time interval of 1400 s, the centre of gravity of the piece of modelling clay in Fig. 1.1 moves through a total distance of 0.44 m.

(i) Calculate the angle through which the minute hand moves in this time interval.

angle = ... rad [1]

(ii) Determine distance *L*.

L = ... m [2]

(iii) Calculate the magnitude of the centripetal acceleration of the piece of modelling clay.

centripetal acceleration = ... $m\,s^{-2}$ [2]

(d) Use your answer in **(c)(iii)** to explain why the variation with time of the magnitude of the force exerted by the minute hand on the piece of modelling clay is negligible as the minute hand undergoes one full revolution.

.....

.....

..... [2]

[Total: 10]

STEMUP.CO.UK

2. June/2023/Paper_9702/41/No.2

A steel sphere of mass 0.29 kg is suspended in equilibrium from a vertical spring. The centre of the sphere is 8.5 cm from the top of the spring, as shown in Fig. 2.1.




Fig. 2.1

The sphere is now set in motion so that it is moving in a horizontal circle at constant speed, as shown in Fig. 2.2.




Fig. 2.2

The distance from the centre of the sphere to the top of the spring is now 10.8 cm.

(a) Explain, with reference to the forces acting on the sphere, why the length of the spring in Fig. 2.2 is greater than in Fig. 2.1.

.....

.....

.....

.....

..... [3]

STEMUP.CO.UK

(b) The angle between the linear axis of the spring and the vertical is 27°.

(i) Show that the radius r of the circle is 4.9 cm.

[1]

(ii) Show that the tension in the spring is 3.2 N.

[2]

(iii) The spring obeys Hooke's law.

Calculate the spring constant, in $\mathrm{N\,cm^{-1}}$, of the spring.

spring constant = ... $\mathrm{N\,cm^{-1}}$ [2]

(c) **(i)** Use the information in **(b)** to determine the centripetal acceleration of the sphere.

centripetal acceleration = ... $\mathrm{m\,s^{-2}}$ [2]

(ii) Calculate the period of the circular motion of the sphere.

period = ... s [2]

[Total: 12]

STEMUP.CO.UK