CAMBRIDGE INTERNATIONAL AS & A LEVEL PHYSICS · 9702

PHYSICS



MARK SCHEME

15 IDEAL GASES

SCORE	PERCENTAGE
/	%

STEMUP

These questions belong to CIE, not to us.

The examination questions, mark schemes and examiner comments in this pack were written and published by Cambridge International (CIE). They are past assessment materials, reproduced here so that students can revise from them.

What we do not own

STEMUP is not affiliated with CIE and does not own, and makes no claim to own, any of the question, mark scheme or examiner report material in this pack. CIE has not endorsed, approved or been involved in the production of it. CIE and the qualification titles used here belong to their respective owners and appear for identification only.

Where to find the originals

CIE publishes its past papers, mark schemes and examiner reports. Always check the current syllabus at cambridgeinternational.org before you revise.

If you hold the rights to any material in this pack and would like it amended or removed, contact admin@stemup.co.uk and we will act on it.

39. 9702_m17_MS_42 Q: 1

	Answer	Marks
(a)	work done per unit mass	M1
	bringing (small test) mass from infinity (to the point)	A1
(b)(i)	$\Delta\phi = (GM/2R) - (GM/5R) = 3GM/10R$	A1
(b)(ii)	change in GPE = $(3 \times 4.0 \times 10^{14} / 10R) \times 4.7 \times 10^4$	C1
	$(3 \times 4.0 \times 10^{14} / 10R) \times 4.7 \times 10^4 = (1.70 - 0.88) \times 10^{12}$ $R = 6.88 \times 10^6$	C1
	distance $= 3 \times 6.88 \times 10^6$ $= 2.1 \times 10^7$ m	A1

40. 9702_w19_MS_41 Q: 2

	Answer	Marks
(a)	(total volume of molecules is) negligible	M1
	compared with volume occupied by the gas	A1
(b)(i)	$pV = NkT$	C1
	$4.60 \times 10^5 \times 2.40 \times 10^{-2} = N \times 1.38 \times 10^{-23} \times (273 + 23)$	C1
	or	
	$pV = nRT$	(C1)
	$4.60 \times 10^5 \times 2.40 \times 10^{-2} = n \times 8.31 \times (273 + 23)$ $n = 4.49$ (mol) $N = nN_A$ $= 4.49 \times 6.02 \times 10^{23}$	(C1)
	$N = 2.7 \times 10^{24}$	A1

	Answer	Marks
(b)(ii)	volume of one atom = d^3 (= 2.7×10^{-29} m^3)	**C1**
	volume of all atoms = $2.7 \times 10^{-29} \times 2.7 \times 10^{24}$	**C1**
	= 7×10^{-5} m^3	**A1**
	or	
	volume of one atom = $(4/3)\pi r^3$ (= 1.41×10^{-29} m^3)	**(C1)**
	volume of all atoms = $2.7 \times 10^{24} \times 1.41 \times 10^{-29}$	**(C1)**
	= 4×10^{-5} m^3	**(A1)**
(c)	numerical comparison between answer to **(b)(ii)** and 2.4×10^{-2} (m^3) showing **(b)(ii)** is <u>much</u> less than 2.4×10^{-2} (m^3)	**B1**

41. 9702_w16_MS_42 Q: 2

(a) (i) number of moles/amount of substance B1 [1]

(ii) kelvin temperature/absolute temperature/thermodynamic temperature B1 [1]

(b) $pV = nRT$

$4.9 \times 10^5 \times 2.4 \times 10^3 \times 10^{-6} = n \times 8.31 \times 373$ B1

$n = 0.38$ (mol) C1

number of molecules or $N = 0.38 \times 6.02 \times 10^{23} = 2.3 \times 10^{23}$ A1 [3]

STEMUP.CO.UK

or

$pV = NkT$ (C1)

$4.9 \times 10^5 \times 2.4 \times 10^3 \times 10^{-6} = N \times 1.38 \times 10^{-23} \times 373$ (M1)

number of molecules or $N = 2.3 \times 10^{23}$ (A1)

volume occupied by one molecule $= (2.4 \times 10^3) / (2.3 \times 10^{23})$ C1

$= 1.04 \times 10^{-20}\ \text{cm}^3$

mean spacing $= (1.04 \times 10^{-20})^{1/3}$ C1

$= 2.2 \times 10^{-7}$ cm (*allow 1 s.f.*) A1 [3]

(*allow other dimensionally correct methods e.g.* $V = (4/3)\pi r^3$)

42. 9702_m21_MS_42 Q: 2

	Answer	Marks
(a)(i)	$pV = NkT$ or $pV = nRT$ and $N = nN_A$ $N = \frac{2.3\times10^5\times3.5\times10^{-3}}{1.38\times10^{-23}\times294}$	**C1**
	= 2.0 × 10²³	**A1**
(a)(ii)	$pV = \frac{1}{3}Nmc^2$ $c^2 = \frac{3\times2.3\times10^5\times3.5\times10^{-3}}{2.0\times10^{23}\times40\times1.66\times10^{-27}}$ = 182 000 r.m.s. speed = **430** m s⁻¹	**C1**
	or $\frac{1}{2}mc^2 = \frac{3}{2}kT$	**A1**
	$c^2 = \frac{3\times1.38\times10^{-23}\times294}{40\times1.66\times10^{-27}}$ = 183 000	**(C1)**
	r.m.s.speed = 430 m s⁻¹	**(A1)**

	Answer	Marks
(b)	$c^2 = \frac{3\times2.0\times10^{23}\times1.38\times10^{-23}\times(294+84)}{2.0\times10^{23}\times40\times1.66\times10^{-27}}$ $c^2 = 236000$ $c = 485$	C1
	$ratio\left(=\frac{485}{430}\right)=1.1$	A1
	OR $v \propto \sqrt{T}$ or $v^2 \propto T$	(C1)
	$ratio = \sqrt{\frac{273+21+84}{273+21}}$ or $\sqrt{\frac{378}{294}}$ ratio = **1.1**	(A1)

43. 9702_s20_MS_42 Q: 2

	Answer	Marks
(a)	ρ: Nm / V	B1
	⅓: molecules move in three dimensions (not one) so ⅓ in any (one) direction	B1
	$<c^2>$: molecules have different speeds so take average	M1
	of (speed)^2	A1
(b)	$pV = NkT$	C1
	$N = (3.0 \times 10^5 \times 6.0 \times 10^{-3}) / (1.38 \times 10^{-23} \times 290)$	C1
	$= 4.5 \times 10^{23}$	C1
	mass $= 20.7 / (4.5 \times 10^{23})$ $= 4.6 \times 10^{-23}$ g	A1

44. 9702_m19_MS_42 Q: 2

	Answer	Marks
(a)(i)	gas obeys formula pV / T = constant	M1
	symbols V and T explained	A1
(a)(ii)	mean-square-speed (of atoms / molecules)	B1
(b)(i)	use of T = 393	C1
	$pV = nRT$	C1
	$2.4 \times 10^5 \times 6.8 \times 10^{-3} = n \times 8.31 \times 393$ and $N = n \times 6.02 \times 10^{23} = 3.0 \times 10^{23}$	A1
	or $pV = NkT$	(C1)
	$2.4 \times 10^5 \times 6.8 \times 10^{-3} = N \times 1.38 \times 10^{-23} \times 393$ hence $N = 3.0 \times 10^{23}$	(A1)
(b)(ii)	volume of one atom = $4 / 3\pi r^3$	C1
	volume occupied $= 3.0 \times 10^{23} \times 4 / 3 \times \pi \times (3.2 \times 10^{-11})^3$ $= 4 \times 10^{-8}$ m^3	A1
(b)(iii)	assumption: volume of atoms negligible compared to volume of container / cylinder	B1
	4×10^{-8} (m^3) << 6.8×10^{-3} (m^3) so yes	B1

45. 9702_w19_MS_43 Q: 2

	Answer	Marks
(a)	(total volume of molecules is) negligible	M1
	compared with volume occupied by the gas	A1
(b)(i)	$pV = NkT$	C1
	$4.60 \times 10^5 \times 2.40 \times 10^{-2} = N \times 1.38 \times 10^{-23} \times (273 + 23)$	C1
	or	
	$pV = nRT$	(C1)
	$4.60 \times 10^5 \times 2.40 \times 10^{-2} = n \times 8.31 \times (273 + 23)$ $n = 4.49$ (mol) $N = nN_A$ $= 4.49 \times 6.02 \times 10^{23}$	(C1)
	$N = 2.7 \times 10^{24}$	A1

	Answer	Marks
(b)(ii)	volume of one atom $= d^3$ ($= 2.7 \times 10^{-29}$ m^3)	C1
	volume of all atoms $= 2.7 \times 10^{-29} \times 2.7 \times 10^{24}$	C1
	$= 7 \times 10^{-5}$ m^3	A1
	or	
	volume of one atom $= (4/3)\pi r^3$ ($= 1.41 \times 10^{-29}$ m^3)	(C1)
	volume of all atoms $= 2.7 \times 10^{24} \times 1.41 \times 10^{-29}$	(C1)
	$= 4 \times 10^{-5}$ m^3	(A1)
(c)	numerical comparison between answer to **(b)(ii)** and 2.4×10^{-2} (m^3) showing **(b)(ii)** is <u>much</u> less than 2.4×10^{-2} (m^3)	B1

46. 9702_w18_MS_42 Q: 2

	Answer	Marks
(a)	gas that obeys equation pV = constant $\times T$	M1
	symbols p, V and T explained	A1
(b)(i)	$pV = \frac{1}{3} Nm\langle c^2 \rangle$ **and** $M = Nm$ (and so) $p = \frac{1}{3}\rho\langle c^2 \rangle$	C1
	$2.12 \times 10^7 = \frac{1}{3} \times [3.20/(1.84 \times 10^{-2})] \times \langle c^2 \rangle$	C1
	$c_{\text{r.m.s.}} = 605\,\text{m s}^{-1}$	A1
(b)(ii)	**1.** $pV = nRT$ **and** $T = (22 + 273)$K	C1
	$n = (2.12 \times 10^7 \times 1.84 \times 10^{-2})/(8.31 \times 295)$ $= 159$ mol	A1
	2. mass $= 3.20/(159 \times 6.02 \times 10^{23})$ **or** mass $= [2 \times (3/2) \times 1.38 \times 10^{-23} \times 295]/605^2$	C1
	mass $= 3.34 \times 10^{-26}$ kg	A1
(c)	$A = (3.34 \times 10^{-26})/(1.66 \times 10^{-27})$ $= 20$	A1

47. 9702_s17_MS_43 Q: 4

	Answer	Marks
(a)	random/haphazard	B1
	constant velocity **or** speed in a straight line between collisions **or** distribution of speeds/different directions	B1
(b)	(small) specks of light/bright specks/pollen grains/dust particles/smoke particles	M1
	moving haphazardly/randomly/jerky/in a zigzag fashion	A1
(c)(i)	$pV = \frac{1}{3} Nm\langle c^2 \rangle$ $1.05 \times 10^5 \times 0.0240 = \frac{1}{3} \times 4.00 \times 10^{-3} \times \langle c^2 \rangle$	C1
	$\langle c^2 \rangle = 1.89 \times 10^6$	C1
	or	
	$\frac{1}{2} m\langle c^2 \rangle = (3/2)\, kT$ $0.5 \times (4.00 \times 10^{-3} / 6.02 \times 10^{23}) \times \langle c^2 \rangle = 1.5 \times 1.38 \times 10^{-23} \times 300$	(C1)
	$\langle c^2 \rangle = 1.87 \times 10^6$	(C1)
	or	
	$nRT = \frac{1}{3} Nm\langle c^2 \rangle$ $1.00 \times 8.31 \times 300 = \frac{1}{3} \times 4.00 \times 10^{-3} \times \langle c^2 \rangle$	(C1)
	$\langle c^2 \rangle = 1.87 \times 10^6$	(C1)
	$c_{r.m.s.} = 1.37 \times 10^3\ \text{m s}^{-1}$	A1

	Answer	Marks
(c)(ii)	$\langle c^2 \rangle \propto T$	C1
	$\langle c^2 \rangle$ at 177 °C $= 1.89 \times 10^6 \times (450 / 300)$	C1
	$c_{r.m.s.}$ at 177 °C $= 1.68 \times 10^3\ \text{m s}^{-1}$	A1

48. 9702_w21_MS_42 Q: 3

	Answer	Marks
(a)(i)	no loss of kinetic energy	B1
(a)(ii)	• molecules have negligible volume (compared with gas/container) • no forces between molecules (except during collisions) • molecules are in random motion • collisions are instantaneous *Any two points, 1 mark each*	B2
(b)(i)	$2mu$	A1
(b)(ii)	$2L / u$	A1
(b)(iii)	force = change in momentum / time $= 2mu / (2L / u)$ $= mu^2 / L$	A1
(b)(iv)	pressure = force / area $= (mu^2 / L) / L^2$ $= mu^2 / L^3$	A1
(c)	$pV = NkT$	C1
	$NkT = \frac{1}{3} Nm\langle c^2 \rangle$ leading to $\frac{1}{2} m\langle c^2 \rangle = (3/2)kT$ and $\frac{1}{2} m\langle c^2 \rangle = E_K$	A1
(d)	$\frac{1}{2} \times 3.34 \times 10^{-27} \times \langle c^2 \rangle = (3/2) \times 1.38 \times 10^{-23} \times (25 + 273)$	C1
	r.m.s. speed $= 1.9 \times 10^3\ \text{m s}^{-1}$	A1

49. 9702_s18_MS_42 Q: 2

	Answer	Marks
(a)	no intermolecular forces (so no potential energy)	B1
(b)(i)	mean square speed (of molecule(s))	B1
(b)(ii)	kelvin/thermodynamic/absolute temperature	B1
(c)(i)1.	$pV = NkT$	C1
	$4.7 \times 10^{-2} \times 2.6 \times 10^{5} = N \times 1.38 \times 10^{-23} \times 446$	C1
	or	
	$pV = nRT$ **and** $N = nN_A$ $4.7 \times 10^{-2} \times 2.6 \times 10^{5} = n \times 8.31 \times 446$ $n = 3.3$ (mol)	(C1)
	$N = 3.3 \times 6.02 \times 10^{23}$	(C1)
	$N = 2.0 \times 10^{24}$	A1
(c)(i)2.	average increase $= 2900 / (2.0 \times 10^{24})$ $= 1.5 \times 10^{-21}$ J	A1
(c)(ii)	$\Delta E_K = (3/2)k(\Delta)T$ $1.5 \times 10^{-21} = (3/2) \times 1.38 \times 10^{-23} \times (\Delta)T$	C1
	$(\Delta)T$ in range 70–72 K	C1
	$T = 173 + 273 + 70$ $= 520$ K	A1

50. 9702_s17_MS_41 Q: 4

	Answer	Marks
(a)	random/haphazard	B1
	constant velocity **or** speed in a straight line between collisions **or** distribution of speeds/different directions	B1
(b)	(small) specks of light/bright specks/pollen grains/dust particles/smoke particles	M1
	moving haphazardly/randomly/jerky/in a zigzag fashion	A1
(c)(i)	$pV = \frac{1}{3}Nm\langle c^2\rangle$ $1.05 \times 10^{5} \times 0.0240 = \frac{1}{3} \times 4.00 \times 10^{-3} \times \langle c^2\rangle$	C1
	$\langle c^2\rangle = 1.89 \times 10^{6}$	C1
	or	
	$\frac{1}{2}m\langle c^2\rangle = (3/2)kT$ $0.5 \times (4.00 \times 10^{-3} / 6.02 \times 10^{23}) \times \langle c^2\rangle = 1.5 \times 1.38 \times 10^{-23} \times 300$	(C1)
	$\langle c^2\rangle = 1.87 \times 10^{6}$	(C1)
	or	
	$nRT = \frac{1}{3}Nm\langle c^2\rangle$ $1.00 \times 8.31 \times 300 = \frac{1}{3} \times 4.00 \times 10^{-3} \times \langle c^2\rangle$	(C1)
	$\langle c^2\rangle = 1.87 \times 10^{6}$	(C1)
	$c_{r.m.s.} = 1.37 \times 10^{3}$ m s^{-1}	A1

	Answer	Marks
(c)(ii)	$\langle c^2 \rangle \propto T$	C1
	$\langle c^2 \rangle$ at 177 °C = $1.89 \times 10^6 \times (450 / 300)$	C1
	$c_{r.m.s.}$ at 177 °C = 1.68×10^3 m s^{-1}	A1

51. 9702_m21_MS_42 Q: 3

	Answer	Marks
(a)	Any 2 from: • particles / atoms / molecules / ions (very) close together / touching • regular, repeating pattern • vibrate about a fixed point	B2
(b)	(much) greater increase in spacing of molecules (for vaporisation compared with fusion)	B1
(c)(i)	–100 °C	B1

	Answer	Marks
(c)(ii)	time = 8.5 – 3.0 = 5.5 min	C1
	$Pt = mL$ energy = power × time = 150 × 5.5 × 60 = 49 500 J $L = \frac{E}{m}$ $= \frac{49500}{0.045}$	C1
	$= 1100\,kJ\,kg^{-1}$	A1
(c)(iii)	gas has a higher specific heat capacity (than liquid)	B1

52. 9702_m20_MS_42 Q: 2

	Answer	Marks
(a)	n = 110 / 0.032 or 110000 / 32 or 3440	C1
	$pV = nRT$	C1
	$T = (1.0 \times 10^5 \times 85) / (8.31 \times (110 / 0.032)) = 300$ K	A1
(b)	$E = mc\Delta\theta$ $= 110 \times 0.66 \times 50$	C1
	= **3600** J	A1
(c)	Any 3 from: • molecule collides with wall • momentum of molecule changes during collision (with wall) • force on molecule so force on wall • many forces act over surface area of container exerting a pressure	B3
(d)	$KE \propto T$ $v \propto \sqrt{T}$	C1
	ratio = √(350 / 300) = **1.1**	A1

Q1.

2	(a)		mean (value of the) square of the speeds (velocities) of the atoms/particles/molecules	M1 A1	[2]
	(b)	(i)	$p = \frac{1}{3} \rho \langle c^2 \rangle$	C1	
			$\langle c^2 \rangle = 3 \times 2 \times 10^5 / 2.4 = 2.5 \times 10^5$	C1	
			r.m.s speed = 500 ms^{-1}	A1	[3]
		(ii)	new $\langle c^2 \rangle = 1.0 \times 10^6$ or $\langle c^2 \rangle$ increases by factor of 4	C1	
			$\langle c^2 \rangle \propto T$ or $3/2\ kT = 1/2\ m\langle c^2 \rangle$	C1	
			$T = \{(1.0 \times 10^6) / (2.5 \times 10^5)\} \times 300$ $= 1200$ K	A1	[3]
				Total	[8]

Q2.

2	(a)		obeys the law pV/T = constant or any two named gas laws at all values of p, V and T or two correct assumptions of kinetic theory of ideal gas **(B1)** third correct assumption **(B1)**	**M1** **A1**	**[2]**
	(b)	(i)	mean square speed	**B1**	**[1]**
		(ii)	mean kinetic energy = $\frac{1}{2}m\langle c^2 \rangle$ $\rho = Nm/V$ and algebra leading to [do not allow if takes $N = 1$] $\frac{1}{2}m\langle c^2 \rangle = 3/2\ kT$	**M1** **M1** **A0**	**[2]**
	(c)	(i)	$\frac{1}{2} \times 6.6 \times 10^{-27} \times (1.1 \times 10^4)^2 = 3/2 \times 1.38 \times 10^{-23} \times T$ $T = 1.9 \times 10^4$ K	**C1** **A1**	**[2]**
		(ii)	Not all atoms have same speed/kinetic energy	**B1**	**[1]**

Q3.

2	(a)		e.g. fixed mass/ amount of gas ideal gas *(any two, 1 each)*	B2	[2]
	(b)	(i)	$n = pV / RT$ $= (2.5 \times 10^7 \times 4.00 \times 10^4 \times 10^{-6}) / (8.31 \times 290)$ $= 415$ mol	C1 C1 A1	[3]
		(ii)	volume of gas at 1.85×10^5 Pa $= (2.5 \times 10^7 \times 4.00 \times 10^4) / (1.85 \times 10^5)$ $= 5.41 \times 10^6$ cm^3 so, $5.41 \times 10^6 = 4.00 \times 10^4 + 7.24 \times 10^3\ N$ $N = 741$ *(answer 740 or fails to allow for gas in cylinder, max 2/3)*	C1 C1 A1	[3]

Q4.

2 (a) molecule(s) rebound from wall of vessel / hits walls — B1
change in momentum gives rise to impulse / force — B1
either (many impulses) averaged to give constant force / pressure
or the molecules are in random motion — B1 [3]

(b) (i) $p = \frac{1}{3}\rho\langle c^2\rangle$ — C1

$1.02 \times 10^5 = \frac{1}{3} \times 0.900 \times \langle c^2\rangle$

$\langle c^2\rangle = 3.4 \times 10^5$ — C1
$c_{RMS} = 580\ m\ s^{-1}$ — A1 [3]

(ii) *either* $\langle c^2\rangle \propto T$ *or* $\langle c^2\rangle = 2 \times 3.4 \times 10^5$ — C1
$c_{RMS} = 830\ m\ s^{-1}$ (allow 820) — A1 [2]

(c) c_{RMS} depends on temperature (alone) — B1
so no effect — B1 [2]

Q5.

2 (a) *either* the half-life of the source is <u>very</u> long
or decay constant is <u>very</u> small
or half-life >> 40 days
or decay constant << 0.02 day^{-1} — B1 [1]

(b) number of helium atoms $= 3.5 \times 10^6 \times 40 \times 24 \times 3600$ — C1
$= 1.21 \times 10^{13}$
either $pV = NkT$ *or* $pV = nRT$ <u>*and*</u> $n = N / N_A$ — C1
$1.5 \times 10^5 \times V = 1.21 \times 10^{13} \times 1.38 \times 10^{-23} \times 290$
$V = 3.2 \times 10^{-13}\ m^3$ — A1 [3]
(if uses T/°C or $n = 1$ or $n = 4$, then 1 mark max for calculation of number of atoms)

Q6.

2	(a) (i)	27.2 + 273.15 or 27.2 + 273.2	C1		
		300.4 K	A1	[2]	
	(ii)	11.6 K	A1	[1]	
	(b) (i)	($<c^2>$ is the) mean / average square speed	B1	[1]	
	(ii)	$\rho = Nm/V$ with N explained	B1		
		so, $pV = 1/3\ Nm<c^2>$	B1		
		and $pV = NkT$ with k explained	B1		
		so mean kinetic energy / $<E_K> = \frac{1}{2}m<c^2> = 3/2\ kT$	B1	[4]	
	(c) (i)	$pV = nRT$			
		$2.1 \times 10^7 \times 7.8 \times 10^{-3} = n \times 8.3 \times 290$	C1		
		$n = 68$ mol	A1	[2]	
	(ii)	mean kinetic energy $= 3/2\ kT$			
		$= 3/2 \times 1.38 \times 10^{-23} \times 290$	C1		
		$= 6.0 \times 10^{-21}$ J	A1	[2]	
	(iii)	realisation that total internal energy is the total kinetic energy	C1		
		energy $= 6.0 \times 10^{-21} \times 68 \times 6.02 \times 10^{23}$	C1		
		$= 2.46 \times 10^5$ J	A1	[3]	

Q7.

2	(a)	number of atoms of carbon-12	M1	
		in 0.012 kg of carbon-12	A1	[2]
	(b)	$pV = NkT$ or $pV = nRT$	C1	
		substitutes temperature as 298 K	C1	
		either $1.1 \times 10^5 \times 6.5 \times 10^{-2} = N \times 1.38 \times 10^{-23} \times 298$		
		or $1.1 \times 10^5 \times 6.5 \times 10^{-2} = n \times 8.31 \times 298$ and $n = N / 6.02 \times 10^{23}$	C1	
		$N = 1.7 \times 10^{24}$	A1	[4]

Q8.

2	(a)	amount of substance	M1	
		containing same number of particles as in 0.012 kg of carbon-12	A1	[2]
	(b)	$pV = nRT$	C1	
		amount = $(2.3 \times 10^5 \times 3.1 \times 10^{-3}) / (8.31 \times 290)$		
		+ $(2.3 \times 10^5 \times 4.6 \times 10^{-3}) / (8.31 \times 303)$	C1	
		= 0.296 + 0.420	C1	
		= 0.716 mol	A1	[4]
		(*give full credit for starting equation* $pV = NkT$ *and* $N = nN_A$)		

Q9.

2	(a)	(i)	*either* random motion *or* constant velocity until hits wall/other molecule	B1	[1]
		(ii)	(total) volume of molecules is negligible compared to volume of containing vessel *or* radius/diameter of a molecule is negligible compared to the average intermolecular distance	M1 A1 (M1) (A1)	[2]
	(b)		*either* molecule has component of velocity in three directions *or* $c^2 = c_X^2 + c_Y^2 + c_Z^2$ random motion and averaging, so $\langle c_X^2 \rangle = \langle c_Y^2 \rangle = \langle c_Z^2 \rangle$ $\langle c^2 \rangle = 3\langle c_X^2 \rangle$ so, $pV = \frac{1}{3}Nm\langle c^2 \rangle$	 M1 M1 A1 A0	[3]
	(c)		$\langle c^2 \rangle \propto T$ or $c_{rms} \propto \sqrt{T}$ temperatures are 300 K and 373 K $c_{rms} = 580\,m\,s^{-1}$ *(Do not allow any marks for use of temperature in units of °C instead of K)*	C1 C1 A1	[3]

Q10.

2	(a)		obeys the equation pV = constant × T or $pV = nRT$ p, V and T explained at all values of p, V and T/fixed mass/n is constant	M1 A1 A1	[3]
	(b)	(i)	$3.4 \times 10^5 \times 2.5 \times 10^3 \times 10^{-6} = n \times 8.31 \times 300$ $n = 0.34$ mol	M1 A0	[1]
		(ii)	for total mass/amount of gas $3.9 \times 10^5 \times (2.5 + 1.6) \times 10^3 \times 10^{-6} = (0.34 + 0.20) \times 8.31 \times T$ $T = 360$ K	C1 A1	[2]
	(c)		when tap opened gas passed (from cylinder B) to cylinder A work done <u>on</u> gas in cylinder A (and no heating) so internal energy and hence temperature increase	 B1 M1 A1	[3]

Q11.

2 (a) (i) 1. $pV = nRT$
$1.80 \times 10^{-3} \times 2.60 \times 105 = n \times 8.31 \times 297$ C1
$n = 0.19$ mol A1 [2]

2. $\Delta q = mc\Delta T$
$95.0 = 0.190 \times 12.5 \times \Delta T$ B1
$\Delta T = 40$ K A1 [2]
(allow 2 marks for correct answer with clear logic shown)

(ii) p/T = constant
$(2.6 \times 10^5) / 297 = p / (297 + 40)$ M1
$p = 2.95 \times 10^5$ Pa A0 [1]

(b) change in internal energy is 120 J / 25 J B1
internal energy decreases / ΔU is negative / kinetic energy of molecules decreases M1
so temperature lower A1 [3]

Q12.

2 (a) obeys the law pV = constant $\times$ T M1
at all values of p, V and T ..... A1 **[2]**

(b) $n = (2.9 \times 10^5 \times 3.1 \times 10^{-2}) / (8.31 \times 290)$ C1
$= 3.73$ mol ..... A1 **[2]**

(c) at new pressure, $n_n = 3.73 \times \frac{3.4}{2.9} \times \frac{290}{300}$
$= 4.23$ mol C1
change = 0.50 mol C1
number of strokes = 0.50 / 0.012 = 42 (must round up for mark) ..... A1 **[3]**

Q13.

2 (a) (i) $pV = nRT$
$V = (8.31 \times 300)/(1.02 \times 10^5)$ C1
$= 0.0244$ m^3 (if uses Celsius, then 0/2) .....A1 [2]

(ii) volume occupied by one atom = $0.0244 / (6.02 \times 10^{23}) = 4.06 \times 10^{-26}$ m^3 .....M1
separation $\approx \sqrt[3]{(4.06 \times 10^{-26})}$ A1
$= 3.44 \times 10^{-9}$ m .....A0 [2]

(b) (i) $F = GMm / r^2$ C1
$= (6.67 \times 10^{-11} \times \{4 \times 1.66 \times 10^{-27}\}^2) / (3.44 \times 10^{-9})^2$ C1
$= 2.49 \times 10^{-46}$ N .....A1 [3]

(ii) ratio $= (4 \times 1.66 \times 10^{-27} \times 9.8) / 2.49 \times 10^{-46}$ C1
$= 2.6 \times 10^{20}$ .....A1 [2]

(c) assumption that forces between atoms are negligible B1
comment e.g. ratio shows gravitational force to be very small
e.g. force is very much less than weight
e.g. if there are forces, they are not gravitational .....B1 [2]

Q14.

5 (a) change/loss in kinetic energy = change/gain in electric potential energy — B1

$2 \times \frac{1}{2}mv^2 = q^2 / 4\pi\varepsilon_0 r$ — C1

$2 \times \frac{1}{2} \times 2 \times 1.67 \times 10^{-27} \times v^2$

$= (1.6 \times 10^{-19})^2 / (4\pi \times 8.85 \times 10^{-12} \times 1.1 \times 10^{-14})$ — M1

$v = 2.5 \times 10^6$ m s^{-1} — A0 [3]

(b) $pV = \frac{1}{2}Nm\langle c^2\rangle$ and $pV = NkT$ — C1

$\frac{1}{2} m\langle c^2\rangle = \frac{3}{2} kT$ (*award 1 mark of first two if $\langle c^2\rangle$ not used*) — C1

$\frac{1}{2} \times 2 \times 1.67 \times 10^{-27} \times (2.5 \times 10^6)^2 = \frac{3}{2} \times 1.38 \times 10^{-23} \times T$ — C1

$T = 5 \times 10^8$ K — A1 [4]

(c) e.g. this is very high temperature
temperature found in stars
(*any sensible comment, 1 mark*)
(*if $T < 10^6$ K, should comment that too low for fusion to occur*) — B1 [1]

Q15.

2 (a) *either* $pV = NkT$ *or* $pV = nRT$ *and* $n = N / N_A$ C1

clear correct substitution e.g.

$2.5 \times 10^5 \times 4.5 \times 10^3 \times 10^{-6} = N \times 1.38 \times 10^{-23} \times 290$ M1

$N = 2.8 \times 10^{23}$ ..... A0 [2]

(*allow 1 mark for calculation of $n = 0.467$ mol*)

(b) (i) volume $= (1.2 \times 10^{-10})^3 \times 2.8 \times 10^{23}$ *or* $\frac{4}{3}\pi r^3 \times 2.8 \times 10^{23}$ C1

$= 4.8 \times 10^{-7}$ m^3 2.53×10^{-7} m^3 ..... A1 [2]

(ii) *either* 4.5×10^3 cm^3 >> 0.48 cm^3 *or* ratio of volumes is about 10^{-4} B1

justified because volume of molecules is negligible ..... B1 [2]

[Total: 6]

Q16.

2 (a) (i) no forces (of attraction or repulsion) between atoms / molecules / particles B1 [1]

(ii) sum of kinetic and potential energy of atoms / molecules M1
due to random motion A1 [2]

(iii) (random) kinetic energy increases with temperature M1
no potential energy
(so increase in temperature increases internal energy) A1 [2]

(b) (i) zero A1 [1]

(ii) work done $= p\Delta V$ C1
$= 4.0 \times 10^5 \times 6 \times 10^{-4}$
$= 240\,\text{J}$ *(ignore any sign)* A1 [2]

(iii)

change	work done / J	heating / J	increase in internal energy / J
P → Q	**+240**	–600	**–360**
Q → R	0	+720	**+720**
R → P	**–840**	+480	**–360**

(*correct signs essential*)
(*each horizontal line correct, 1 mark – max 3*) B3 [3]

Q17.

2 (a) atoms / molecules / particles behave as elastic (identical) spheres (1)
volume of atoms / molecules negligible compared to volume of containing vessel (1)
time of collision negligible to time between collisions (1)
no forces of attraction or repulsion between atoms / molecules (1)
atoms / molecules / particles are in (continuous) random motion (1)
(*any four, 1 each*) B4 [4]

(b) $pV = \frac{1}{3}Nm\langle c^2\rangle$ and $pV = nRT$ *or* $pV = NkT$ B1
$\frac{1}{3}Nm\langle c^2\rangle = nRT$ or $= NkT$ and $\langle E_K\rangle = \frac{1}{2}m\langle c^2\rangle$ B1
$n = N/N_A$ *or* $k = R/N_A$ B1
$\langle E_K\rangle = \frac{3}{2} \times R/N_A \times T$ A0 [3]

(c) (i) reaction represents *either* build-up of nucleus from light nuclei
or build-up of heavy nucleus from nuclei M1
so fusion reaction A1 [2]

(ii) proton and deuterium nucleus will have equal kinetic energies B1
$1.2 \times 10^{-14} = \frac{3}{2} \times 8.31 / (6.02 \times 10^{23}) \times T$ C1
$T = 5.8 \times 10^8\,\text{K}$ A1 [3]
(*use of* $E = 2.4 \times 10^{-14}$ *giving* 1.16×10^9 *K scores 1 mark*)

(iii) *either* inter-molecular / atomic / nuclear forces exist
or proton and deuterium nucleus are positively charged / repel B1 [1]

Q18.

2 **(a)** e.g. moving in random (rapid) motion of molecules/atoms/particles
no intermolecular forces of attraction/repulsion
volume of molecules/atoms/particles negligible compared to volume of container
time of collision negligible to time between collisions
(1 each, max 2) B2 [2]

(b) **(i)** **1.** number of (gas) molecules B1 [1]

2. mean square speed/velocity (of gas molecules) B1 [1]

(ii) either $pV = NkT$ *or* $pV = nRT$ and links n and k
and $\langle E_K \rangle = \frac{1}{2}m\langle c^2 \rangle$ M1

clear algebra leading to $\langle E_K \rangle = \frac{3}{2}kT$ A1 [2]

(c) **(i)** sum of potential energy and kinetic energy of molecules/atoms/particles M1
reference to random (distribution) A1 [2]

(ii) no intermolecular forces so no potential energy B1
(change in) internal energy is (change in) kinetic energy and this is proportional to (change in) T B1 [2]

Q19.

(c) **(i)** $\frac{1}{2} \times 2 \times 1.66 \times 10^{-27} \times (5.03 \times 10^{3})^2 = \frac{3}{2} \times 1.38 \times 10^{-23} \times T$ C1

$T = 2030$ K A1 [2]

(ii) *either* because there is a range of speeds M1
some molecules have a higher speed A1
or some escape from point above planet surface (M1)
so initial potential energy is higher (A1) [2]

Q20.

2 **(a)** **(i)** sum of potential energy and kinetic energy of atoms/molecules/particles M1
reference to random A1 [2]

(ii) no intermolecular forces B1
no potential energy B1
internal energy is kinetic energy (of random motion) of molecules B1 [3]
(*reference to random motion here then allow back credit to* ***(i)*** *if M1 scored*)

(b) kinetic energy $\propto$ thermodynamic temperature B1
either temperature in Celsius, not kelvin so incorrect
or temperature in kelvin is not doubled B1 [2]

Q21.

1	(a)	(i)	number of molecules	B1	[1]
		(ii)	mean square speed	B1	[1]
	(b)	(i)	1. $pV = nRT$	C1	
			$n = (6.1 \times 10^5 \times 2.1 \times 10^4 \times 10^{-6}) / (8.31 \times 285)$	C1	
			$n = 5.4$ mol	A1	[3]
			2. *either* $N = nN_A$		
			$= 5.4 \times 6.02 \times 10^{23}$	C1	
			$= 3.26 \times 10^{24}$	A1	
			or		
			$pV = NkT$		
			$N = (6.1 \times 10^5 \times 2.1 \times 10^4 \times 10^{-6}) / (1.38 \times 10^{-23} \times 285)$	(C1)	
			$N = 3.26 \times 10^{24}$	(A1)	[2]
		(ii)	*either* $6.1 \times 10^5 \times 2.1 \times 10^{-2} = {}^1/_3 \times 3.25 \times 10^{24} \times 4 \times 1.66 \times 10^{-27} \times \langle c^2 \rangle$	C1	
			$\langle c^2 \rangle = 1.78 \times 10^6$	C1	
			$c_{RMS} = 1.33 \times 10^3$ m s^{-1}	A1	
			or		
			${}^1/_2 \times 4 \times 1.66 \times 10^{-27} \times \langle c^2 \rangle = {}^3/_2 \times 1.38 \times 10^{-23} \times 285$	(C1)	
			$\langle c^2 \rangle = 1.78 \times 10^6$	(C1)	
			$c_{RMS} = 1.33 \times 10^3$ m s^{-1}	(A1)	[3]

Q22.

2	(a)	(i)	N: (total) number of molecules	B1	[1]
		(ii)	$\langle c^2 \rangle$: mean square speed/velocity	B1	[1]
	(b)		$pV = \frac{1}{3}Nm\langle c^2 \rangle = NkT$		
			(mean) kinetic energy = $\frac{1}{2} m\langle c^2 \rangle$	C1	
			algebra clear leading to $\frac{1}{2} m\langle c^2 \rangle = (3/2)kT$	A1	[2]
	(c)	(i)	*either* energy required = $(3/2) \times 1.38 \times 10^{-23} \times 1.0 \times 6.02 \times 10^{23}$	C1	
			$= 12.5$ J (*12J if 2 s.f.*)	A1	[2]
			or energy = $(3/2) \times 8.31 \times 1.0$	(C1)	
			$= 12.5$ J	(A1)	
		(ii)	energy is needed to push back atmosphere/do work against atmosphere	M1	
			so total energy required is greater	A1	[2]

Q23.

2 (a) (i) sum of kinetic and potential energies of the molecules — M1
reference to random distribution — A1 [2]

(ii) for ideal gas, no intermolecular forces — M1
so no potential energy (only kinetic) — A1 [2]

(b) (i) *either* change in kinetic energy $= 3/2 \times 1.38 \times 10^{-23} \times 1.0 \times 6.02 \times 10^{23} \times 180$ — C1
$= 2240\,\text{J}$ — A1 [2]

or $R = kN_A$
energy $= 3/2 \times 1.0 \times 8.31 \times 180$ — (C1)
$= 2240\ \text{J}$ — (A1)

(ii) increase in internal energy = heat supplied + work done on system — B1
2240 = energy supplied – 1500 — C1
energy supplied = 3740 J — A1 [3]

Q24.

2 (a) the number of atoms
in 12 g of carbon-12 — M1 A1 [2]

(b) (i) amount = 3.2/40
$= 0.080$ mol — A1 [1]

(ii) $pV = nRT$
$p \times 210 \times 10^{-6} = 0.080 \times 8.31 \times 310$ — C1
$p = 9.8 \times 10^5$ Pa — A1 [2]
(do not credit if T in °C not K)

(iii) *either* $pV = 1/3 \times Nm \langle c^2 \rangle$
$N = 0.080 \times 6.02 \times 10^{23}$ $(= 4.82 \times 10^{22})$
and $m = 40 \times 1.66 \times 10^{-27}$ $(= 6.64 \times 10^{-26})$ — C1
$9.8 \times 10^5 \times 210 \times 10^{-6} = 1/3 \times 4.82 \times 10^{22} \times 6.64 \times 10^{-26} \times \langle c^2 \rangle$ — C1
$\langle c^2 \rangle = 1.93 \times 10^5$
$c_{RMS} = 440\ \text{m s}^{-1}$ — A1 [3]

or $Nm = 3.2 \times 10^{-3}$ — (C1)
$9.8 \times 10^5 \times 210 \times 10^{-6} = 1/3 \times 3.2 \times 10^{-3} \times \langle c^2 \rangle$ — (C1)
$\langle c^2 \rangle = 1.93 \times 10^5$
$c_{RMS} = 440\ \text{m s}^{-1}$ — (A1)

or $1/2\ m\langle c^2 \rangle = 3/2\ kT$ — (C1)
$1/2 \times 40 \times 1.66 \times 10^{-27} \langle c^2 \rangle = 3/2 \times 1.38 \times 10^{-23} \times 310$ — (C1)
$\langle c^2 \rangle = 1.93 \times 10^5$
$c_{RMS} = 440\ \text{m s}^{-1}$ — (A1)

(*if T in °C not K award max 1/3, unless already penalised in* ***(b)(ii)***)

Q25.

2 (a) use of kelvin temperatures B1
both values of (V / T) correct (11.87), V / T is constant so pressure is constant M1 [2]

(allow use of n =1. Do not allow other values of n.)

(b) (i) work done $= p\Delta V$
$= 4.2 \times 10^5 \times (3.87 - 3.49) \times 10^3 \times 10^{-6}$ C1
$= 160$ J A1 [2]

(do not allow use of V instead of ΔV)

(ii) increase / change in internal energy = heating of system
+ work done on system C1
$= 565 - 160$
$= 405$ J A1 [2]

(c) internal energy = sum of kinetic energy and potential energy / $E_K + E_P$ B1
no intermolecular forces M1
no potential energy (so $\Delta U = \Delta E_K$) A1 [3]

Q26.

3 (a) obeys the equation pV/T = constant B1 [1]
(accept $pV = nRT$)

(b) (i) $pV = nRT$ C1
$5.0 \times 10^7 \times 3.0 \times 10^{-4} = n \times 8.31 \times 296$ giving $n = 6.1$ mol A1 [2]

(ii) pressure $\propto$ amount of substance
loss = 0.40 / 100 × 6.1 mol = 0.0244 mol C1
$= 0.0244 \times 6.02 \times 10^{23}$ (atoms) C1
$= 1.47 \times 10^{22}$ atoms C1

rate = $(1.47 \times 10^{22}) / (35 \times 24 \times 60 \times 60)$
$= 4.9 \times 10^{15}\ s^{-1}$ A1 [4]