



Cambridge International AS & A Level

CHEMISTRY

9701/11

Paper 1 Multiple Choice

October/November 2024

1 hour 15 minutes

You must answer on the multiple choice answer sheet.

You will need: Multiple choice answer sheet
Soft clean eraser
Soft pencil (type B or HB is recommended)

INSTRUCTIONS

- There are **forty** questions on this paper. Answer **all** questions.
- For each question there are four possible answers **A**, **B**, **C** and **D**. Choose the **one** you consider correct and record your choice in soft pencil on the multiple choice answer sheet.
- Follow the instructions on the multiple choice answer sheet.
- Write in soft pencil.
- Write your name, centre number and candidate number on the multiple choice answer sheet in the spaces provided unless this has been done for you.
- Do **not** use correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.

INFORMATION

- The total mark for this paper is 40.
- Each correct answer will score one mark.
- Any rough working should be done on this question paper.
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **20** pages. Any blank pages are indicated.



- 1 In this question Q is used to represent a halogen atom.

Magnesium and calcium each form a compound with chlorine and a compound with bromine.

One of these compounds contains:

- the element in Group 2 with the higher first ionisation energy **and**
- the element in Group 17 with the higher Q–Q bond energy.

What is the formula of this compound?

- A** MgCl_2 **B** MgBr_2 **C** CaCl_2 **D** CaBr_2

- 2 Compound X contains two elements, Y and Z.

Element Y is in Period 2 of the Periodic Table. In one atom of element Y, the p sub-shell has all three orbitals occupied; only one of these three orbitals is fully occupied.

Element Z is in Period 3 of the Periodic Table. In one atom of element Z, the p sub-shell has only two orbitals occupied.

What is the formula of compound X?

- A** CCl_4 **B** SiCl_4 **C** SiO_2 **D** SO_2

- 3 Glauber's salt consists of crystals of hydrated sodium sulfate, $\text{Na}_2\text{SO}_4 \cdot x\text{H}_2\text{O}$, which can be used for the manufacture of detergents.

When a sample of Glauber's salt was heated, 1.91 g of water was removed leaving 1.51 g of anhydrous Na_2SO_4 .

What is the value of x in $\text{Na}_2\text{SO}_4 \cdot x\text{H}_2\text{O}$?

- A** 1 **B** 8.85 **C** 10 **D** 11.25

- 4 What contains the greatest number of the named particles?

- A** 6.0 dm^3 of argon atoms at room conditions
B 6.0 g of carbon dioxide molecules
C 6.0 g of magnesium atoms
D 6.0 g of water molecules





- 5 Phosphorus forms a compound with hydrogen called phosphine, PH_3 . This compound can react with a hydrogen ion, H^+ .

Which type of interaction occurs between PH_3 and H^+ ?

- A dative covalent bond
- B dipole–dipole forces
- C hydrogen bond
- D ionic bond

- 6 The graphs show trends in four physical properties of elements in Period 3, excluding argon.

Which graph has electronegativity on the y -axis?



- 7 The element tin exists in two forms, grey tin and white tin.

Some properties of grey tin and white tin are shown.

	grey tin	white tin
boiling point/K	2543	2533
electrical conductivity	none in solid or liquid	good in solid and liquid
malleability	brittle	malleable

Which structural change might take place when grey tin changes to white tin?

- A** giant covalent to giant ionic
B giant covalent to giant metallic
C giant ionic to giant covalent
D giant ionic to giant metallic
- 8 Which solid has a simple molecular lattice?
- A** calcium fluoride
B nickel
C silicon(IV) oxide
D sulfur

- 9 The standard enthalpy change of combustion of carbon is -394 kJ mol^{-1} .

The standard enthalpy change of combustion of hydrogen is -286 kJ mol^{-1} .

The standard enthalpy change of formation of butane is -129 kJ mol^{-1} .

What is the standard enthalpy change of combustion of butane?

- A** -551 kJ mol^{-1}
B $-2877 \text{ kJ mol}^{-1}$
C $-3135 \text{ kJ mol}^{-1}$
D $-4307 \text{ kJ mol}^{-1}$

10 Three processes are described.

- 1 $\text{H}^+(\text{aq}) + \text{OH}^-(\text{aq}) \rightarrow \text{H}_2\text{O}(\text{l})$
- 2 $\text{CH}_4(\text{g}) + 2\text{O}_2(\text{g}) \rightarrow \text{CO}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l})$
- 3 $\text{NH}_3(\text{g}) \rightarrow \text{NH}_3(\text{l})$

Which statement is correct?


- A** None of the processes have a positive enthalpy change.
- B** Only process 1 has a positive enthalpy change.
- C** Only process 2 has a positive enthalpy change.
- D** Only process 3 has a positive enthalpy change.
- 11 In alkaline solution, MnO_4^- ions oxidise SO_3^{2-} ions to SO_4^{2-} ions. The MnO_4^- ions are reduced to MnO_2 .

What is the ratio of the two ions in the balanced chemical equation for this reaction?

	MnO_4^-	SO_3^{2-}
A	2	3
B	3	2
C	4	7
D	7	4

12 Lithium reacts with nitrogen at room temperature to form solid Li_3N .

Three vessels of equal volume are connected by taps 1 and 2 as shown.




At the start, taps 1 and 2 are closed, the left-hand vessel is evacuated, the middle vessel has the indicated reaction at equilibrium and the right-hand vessel contains lithium only.

Which action would allow the equilibrium mixture to contain the **most** ammonia?

- A** Keep both taps 1 and 2 closed.
- B** Open both taps 1 and 2.
- C** Open tap 1 only.
- D** Open tap 2 only.

- 13 When 0.20 mol of hydrogen gas and 0.15 mol of iodine gas are heated at 723 K until equilibrium is established, the equilibrium mixture contains 0.26 mol of hydrogen iodide.


The equation for the reaction is as follows.





What is the correct expression for the equilibrium constant K_c ?

- A $\frac{2 \times 0.26}{0.20 \times 0.15}$ B $\frac{(2 \times 0.26)^2}{0.20 \times 0.15}$ C $\frac{(0.26)^2}{0.07 \times 0.02}$ D $\frac{(0.26)^2}{0.13 \times 0.13}$

- 14 In acidic conditions, iodine reacts with propanone in a substitution reaction.



The kinetics of the reaction are investigated using a colorimeter. As the I_2 reacts, the yellow/brown colour of the $\text{I}_2(\text{aq})$ fades to colourless, changing the absorbance of the solution. Known concentrations of $\text{I}_2(\text{aq})$ are used to prepare a calibration curve graph and the absorbance is then measured as the reaction proceeds.




What is the rate of reaction at 20 s?

- A $5 \times 10^{-6} \text{ mol dm}^{-3} \text{ s}^{-1}$
 B $1 \times 10^{-5} \text{ mol dm}^{-3} \text{ s}^{-1}$
 C $5 \times 10^{-3} \text{ mol dm}^{-3} \text{ s}^{-1}$
 D $1 \times 10^{-2} \text{ mol dm}^{-3} \text{ s}^{-1}$

15 The diagram shows a Boltzmann distribution curve.

The axes are not labelled.



Points X and Y are points on the vertical axis.

What is represented by both points X and Y?

	point X	point Y
A	number of molecules with energy equal to E_A	largest number of molecules with the same energy
B	number of molecules with energy equal to or greater than E_A	largest number of molecules with the same energy
C	number of molecules with energy equal to E_A	the amount of energy of the greatest number of molecules
D	number of molecules with energy equal to or greater than E_A	the amount of energy of the greatest number of molecules

16 What are the acid–base nature and structure of SO_2 ?

	acid–base nature	structure
A	acidic	giant covalent lattice
B	acidic	simple molecular
C	basic	giant covalent lattice
D	basic	simple molecular

- 17** Elements X and Y are in Period 3 of the Periodic Table. Element X is either phosphorus or sulfur. Element Y is either sodium or magnesium.

Element X forms an oxide that reacts with water to give a solution containing the aqueous anion XO_4^{2-} .

One mole of element Y reacts with one mole of chlorine molecules. At the end of the reaction, all of the element Y and all of the chlorine molecules have been used up.

What are elements X and Y?

	X	Y
A	phosphorus	sodium
B	phosphorus	magnesium
C	sulfur	sodium
D	sulfur	magnesium

- 18** Q is a semi-conductor. The chloride of Q reacts with water to form white fumes and an acidic solution.

Which Period 3 element is Q?

- A** magnesium
B aluminium
C silicon
D phosphorus

- 19** V and W are two compounds. Each one contains a different Group 2 element.

A sample of each solid is added to water, shaken, and the pH of the resulting solution is measured.

compound	V	W
pH	13.6	9.4

Which row could identify V and W?

	V	W
A	BaSO_4	MgSO_4
B	MgSO_4	BaSO_4
C	Ba(OH)_2	Mg(OH)_2
D	Mg(OH)_2	Ba(OH)_2

- 20 Compound L decomposes on heating. One of the products is gas M.

M reacts with unburned hydrocarbons to form peroxyacetyl nitrate, PAN.

What could be the formula of L?

- A** CaNO_3 **B** $\text{Ca}(\text{NO}_3)_2$ **C** MgCO_3 **D** $\text{Mg}(\text{CO}_3)_2$

- 21 In reaction 1, concentrated sulfuric acid is added to potassium chloride and the fumes produced are bubbled into aqueous potassium iodide solution.

In reaction 2, potassium chloride is dissolved in aqueous ammonia and this is then added to aqueous silver nitrate.

What are the observations for reactions 1 and 2?

	observation for reaction 1	observation for reaction 2
A	brown solution	colourless solution
B	brown solution	white precipitate
C	colourless solution	colourless solution
D	colourless solution	white precipitate

- 22 The table refers to the hydrogen halides.

Which row is correct?

	oxidation	thermal stability
A	easier to oxidise down the group	increases down the group
B	more difficult to oxidise down the group	increases down the group
C	easier to oxidise down the group	decreases down the group
D	more difficult to oxidise down the group	decreases down the group

- 23 7.5 g of nitrogen monoxide reacts with 7.0 g of carbon monoxide on the surface of the catalytic converter in the exhaust system of a car.

What is the total volume of the product gases measured at room conditions?

- A** 3.0 dm^3 **B** 6.0 dm^3 **C** 9.0 dm^3 **D** 12.0 dm^3


24 Three statements about ammonia molecules and ammonium ions are given.

- 1 In aqueous solution, ammonia molecules form coordinate bonds with hydroxide ions.
- 2 Ammonium ions are Brønsted–Lowry acids.
- 3 The H–N–H bond angle is larger in the ammonium ion than in the ammonia molecule.

Which statements are correct?

- A** 1 and 2 only **B** 1 and 3 only **C** 2 and 3 only **D** 1, 2 and 3

25 Ethene reacts with steam in the presence of sulfuric acid.




Which type of reaction is this?

- A** acid–base
B addition
C hydrolysis
D substitution

26 Compound Z has the molecular formula $\text{C}_4\text{H}_8\text{O}_2$.

Compound Z reacts with propan-1-ol in the presence of concentrated H_2SO_4 .



The diagram shows the skeletal formulae of three compounds, S, T and U.



What are the possible skeletal formulae of the products of the reaction between compound Z and propan-1-ol?

- A** S and T **B** U only **C** S and U **D** T only

27 Geraniol and nerol are isomers of each other.



Which type of isomerism is shown here?


- A chain
- B geometrical (cis / trans)
- C optical
- D positional

28 Which compound has the greatest number of stereoisomers?

- A 2-methylhex-2-ene
- B 3-methylhex-2-ene
- C 4-methylhex-2-ene
- D 5-methylhex-2-ene

29 Vitamin A contains retinol.

retinol



Under appropriate conditions, acidified $\text{KMnO}_4(\text{aq})$ can be used to break $\text{C}=\text{C}$ bonds.


After these bonds have been broken, further oxidation of the fragments may occur.

Under which conditions is the acidified $\text{KMnO}_4(\text{aq})$ used and what do the final oxidation products include?

	conditions	final oxidation products
A	cold, dilute	aldehydes and carboxylic acids
B	cold, dilute	ketones and carboxylic acids
C	hot, concentrated	aldehydes and carboxylic acids
D	hot, concentrated	ketones and carboxylic acids

30 The structure of limonene is shown.

limonene



What are the number of moles of carbon dioxide and water produced when a sample of limonene is completely combusted in oxygen?

	number of moles of carbon dioxide	number of moles of water
A	4	3
B	5	4
C	5	8
D	9	7


- 31 The reaction of chlorine with methane is carried out in the presence of light.

What is the function of the light?

- A to break the C–H bonds in methane
- B to break the chlorine molecules into atoms
- C to break the chlorine molecules into ions
- D to heat the mixture

- 32 When X is added to NaOH(aq) and heated under reflux, pentan-2-ol is made.

Which organic product is made when X is heated with a solution of KCN dissolved in ethanol?



- 33 1-chlorobutane and 1-iodobutane both react with aqueous sodium hydroxide by a nucleophilic substitution mechanism.

Which reaction has the greatest rate under the same conditions and which mechanism is followed by this reaction?


	greatest rate	mechanism
A	1-chlorobutane	S _N 1
B	1-chlorobutane	S _N 2
C	1-iodobutane	S _N 1
D	1-iodobutane	S _N 2

- 34 Compound Y reacts with alkaline I₂(aq). When the products of this reaction are acidified, a dicarboxylic acid is produced. The formula of the dicarboxylic acid is HOOC–R–COOH where R consists of one or more CH₂ groups.

Which compound is Y?

- A pentan-1,4-diol
- B pentan-1,5-diol
- C pentan-2,3-diol
- D pentan-2,4-diol

- 35 Which alcohol gives only **one** possible oxidation product when warmed with dilute acidified potassium dichromate(VI)?
- A butan-1-ol
 B butan-2-ol
 C 2-methylpropan-1-ol
 D 2-methylpropan-2-ol
- 36 Which compound, on reaction with hydrogen cyanide, produces a compound with a chiral centre?
- A CH_3CHO
 B $\text{CH}_3\text{CH}_2\text{COCH}_2\text{CH}_3$
 C $\text{CH}_3\text{CO}_2\text{CH}_3$
 D HCHO
- 37 The diagram shows three reactions of ethanal. In each case, an excess of ethanal is used.




Observations are made after each of the three reactions.

What are the colours of solution 1 and solids 2 and 3?

	solution 1	solid 2	solid 3
A	green	yellow	silver mirror
B	green	yellow	red
C	orange	red	silver mirror
D	orange	red	red


38 $(\text{CH}_3)_3\text{CCN}$ reacts to form alcohol Y via the reaction sequence shown.



Which row names the molecule X and the class of alcohol Y?

	name of molecule X	class of alcohol Y
A	2,2-dimethylbutanoic acid	primary
B	3,3-dimethylbutanoic acid	tertiary
C	dimethylpropanoic acid	primary
D	dimethylpropanoic acid	tertiary


39 The diagram shows a section of an addition polymer. The polymer is made using two different monomers.



What are the names of the two monomers needed to make this polymer?

- A** 1,2-dichloropropene and 2-chlorobut-2-ene
- B** 2,3-dichlorobut-2-ene and chloropropene
- C** 1,2-dichloropropene and chloroethene
- D** chloropropene and 2-chlorobut-2-ene

- 40 The diagram shows the mass spectrum of a sample of chlorine. Peaks V, W, X, Y and Z are labelled.



Which statements about this spectrum are correct?

- 1 The relative atomic mass of chlorine can be calculated from the abundances and m/e values of 2 of the 5 peaks.
- 2 37.0 g of the species responsible for peak Z contains 3.011×10^{23} molecules.
- 3 The relative molecular mass of chlorine can be calculated from the abundances and m/e values of peaks X, Y and Z.

A 1, 2 and 3 **B** 1 and 2 only **C** 1 and 3 only **D** 2 and 3 only

BLANK PAGE

BLANK PAGE

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 $\text{J g}^{-1} \text{ K}^{-1}$)

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

The Periodic Table of Elements

Group																							
1	2													13	14	15	16	17	18				
		Key atomic number atomic symbol name relative atomic mass																1 H hydrogen 1.0					
3 Li lithium 6.9	4 Be beryllium 9.0																	5 B boron 10.8	6 C carbon 12.0	7 N nitrogen 14.0	8 O oxygen 16.0	9 F fluorine 19.0	10 Ne neon 20.2
11 Na sodium 23.0	12 Mg magnesium 24.3																	13 Al aluminium 27.0	14 Si silicon 28.1	15 P phosphorus 31.0	16 S sulfur 32.1	17 Cl chlorine 35.5	18 Ar argon 39.9
19 K potassium 39.1	20 Ca calcium 40.1	21 Sc scandium 45.0	22 Ti titanium 47.9	23 V vanadium 50.9	24 Cr chromium 52.0	25 Mn manganese 54.9	26 Fe iron 55.8	27 Co cobalt 58.9	28 Ni nickel 58.7	29 Cu copper 63.5	30 Zn zinc 65.4	31 Ga gallium 69.7	32 Ge germanium 72.6	33 As arsenic 74.9	34 Se selenium 79.0	35 Br bromine 79.9	36 Kr krypton 83.8						
37 Rb rubidium 85.5	38 Sr strontium 87.6	39 Y yttrium 88.9	40 Zr zirconium 91.2	41 Nb niobium 92.9	42 Mo molybdenum 95.9	43 Tc technetium —	44 Ru ruthenium 101.1	45 Rh rhodium 102.9	46 Pd palladium 106.4	47 Ag silver 107.9	48 Cd cadmium 112.4	49 In indium 114.8	50 Sn tin 118.7	51 Sb antimony 121.8	52 Te tellurium 127.6	53 I iodine 126.9	54 Xe xenon 131.3						
55 Cs caesium 132.9	56 Ba barium 137.3	57–71 lanthanoids		72 Hf hafnium 178.5	73 Ta tantalum 180.9	74 W tungsten 183.8	75 Re rhenium 186.2	76 Os osmium 190.2	77 Ir iridium 192.2	78 Pt platinum 195.1	79 Au gold 197.0	80 Hg mercury 200.6	81 Tl thallium 204.4	82 Pb lead 207.2	83 Bi bismuth 209.0	84 Po polonium —	85 At astatine —	86 Rn radon —					
87 Fr francium —	88 Ra radium —	89–103 actinoids		104 Rf rutherfordium —	105 Db dubnium —	106 Sg seaborgium —	107 Bh bohrium —	108 Hs hassium —	109 Mt meitnerium —	110 Ds darmstadtium —	111 Rg roentgenium —	112 Cn copernicium —	113 Nh nihonium —	114 Fl flerovium —	115 Mc moscovium —	116 Lv livermorium —	117 Ts tennessine —	118 Og oganesson —					

lanthanoids

57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.2	61 Pm promethium —	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0
--------------------------------	-----------------------------	-----------------------------------	--------------------------------	-----------------------------	-------------------------------	-------------------------------	---------------------------------	------------------------------	---------------------------------	------------------------------	-----------------------------	------------------------------	--------------------------------	-------------------------------

actinoids

89 Ac actinium —	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium —	94 Pu plutonium —	95 Am americium —	96 Cm curium —	97 Bk berkelium —	98 Cf californium —	99 Es einsteinium —	100 Fm fermium —	101 Md mendelevium —	102 No nobelium —	103 Lr lawrencium —
---------------------------	------------------------------	-----------------------------------	-----------------------------	----------------------------	----------------------------	----------------------------	-------------------------	----------------------------	------------------------------	------------------------------	---------------------------	-------------------------------	----------------------------	------------------------------



Cambridge International AS & A Level

CHEMISTRY

9701/12

Paper 1 Multiple Choice

October/November 2024

1 hour 15 minutes

You must answer on the multiple choice answer sheet.

You will need: Multiple choice answer sheet
Soft clean eraser
Soft pencil (type B or HB is recommended)

INSTRUCTIONS

- There are **forty** questions on this paper. Answer **all** questions.
- For each question there are four possible answers **A**, **B**, **C** and **D**. Choose the **one** you consider correct and record your choice in soft pencil on the multiple choice answer sheet.
- Follow the instructions on the multiple choice answer sheet.
- Write in soft pencil.
- Write your name, centre number and candidate number on the multiple choice answer sheet in the spaces provided unless this has been done for you.
- Do **not** use correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.

INFORMATION

- The total mark for this paper is 40.
- Each correct answer will score one mark.
- Any rough working should be done on this question paper.
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **16** pages.



1 Which species contains a different number of electrons from the other three?

- A** ClO_4^- **B** H_2SO_4 **C** SO_4^{2-} **D** Te^{2-}

2 Which factor causes helium to have a higher first ionisation energy than hydrogen?

- A** In the 1s orbital in helium, electrons are paired.
B The lowest energy level in helium is filled.
C The nuclear charge in helium is higher than in hydrogen.
D There is less shielding of the outer shell in helium.

3 A 0.216 g sample of aluminium carbide reacts with an excess of water to produce methane gas. This is the only carbon-containing product formed in the reaction. This methane gas burns completely in O_2 to form H_2O and CO_2 only. The volume of CO_2 produced at room temperature and pressure is 108 cm^3 .

What is the formula of aluminium carbide?

- A** Al_2C_3 **B** Al_3C_2 **C** Al_3C_4 **D** Al_4C_3

4 A reaction between two gases takes place on the surface of the catalytic converter of a petrol-engined car.

In this reaction, four reactant molecules produce three product molecules.

What could be the two reactant gases in this reaction?

- A** nitrogen and carbon dioxide
B nitrogen monoxide and carbon dioxide
C nitrogen monoxide and carbon monoxide
D nitrogen dioxide and carbon monoxide

5 An ion contains 1 nitrogen atom and 2 hydrogen atoms. It has an H–N–H bond angle of approximately 105° .

Which row is correct?

	number of lone pairs around N in ion	overall charge on ion
A	1	+1
B	2	+1
C	1	–1
D	2	–1

- 6 Why does ICl have a higher boiling point than Br_2 ?
- A because of the difference in the bond energies of the covalent bonds within ICl and Br_2
- B because of the difference in the polar nature of ICl and Br_2
- C because of the difference in the number of electrons contained within ICl and Br_2
- D because of the difference in the relative molecular mass of ICl and Br_2
- 7 In this question you may assume that nitrogen behaves as an ideal gas. One atmosphere pressure = 101 kPa.
- Which volume does 1.0 g of nitrogen occupy at 50°C and a pressure of 2.0 atmospheres?
- A 70 cm^3 B 150 cm^3 C 470 cm^3 D 950 cm^3
- 8 Which statement about the properties associated with the different types of bonding involved is correct?
- A Any covalent compound that contains both oxygen and hydrogen in its molecule forms hydrogen bonds.
- B Ionic bonds and covalent bonds cannot both occur in the same compound.
- C Ionic compounds differ from metals in that ionic compounds do not conduct electricity in the solid state.
- D The only covalent compounds with high melting points are those in which hydrogen bonds occur.
- 9 For which reaction is the enthalpy change an enthalpy change of formation?
- A $\text{C(g)} + 2\text{H}_2\text{(g)} \rightarrow \text{CH}_4\text{(g)}$
- B $\frac{1}{2}\text{N}_2\text{(g)} + \frac{1}{2}\text{O}_2\text{(g)} \rightarrow \text{NO(g)}$
- C $\text{Na}_2\text{O(s)} + \text{SO}_3\text{(g)} \rightarrow \text{Na}_2\text{SO}_4\text{(s)}$
- D $\text{PCl}_3\text{(g)} + \text{Cl}_2\text{(g)} \rightarrow \text{PCl}_5\text{(g)}$

- 10 Two standard enthalpy change of formation values are given.




$$\Delta H_f^\ominus [\text{VCl}_2] = -452\text{ kJ mol}^{-1}$$

$$\Delta H_f^\ominus [\text{VCl}_3] = -573\text{ kJ mol}^{-1}$$

What is the enthalpy change for the reaction $3\text{VCl}_2 \rightarrow 2\text{VCl}_3 + \text{V}$?

- A -210 kJ mol^{-1} B -121 kJ mol^{-1} C $+121\text{ kJ mol}^{-1}$ D $+210\text{ kJ mol}^{-1}$


11 Equations for some reactions of hydrogen peroxide are given.



In which reactions is hydrogen peroxide acting as a reducing agent?

- A** 1 and 3 **B** 1 only **C** 2 and 3 **D** 2 only

12 The equation for the reaction of aqueous thiosulfate ions, $\text{S}_2\text{O}_3^{2-}$, and aqueous dioxo-vanadium ions, VO_2^+ , is shown.



Which row shows two correct statements about the equation for this reaction?

	comparison of x and y to z	change in oxidation number of vanadium
A	x and z are the same value and quarter the value of y	from +4 to +5
B	x and z are the same value and quarter the value of y	from +5 to +4
C	x and z are the same value and half the value of y	from +5 to +4
D	x and z are the same value and half the value of y	from +4 to +5


13 When some solid $\text{Ca}_5(\text{PO}_4)_3\text{OH}$ is added to a beaker of water, an equilibrium is set up.



Which compound, when added to the equilibrium mixture, increases the amount of $\text{Ca}_5(\text{PO}_4)_3\text{OH}(\text{s})$ present?

- A** NH_3 **B** NH_4Cl **C** $\text{CH}_3\text{CO}_2\text{H}$ **D** NaCl

- 14 Gaseous hydrogen and gaseous iodine react to form gaseous hydrogen iodide.



In an experiment, 2.0 mol of hydrogen and 2.0 mol of iodine are placed in a sealed container of volume 1.0 dm^3 .

The K_c value for this reaction under the conditions used is 9.0.

How many moles of hydrogen iodide are present at equilibrium?


- A** 0.57 mol **B** 1.2 mol **C** 1.5 mol **D** 2.4 mol

- 15 Why does the rate of a gaseous reaction increase when the pressure is increased at a constant temperature?

- A** More particles have energy that exceeds the activation energy.
B The particles have more space in which to move.
C The particles move faster.
D There are more frequent collisions between particles.

- 16 The Boltzmann distribution for a mixture of gases capable of reaction is shown.

The two curves represent the mixture of gases at 25°C and at 35°C . The activation energies for the catalysed and uncatalysed reactions are shown.



Which row is correct?

	number of particles with enough energy to react at 25°C in the catalysed reaction	number of particles with enough energy to react at 35°C in the uncatalysed reaction
A	$w + x + y + z$	z
B	$w + x + y + z$	$x + z$
C	$y + z$	z
D	$y + z$	$x + z$

17 Which oxide is insoluble in aqueous sodium hydroxide?

- A** MgO **B** Al_2O_3 **C** P_4O_{10} **D** SO_2

18 Sodium and sulfur are burned separately in oxygen.

Each reaction has a distinctive coloured flame.

Which row is correct?

	Na + O ₂	S + O ₂
A	white flame	blue flame
B	white flame	yellow flame
C	yellow flame	blue flame
D	yellow flame	yellow flame

19 X and Y are elements in Period 3 of the Periodic Table.

Y has a greater atomic number than X.

The stable ion formed by Y has a greater radius than the stable ion formed by X.

The stable ion formed by Y has 18 electrons.

Which row is correct?

	number of electrons in the stable ion of X	element with the greater atomic radius
A	10	X
B	10	Y
C	18	X
D	18	Y

- 20 X is a Group 2 element in either Period 3 or Period 5. $X(OH)_2$ is less soluble in water than $Ca(OH)_2$.

When $X(NO_3)_2$ is heated, it decomposes.

Which row is correct?

	identity of X	equation describing decomposition of $X(NO_3)_2$
A	Mg	$X(NO_3)_2 \rightarrow X + 2NO_2 + O_2$
B	Mg	$2X(NO_3)_2 \rightarrow 2XO + 4NO_2 + O_2$
C	Sr	$X(NO_3)_2 \rightarrow X + 2NO_2 + O_2$
D	Sr	$2X(NO_3)_2 \rightarrow 2XO + 4NO_2 + O_2$

- 21 Which statement comparing magnesium and barium, or their compounds, is correct?

- A Magnesium reacts with dilute hydrochloric acid more rapidly than barium does.
- B One mole of magnesium carbonate gives off a greater amount of gas when it reacts with an excess of dilute hydrochloric acid than one mole of barium carbonate does.
- C The solubility of magnesium sulfate in water is greater than the solubility of barium sulfate in water.
- D Magnesium carbonate undergoes thermal decomposition **less** readily than barium carbonate does.

- 22 The colours of the silver halides $AgCl$, $AgBr$ and AgI differ.

The solubilities of these halides in aqueous ammonia also differ.

Which row is correct?

	colour of $AgBr$	silver halide that is most soluble in $NH_3(aq)$
A	cream	$AgCl$
B	cream	AgI
C	yellow	$AgCl$
D	yellow	AgI

- 23** The name 'chlorate' is used for an anion consisting of chlorine and oxygen only.

In a molecule of ICl , the iodine atom has oxidation number x and the chlorine atom has oxidation number y .

When ICl is added to H_2O , iodine is reduced.




Which statement about the value of x or y is correct?

- A** x is the same as the oxidation number of Cl in the chlorate ion formed when $Cl_2(aq)$ is added to cold $NaOH(aq)$.
- B** x is the same as the oxidation number of Cl in the chlorate ion formed when $Cl_2(aq)$ is added to hot $NaOH(aq)$.
- C** y is the same as the oxidation number of Cl in the chlorate ion formed when $Cl_2(aq)$ is added to cold $NaOH(aq)$.
- D** y is the same as the oxidation number of Cl in the chlorate ion formed when $Cl_2(aq)$ is added to hot $NaOH(aq)$.
- 24** Which statement is correct?
- A** An ammonium ion is basic due to a lone pair of electrons on the nitrogen atom.
- B** Nitrogen monoxide, NO , reacts with peroxyacetyl nitrate to produce a component of photochemical smog.
- C** Nitrogen dioxide catalyses the oxidation of atmospheric sulfur dioxide.
- D** Nitrogen is very unreactive due to the very strong permanent dipole–permanent dipole attractions between the nitrogen atoms.

- 25** The diagram shows the structural formula of a hydrocarbon molecule Q.

molecule Q




How many of the carbon atoms in molecule Q are sp^2 hybridised?

- A** 3 **B** 4 **C** 7 **D** 10

- 26 Compound X is found in cell walls of some bacteria. Its structural formula is shown.

compound X



How many stereoisomers are there with this structural formula?


- A 2 B 4 C 6 D 8
- 27 Structural isomerism **only** should be considered when answering this question.

How many straight-chain isomers are there with molecular formula $\text{C}_4\text{H}_8\text{Cl}_2$?

- A 6 B 7 C 8 D 9
- 28 What is true of **every** nucleophile?

- A It attacks a double bond.
 B It donates a lone pair of electrons.
 C It is a single atom.
 D It is negatively charged.


- 29 The diagram shows a synthetic route to produce 1-methylcyclohexanol.



What is reagent Y?

- A aqueous NaOH
 B cold dilute KMnO_4
 C ethanolic NaOH
 D hot concentrated KMnO_4

30 X and Y are the reagents required to convert 1-bromopropane into butanoic acid.



What are the correct identities of reagents X and Y?

	X	Y
A	NH ₃	HCl(aq)
B	KCN in C ₂ H ₅ OH	NaOH(aq)
C	KCN in C ₂ H ₅ OH	HCl(aq)
D	HCN	NaOH(aq)

31 The table shows three sets of reagents and reaction conditions.

	reagents	reaction conditions
1	CH ₂ C(CH ₃)CH ₃ and HCl(g)	room temperature
2	CH ₃ C(CH ₃)(OH)CH ₃ and SOCl ₂	room temperature
3	CH ₃ CH(CH ₃)CH ₃ and Cl ₂	the presence of ultraviolet light

Which sets of reagents and conditions can be used to produce 2-chloro-2-methylpropane as one of the organic products?


A 1, 2 and 3 **B** 1 and 2 only **C** 1 and 3 only **D** 2 and 3 only

32 What are the **only** structures formed when butan-2-ol is heated with concentrated H_2SO_4 ?




A	$\begin{array}{c} \text{CH}_3\text{CH}_2 \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{H} \end{array}$ $\begin{array}{c} \text{CH}_3 \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{CH}_3 \end{array}$
B	$\begin{array}{c} \text{CH}_3 \quad \text{CH}_3 \\ \diagdown \quad \diagup \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{H} \end{array}$ $\begin{array}{c} \text{CH}_3 \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{CH}_3 \end{array}$ $\begin{array}{c} \text{H} \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{CH}_3\text{CH}_2 \quad \text{H} \end{array}$
C	$\begin{array}{c} \text{H} \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{CH}_3\text{CH}_2 \quad \text{H} \end{array}$ $\begin{array}{c} \text{H} \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{CH}_3 \quad \text{CH}_3 \end{array}$
D	$\begin{array}{c} \text{CH}_3 \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{CH}_3 \quad \text{H} \end{array}$ $\begin{array}{c} \text{CH}_3\text{CH}_2 \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{H} \end{array}$ $\begin{array}{c} \text{H} \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{CH}_3 \quad \text{CH}_3 \end{array}$

33 The compound 'leaf alcohol' is partly responsible for the smell of new-mown grass.

leaf alcohol



What will be formed when 'leaf alcohol' is oxidised using an excess of hot acidified $\text{K}_2\text{Cr}_2\text{O}_7(\text{aq})$?



34 Compound X:

- does **not** react with Tollens' reagent
- forms a yellow precipitate with alkaline $I_2(aq)$
- does **not** react with sodium.

What could be the identity of X?

- A CH_3CHO
- B $C_2H_5COCH_3$
- C $CH_3COOC_2H_5$
- D $CH_3CHOHCH_3$

35 Which compound can undergo nucleophilic addition?

- A bromoethane, C_2H_5Br
- B ethanal, CH_3CHO
- C ethane, C_2H_6
- D ethene, C_2H_4

36 $C_2H_5COOCH_3$ is reacted with aqueous acid.

The products from this reaction are reacted with $LiAlH_4$ to form two molecules Y and Z.

What are the identities of molecules Y and Z?

- A both molecules are C_2H_5OH
- B CH_3OH and $CH_3CHOHCH_3$
- C CH_3OH and C_2H_5OH
- D CH_3OH and $C_2H_5CH_2OH$

37 A sample of propanoic acid of mass 3.70 g reacts with an excess of magnesium.

A second sample of propanoic acid of mass 3.70 g reacts with an excess of sodium.

Both reactions go to completion forming a gas.

Which row is correct?





	volume of gas formed with magnesium at s.t.p. / cm ³	volume of gas formed with sodium at s.t.p. / cm ³
A	560	560
B	560	1120
C	1120	560
D	1120	1120

38 Which statement about $\text{H}_2\text{C}=\text{C}(\text{CH}_3)\text{CH}_2\text{CO}_2\text{CH}_3$ is correct?


- A** It can be hydrolysed to a secondary alcohol.
- B** It can be made using ethanoic acid and a suitable alcohol.
- C** It gives a positive test with alkaline $\text{I}_2(\text{aq})$.
- D** When treated with hot concentrated acidified KMnO_4 it gives $\text{CH}_3\text{COCH}_2\text{COOH}$ as one product.

39 Synthetic resins can be made by polymerisation of a variety of monomers including prop-2-en-1-ol, $\text{CH}_2=\text{CHCH}_2\text{OH}$.

Which structure represents the repeat unit in the polymer poly(prop-2-en-1-ol)?



40 Vitamin C has the structure shown.



The mass spectrum of vitamin C has a molecular ion peak with an m/e value of 176 and a relative abundance of 7.0%.

What is the abundance of the $M + 1$ peak?

- A** 0.462% **B** 0.539% **C** 0.616% **D** 0.693%

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ ($4.18 \text{ J g}^{-1} \text{ K}^{-1}$)

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.



Cambridge International AS & A Level

CHEMISTRY

9701/13

Paper 1 Multiple Choice

October/November 2024

1 hour 15 minutes

You must answer on the multiple choice answer sheet.

You will need: Multiple choice answer sheet
Soft clean eraser
Soft pencil (type B or HB is recommended)

INSTRUCTIONS

- There are **forty** questions on this paper. Answer **all** questions.
- For each question there are four possible answers **A**, **B**, **C** and **D**. Choose the **one** you consider correct and record your choice in soft pencil on the multiple choice answer sheet.
- Follow the instructions on the multiple choice answer sheet.
- Write in soft pencil.
- Write your name, centre number and candidate number on the multiple choice answer sheet in the spaces provided unless this has been done for you.
- Do **not** use correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.

INFORMATION

- The total mark for this paper is 40.
- Each correct answer will score one mark.
- Any rough working should be done on this question paper.
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **20** pages. Any blank pages are indicated.



- 1 In this question Q is used to represent a halogen atom.

Magnesium and calcium each form a compound with chlorine and a compound with bromine.

One of these compounds contains:

- the element in Group 2 with the higher first ionisation energy **and**
- the element in Group 17 with the higher Q–Q bond energy.

What is the formula of this compound?

- A** MgCl_2 **B** MgBr_2 **C** CaCl_2 **D** CaBr_2

- 2 Compound X contains two elements, Y and Z.

Element Y is in Period 2 of the Periodic Table. In one atom of element Y, the p sub-shell has all three orbitals occupied; only one of these three orbitals is fully occupied.

Element Z is in Period 3 of the Periodic Table. In one atom of element Z, the p sub-shell has only two orbitals occupied.

What is the formula of compound X?

- A** CCl_4 **B** SiCl_4 **C** SiO_2 **D** SO_2

- 3 Glauber's salt consists of crystals of hydrated sodium sulfate, $\text{Na}_2\text{SO}_4 \cdot x\text{H}_2\text{O}$, which can be used for the manufacture of detergents.

When a sample of Glauber's salt was heated, 1.91 g of water was removed leaving 1.51 g of anhydrous Na_2SO_4 .

What is the value of x in $\text{Na}_2\text{SO}_4 \cdot x\text{H}_2\text{O}$?

- A** 1 **B** 8.85 **C** 10 **D** 11.25

- 4 What contains the greatest number of the named particles?

- A** 6.0 dm^3 of argon atoms at room conditions
B 6.0 g of carbon dioxide molecules
C 6.0 g of magnesium atoms
D 6.0 g of water molecules





- 5 Phosphorus forms a compound with hydrogen called phosphine, PH_3 . This compound can react with a hydrogen ion, H^+ .

Which type of interaction occurs between PH_3 and H^+ ?

- A dative covalent bond
- B dipole–dipole forces
- C hydrogen bond
- D ionic bond

- 6 The graphs show trends in four physical properties of elements in Period 3, excluding argon.

Which graph has electronegativity on the y -axis?



- 7 The element tin exists in two forms, grey tin and white tin.

Some properties of grey tin and white tin are shown.

	grey tin	white tin
boiling point/K	2543	2533
electrical conductivity	none in solid or liquid	good in solid and liquid
malleability	brittle	malleable

Which structural change might take place when grey tin changes to white tin?

- A** giant covalent to giant ionic
B giant covalent to giant metallic
C giant ionic to giant covalent
D giant ionic to giant metallic
- 8 Which solid has a simple molecular lattice?
- A** calcium fluoride
B nickel
C silicon(IV) oxide
D sulfur

- 9 The standard enthalpy change of combustion of carbon is -394 kJ mol^{-1} .

The standard enthalpy change of combustion of hydrogen is -286 kJ mol^{-1} .

The standard enthalpy change of formation of butane is -129 kJ mol^{-1} .

What is the standard enthalpy change of combustion of butane?

- A** -551 kJ mol^{-1}
B $-2877 \text{ kJ mol}^{-1}$
C $-3135 \text{ kJ mol}^{-1}$
D $-4307 \text{ kJ mol}^{-1}$

10 Three processes are described.

- 1 $\text{H}^+(\text{aq}) + \text{OH}^-(\text{aq}) \rightarrow \text{H}_2\text{O}(\text{l})$
- 2 $\text{CH}_4(\text{g}) + 2\text{O}_2(\text{g}) \rightarrow \text{CO}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l})$
- 3 $\text{NH}_3(\text{g}) \rightarrow \text{NH}_3(\text{l})$

Which statement is correct?


- A** None of the processes have a positive enthalpy change.
- B** Only process 1 has a positive enthalpy change.
- C** Only process 2 has a positive enthalpy change.
- D** Only process 3 has a positive enthalpy change.
- 11 In alkaline solution, MnO_4^- ions oxidise SO_3^{2-} ions to SO_4^{2-} ions. The MnO_4^- ions are reduced to MnO_2 .

What is the ratio of the two ions in the balanced chemical equation for this reaction?

	MnO_4^-	SO_3^{2-}
A	2	3
B	3	2
C	4	7
D	7	4

12 Lithium reacts with nitrogen at room temperature to form solid Li_3N .

Three vessels of equal volume are connected by taps 1 and 2 as shown.




At the start, taps 1 and 2 are closed, the left-hand vessel is evacuated, the middle vessel has the indicated reaction at equilibrium and the right-hand vessel contains lithium only.

Which action would allow the equilibrium mixture to contain the **most** ammonia?

- A** Keep both taps 1 and 2 closed.
- B** Open both taps 1 and 2.
- C** Open tap 1 only.
- D** Open tap 2 only.

- 13 When 0.20 mol of hydrogen gas and 0.15 mol of iodine gas are heated at 723 K until equilibrium is established, the equilibrium mixture contains 0.26 mol of hydrogen iodide.


The equation for the reaction is as follows.





What is the correct expression for the equilibrium constant K_c ?

- A $\frac{2 \times 0.26}{0.20 \times 0.15}$ B $\frac{(2 \times 0.26)^2}{0.20 \times 0.15}$ C $\frac{(0.26)^2}{0.07 \times 0.02}$ D $\frac{(0.26)^2}{0.13 \times 0.13}$

- 14 In acidic conditions, iodine reacts with propanone in a substitution reaction.



The kinetics of the reaction are investigated using a colorimeter. As the I_2 reacts, the yellow/brown colour of the $\text{I}_2(\text{aq})$ fades to colourless, changing the absorbance of the solution. Known concentrations of $\text{I}_2(\text{aq})$ are used to prepare a calibration curve graph and the absorbance is then measured as the reaction proceeds.




What is the rate of reaction at 20 s?

- A $5 \times 10^{-6} \text{ mol dm}^{-3} \text{ s}^{-1}$
 B $1 \times 10^{-5} \text{ mol dm}^{-3} \text{ s}^{-1}$
 C $5 \times 10^{-3} \text{ mol dm}^{-3} \text{ s}^{-1}$
 D $1 \times 10^{-2} \text{ mol dm}^{-3} \text{ s}^{-1}$

15 The diagram shows a Boltzmann distribution curve.

The axes are not labelled.



Points X and Y are points on the vertical axis.

What is represented by both points X and Y?

	point X	point Y
A	number of molecules with energy equal to E_A	largest number of molecules with the same energy
B	number of molecules with energy equal to or greater than E_A	largest number of molecules with the same energy
C	number of molecules with energy equal to E_A	the amount of energy of the greatest number of molecules
D	number of molecules with energy equal to or greater than E_A	the amount of energy of the greatest number of molecules

16 What are the acid–base nature and structure of SO_2 ?

	acid–base nature	structure
A	acidic	giant covalent lattice
B	acidic	simple molecular
C	basic	giant covalent lattice
D	basic	simple molecular

- 17** Elements X and Y are in Period 3 of the Periodic Table. Element X is either phosphorus or sulfur. Element Y is either sodium or magnesium.

Element X forms an oxide that reacts with water to give a solution containing the aqueous anion XO_4^{2-} .

One mole of element Y reacts with one mole of chlorine molecules. At the end of the reaction, all of the element Y and all of the chlorine molecules have been used up.

What are elements X and Y?

	X	Y
A	phosphorus	sodium
B	phosphorus	magnesium
C	sulfur	sodium
D	sulfur	magnesium

- 18** Q is a semi-conductor. The chloride of Q reacts with water to form white fumes and an acidic solution.

Which Period 3 element is Q?

- A** magnesium
B aluminium
C silicon
D phosphorus

- 19** V and W are two compounds. Each one contains a different Group 2 element.

A sample of each solid is added to water, shaken, and the pH of the resulting solution is measured.

compound	V	W
pH	13.6	9.4

Which row could identify V and W?

	V	W
A	BaSO_4	MgSO_4
B	MgSO_4	BaSO_4
C	Ba(OH)_2	Mg(OH)_2
D	Mg(OH)_2	Ba(OH)_2

- 20 Compound L decomposes on heating. One of the products is gas M.

M reacts with unburned hydrocarbons to form peroxyacetyl nitrate, PAN.

What could be the formula of L?

- A** CaNO_3 **B** $\text{Ca}(\text{NO}_3)_2$ **C** MgCO_3 **D** $\text{Mg}(\text{CO}_3)_2$

- 21 In reaction 1, concentrated sulfuric acid is added to potassium chloride and the fumes produced are bubbled into aqueous potassium iodide solution.

In reaction 2, potassium chloride is dissolved in aqueous ammonia and this is then added to aqueous silver nitrate.

What are the observations for reactions 1 and 2?

	observation for reaction 1	observation for reaction 2
A	brown solution	colourless solution
B	brown solution	white precipitate
C	colourless solution	colourless solution
D	colourless solution	white precipitate

- 22 The table refers to the hydrogen halides.

Which row is correct?

	oxidation	thermal stability
A	easier to oxidise down the group	increases down the group
B	more difficult to oxidise down the group	increases down the group
C	easier to oxidise down the group	decreases down the group
D	more difficult to oxidise down the group	decreases down the group

- 23 7.5 g of nitrogen monoxide reacts with 7.0 g of carbon monoxide on the surface of the catalytic converter in the exhaust system of a car.

What is the total volume of the product gases measured at room conditions?

- A** 3.0 dm^3 **B** 6.0 dm^3 **C** 9.0 dm^3 **D** 12.0 dm^3


24 Three statements about ammonia molecules and ammonium ions are given.

- 1 In aqueous solution, ammonia molecules form coordinate bonds with hydroxide ions.
- 2 Ammonium ions are Brønsted–Lowry acids.
- 3 The H–N–H bond angle is larger in the ammonium ion than in the ammonia molecule.

Which statements are correct?

- A** 1 and 2 only **B** 1 and 3 only **C** 2 and 3 only **D** 1, 2 and 3

25 Ethene reacts with steam in the presence of sulfuric acid.




Which type of reaction is this?

- A** acid–base
B addition
C hydrolysis
D substitution

26 Compound Z has the molecular formula $\text{C}_4\text{H}_8\text{O}_2$.

Compound Z reacts with propan-1-ol in the presence of concentrated H_2SO_4 .



The diagram shows the skeletal formulae of three compounds, S, T and U.



What are the possible skeletal formulae of the products of the reaction between compound Z and propan-1-ol?

- A** S and T **B** U only **C** S and U **D** T only

27 Geraniol and nerol are isomers of each other.



Which type of isomerism is shown here?


- A chain
- B geometrical (cis / trans)
- C optical
- D positional

28 Which compound has the greatest number of stereoisomers?

- A 2-methylhex-2-ene
- B 3-methylhex-2-ene
- C 4-methylhex-2-ene
- D 5-methylhex-2-ene

29 Vitamin A contains retinol.

retinol



Under appropriate conditions, acidified $\text{KMnO}_4(\text{aq})$ can be used to break $\text{C}=\text{C}$ bonds.


After these bonds have been broken, further oxidation of the fragments may occur.

Under which conditions is the acidified $\text{KMnO}_4(\text{aq})$ used and what do the final oxidation products include?

	conditions	final oxidation products
A	cold, dilute	aldehydes and carboxylic acids
B	cold, dilute	ketones and carboxylic acids
C	hot, concentrated	aldehydes and carboxylic acids
D	hot, concentrated	ketones and carboxylic acids

30 The structure of limonene is shown.

limonene



What are the number of moles of carbon dioxide and water produced when a sample of limonene is completely combusted in oxygen?

	number of moles of carbon dioxide	number of moles of water
A	4	3
B	5	4
C	5	8
D	9	7


- 31 The reaction of chlorine with methane is carried out in the presence of light.

What is the function of the light?

- A to break the C–H bonds in methane
- B to break the chlorine molecules into atoms
- C to break the chlorine molecules into ions
- D to heat the mixture

- 32 When X is added to NaOH(aq) and heated under reflux, pentan-2-ol is made.

Which organic product is made when X is heated with a solution of KCN dissolved in ethanol?



- 33 1-chlorobutane and 1-iodobutane both react with aqueous sodium hydroxide by a nucleophilic substitution mechanism.

Which reaction has the greatest rate under the same conditions and which mechanism is followed by this reaction?


	greatest rate	mechanism
A	1-chlorobutane	S _N 1
B	1-chlorobutane	S _N 2
C	1-iodobutane	S _N 1
D	1-iodobutane	S _N 2

- 34 Compound Y reacts with alkaline I₂(aq). When the products of this reaction are acidified, a dicarboxylic acid is produced. The formula of the dicarboxylic acid is HOOC–R–COOH where R consists of one or more CH₂ groups.

Which compound is Y?

- A pentan-1,4-diol
- B pentan-1,5-diol
- C pentan-2,3-diol
- D pentan-2,4-diol

- 35 Which alcohol gives only **one** possible oxidation product when warmed with dilute acidified potassium dichromate(VI)?
- A butan-1-ol
 B butan-2-ol
 C 2-methylpropan-1-ol
 D 2-methylpropan-2-ol
- 36 Which compound, on reaction with hydrogen cyanide, produces a compound with a chiral centre?
- A CH_3CHO
 B $\text{CH}_3\text{CH}_2\text{COCH}_2\text{CH}_3$
 C $\text{CH}_3\text{CO}_2\text{CH}_3$
 D HCHO
- 37 The diagram shows three reactions of ethanal. In each case, an excess of ethanal is used.




Observations are made after each of the three reactions.

What are the colours of solution 1 and solids 2 and 3?

	solution 1	solid 2	solid 3
A	green	yellow	silver mirror
B	green	yellow	red
C	orange	red	silver mirror
D	orange	red	red


38 $(\text{CH}_3)_3\text{CCN}$ reacts to form alcohol Y via the reaction sequence shown.



Which row names the molecule X and the class of alcohol Y?

	name of molecule X	class of alcohol Y
A	2,2-dimethylbutanoic acid	primary
B	3,3-dimethylbutanoic acid	tertiary
C	dimethylpropanoic acid	primary
D	dimethylpropanoic acid	tertiary


39 The diagram shows a section of an addition polymer. The polymer is made using two different monomers.



What are the names of the two monomers needed to make this polymer?

- A** 1,2-dichloropropene and 2-chlorobut-2-ene
- B** 2,3-dichlorobut-2-ene and chloropropene
- C** 1,2-dichloropropene and chloroethene
- D** chloropropene and 2-chlorobut-2-ene

- 40 The diagram shows the mass spectrum of a sample of chlorine. Peaks V, W, X, Y and Z are labelled.



Which statements about this spectrum are correct?

- 1 The relative atomic mass of chlorine can be calculated from the abundances and m/e values of 2 of the 5 peaks.
- 2 37.0 g of the species responsible for peak Z contains 3.011×10^{23} molecules.
- 3 The relative molecular mass of chlorine can be calculated from the abundances and m/e values of peaks X, Y and Z.

A 1, 2 and 3 **B** 1 and 2 only **C** 1 and 3 only **D** 2 and 3 only

BLANK PAGE

BLANK PAGE

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 $\text{J g}^{-1} \text{ K}^{-1}$)

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

The Periodic Table of Elements

Group																									
1	2	Key										13	14	15	16	17	18								
		atomic number atomic symbol name relative atomic mass												1 H hydrogen 1.0											
3 Li lithium 6.9	4 Be beryllium 9.0																	5 B boron 10.8	6 C carbon 12.0	7 N nitrogen 14.0	8 O oxygen 16.0	9 F fluorine 19.0	10 Ne neon 20.2		
11 Na sodium 23.0	12 Mg magnesium 24.3																	13 Al aluminium 27.0	14 Si silicon 28.1	15 P phosphorus 31.0	16 S sulfur 32.1	17 Cl chlorine 35.5	18 Ar argon 39.9		
19 K potassium 39.1	20 Ca calcium 40.1	21 Sc scandium 45.0	22 Ti titanium 47.9	23 V vanadium 50.9	24 Cr chromium 52.0	25 Mn manganese 54.9	26 Fe iron 55.8	27 Co cobalt 58.9	28 Ni nickel 58.7	29 Cu copper 63.5	30 Zn zinc 65.4	31 Ga gallium 69.7	32 Ge germanium 72.6	33 As arsenic 74.9	34 Se selenium 79.0	35 Br bromine 79.9	36 Kr krypton 83.8								
37 Rb rubidium 85.5	38 Sr strontium 87.6	39 Y yttrium 88.9	40 Zr zirconium 91.2	41 Nb niobium 92.9	42 Mo molybdenum 95.9	43 Tc technetium —	44 Ru ruthenium 101.1	45 Rh rhodium 102.9	46 Pd palladium 106.4	47 Ag silver 107.9	48 Cd cadmium 112.4	49 In indium 114.8	50 Sn tin 118.7	51 Sb antimony 121.8	52 Te tellurium 127.6	53 I iodine 126.9	54 Xe xenon 131.3								
55 Cs caesium 132.9	56 Ba barium 137.3	57–71 lanthanoids		72 Hf hafnium 178.5	73 Ta tantalum 180.9	74 W tungsten 183.8	75 Re rhenium 186.2	76 Os osmium 190.2	77 Ir iridium 192.2	78 Pt platinum 195.1	79 Au gold 197.0	80 Hg mercury 200.6	81 Tl thallium 204.4	82 Pb lead 207.2	83 Bi bismuth 209.0	84 Po polonium —	85 At astatine —	86 Rn radon —							
87 Fr francium —	88 Ra radium —	89–103 actinoids		104 Rf rutherfordium —	105 Db dubnium —	106 Sg seaborgium —	107 Bh bohrium —	108 Hs hassium —	109 Mt meitnerium —	110 Ds darmstadtium —	111 Rg roentgenium —	112 Cn copernicium —	113 Nh nihonium —	114 Fl flerovium —	115 Mc moscovium —	116 Lv livermorium —	117 Ts tennessine —	118 Og oganesson —							