



Cambridge International AS & A Level

CANDIDATE
NAME



CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

CHEMISTRY

9701/21

Paper 2 AS Level Structured Questions

October/November 2024

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 60.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **20** pages. Any blank pages are indicated.



1 Cobalt, rhodium and iridium are metals in the same group of the Periodic Table.

(a) The shorthand electronic configuration of cobalt is $[\text{Ar}]3d^74s^2$.

(i) Identify what is meant by $[\text{Ar}]$ by giving its full electronic configuration.

..... [1]

(ii) The lowest-energy electrons in cobalt are in the 1s orbital.

Draw the shape of a 1s orbital.

[1]

(iii) Deduce the number of unpaired electrons in a cobalt atom.

..... [1]

(b) Table 1.1 gives some details of the stable naturally occurring isotopes of rhodium and iridium.

Table 1.1

isotope	number of protons	number of neutrons	total number of electron shells
$^{103}_{45}\text{Rh}$		58	
$^{191}_{77}\text{Ir}$			6
$^{193}_{77}\text{Ir}$			6

Complete Table 1.1.

[3]



- (c) Table 1.2 shows the relative abundances of isotopes in a sample of an alloy containing rhodium and iridium only.

Table 1.2

isotope	relative isotopic mass	relative abundance in alloy
$^{103}_{45}\text{Rh}$	102.91	50.00
$^{191}_{77}\text{Ir}$	190.96	15.18
$^{193}_{77}\text{Ir}$	192.96	34.82

- (i) Define relative isotopic mass.

.....

 [2]

- (ii) Use Table 1.2 to calculate the relative atomic mass, A_r , of iridium in the alloy.

Give your answer to **two** decimal places.

relative atomic mass of iridium = ... [2]





- (d) Hydrated rhodium(III) chloride, $\text{RhCl}_3 \cdot x\text{H}_2\text{O}$, catalyses the conversion of ethene to but-2-ene.

Both stereoisomers of but-2-ene are formed in the reaction.

- (i) Hydrated rhodium(III) chloride contains 20.5% by mass of water of crystallisation.

Deduce the integer value of x in $\text{RhCl}_3 \cdot x\text{H}_2\text{O}$.

Show your working.

$x = \dots\dots\dots$
[2]



- (ii) Define stereoisomers.

.....
..... [1]

- (iii) Explain how the conversion of ethene to but-2-ene can be described as an addition reaction.

.....
..... [1]

- (iv) Draw the two stereoisomers of but-2-ene.



[2]

[Total: 16]





BLANK PAGE

DO NOT WRITE IN THIS MARGIN





2 Chlorine is one of the elements in Group 17 of the Periodic Table.

(a) (i) Describe the colours of the Group 17 elements, chlorine to iodine, at room temperature.

.....
 [1]

(ii) Describe the relative reactivity of the elements chlorine to iodine as oxidising agents.

..... [1]

(iii) State what is observed when chlorine reacts with hydrogen.

..... [1]

(iv) Explain why the thermal stability of the hydrogen halides decreases down the group.

.....
 [1]

(b) The halogenoalkane $\text{CH}_3\text{CH}_2\text{Cl}$ forms when chlorine reacts with C_2H_6 via a free-radical substitution mechanism.

(i) Define free radical.

.....
 [1]

(ii) State the essential condition for chlorine to react with C_2H_6 at room temperature.

..... [1]

(iii) Write **two** equations to show the propagation steps in this reaction.

1
 2 [2]

(c) CHCl_3 is another halogenoalkane. CHCl_3 forms when propanone reacts with NaClO .

NaClO is made from chlorine in a disproportionation reaction.

(i) Identify a reagent and conditions that can be used to convert chlorine to NaClO .

..... [1]

(ii) Define disproportionation.

.....
 [1]





- (iii) Write numbers in the boxes to balance the equation showing the reaction of propanone with NaClO .



[1]

- (iv) Aqueous AgNO_3 dissolved in ethanol reacts with an aqueous solution of CHCl_3 .

State what is observed in this reaction. Explain your answer.

.....

 [2]

[Total: 13]





3 The Group 14 elements show a change from non-metallic to metallic character down the group.

- (a) Table 3.1 shows some properties of two Group 14 elements, C and Sn, in their standard states. The table is incomplete.

Table 3.1

	C (graphite)	Sn
state and appearance in standard state	grey shiny solid	silvery solid
electrical conductivity		good
type of bonding		metallic
type of structure	giant	

- (i) Complete Table 3.1. [3]

- (ii) Identify the lattice structure shown by graphite.

..... [1]

- (iii) Explain why Sn has good electrical conductivity.

..... [1]

- (b) Carbon is found in inorganic compounds such as carbonates.

- (i) Write an equation for the reaction of magnesium carbonate with dilute HCl(aq) .

..... [1]

- (ii) Describe the thermal stability of the carbonates down Group 2.

..... [1]

- (iii) Ammonium carbonate undergoes an acid–base reaction with NaOH(aq) .

Explain this statement.

..... [2]



(c) Fig. 3.1 shows a sketch of some of the ionisation energies of silicon, Si.




Fig. 3.1

(i) Complete the graph in Fig. 3.1 to show the third to sixth ionisation energies of Si.

[2]

(ii) Construct an equation to represent the second ionisation energy of Si.

..... [1]





(d) Fig. 3.2 shows the boiling points of the simplest hydrides of the Group 14 elements, C to Pb.




Fig. 3.2

(i) Explain the trend in the boiling points of the Group 14 hydrides shown in Fig. 3.2.

.....
.....
.....
..... [2]

(ii) Deduce the shape of a molecule of SiH₄.

..... [1]



(e) Silicon readily reacts with elements of high electronegativity.

(i) Write an equation for the formation of SiCl_4 from its constituent elements.

..... [1]

(ii) Describe what is observed when a small sample of SiCl_4 is added to water.

.....
..... [1]

(iii) SiO_2 is a white solid that melts above 1700°C .

SiCl_4 is a colourless liquid at room temperature.

Explain the difference in the melting points of these two compounds with reference to their structure and bonding.

.....
.....
.....
.....
..... [2]

(f) Tin forms an amphoteric oxide, SnO_2 .

Suggest the formula of the tin compound that forms when SnO_2 reacts with H_2SO_4 in an acid–base reaction.

..... [1]

[Total: 20]





- 4 Propanone, CH_3COCH_3 , is an important organic reagent. Fig. 4.1 shows some reactions of propanone and its derivatives.




Fig. 4.1

- (a) Reaction 1 is a nucleophilic addition reaction.

- (i) Complete Fig. 4.2 to show the mechanism for the formation of **A** from propanone.

Include charges, dipoles, lone pairs of electrons and curly arrows as appropriate.



Fig. 4.2

[3]

- (ii) Explain why **A** does **not** show optical isomerism.

.....
 [1]

- (b) Suggest the reagents and conditions for reaction 2.

..... [1]





(c) Reaction 3 is a reduction reaction.

(i) Construct an equation to represent reaction 3.

Use [H] to represent one atom of hydrogen from the reducing agent.

..... [1]

(ii) Name C.

..... [1]

(d) State what is observed in reaction 4.

..... [1]

(e) Explain why Fehling's reagent does **not** react with propanone.

.....
..... [1]





(f) Compounds **A**, **B** and **C** can be distinguished using infrared spectroscopy.






Fig. 4.3 shows the infrared spectrum of one of the compounds.




Fig. 4.3

Table 4.1

bond	functional groups containing the bond	characteristic infrared absorption range (in wavenumbers)/cm ⁻¹
C–O	hydroxy, ester	1040–1300
C=C	aromatic compound, alkene	1500–1680
C=O	amide carbonyl, carboxyl ester	1640–1690 1670–1740 1710–1750
C≡N	nitrile	2200–2250
C–H	alkane	2850–2950
N–H	amine, amide	3300–3500
O–H	carboxyl hydroxy	2500–3000 3200–3600





- (i) Explain why the absorptions at $2850\text{--}2950\text{ cm}^{-1}$ are **not** useful to help determine which of the compounds **A**, **B** or **C** produces the infrared spectrum in Fig. 4.3.

Use Table 4.1 to answer this question.

.....
 [1]

- (ii) Identify which of compounds **A**, **B** or **C** produces the infrared spectrum in Fig. 4.3. Explain your answer.

compound

explanation

.....
 [1]

[Total: 11]







DO NOT WRITE IN THIS MARGIN







Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 J g ⁻¹ K ⁻¹)





The Periodic Table of Elements

Group																		
1	2	Key										13	14	15	16	17	18	
		atomic number atomic symbol name relative atomic mass																
		1 H hydrogen 1.0																
3 Li lithium 6.9	4 Be beryllium 9.0											5 B boron 10.8	6 C carbon 12.0	7 N nitrogen 14.0	8 O oxygen 16.0	9 F fluorine 19.0	10 Ne neon 20.2	
11 Na sodium 23.0	12 Mg magnesium 24.3											13 Al aluminium 27.0	14 Si silicon 28.1	15 P phosphorus 31.0	16 S sulfur 32.1	17 Cl chlorine 35.5	18 Ar argon 39.9	
19 K potassium 39.1	20 Ca calcium 40.1	21 Sc scandium 45.0	22 Ti titanium 47.9	23 V vanadium 50.9	24 Cr chromium 52.0	25 Mn manganese 54.9	26 Fe iron 55.8	27 Co cobalt 58.9	28 Ni nickel 58.7	29 Cu copper 63.5	30 Zn zinc 65.4	31 Ga gallium 69.7	32 Ge germanium 72.6	33 As arsenic 74.9	34 Se selenium 79.0	35 Br bromine 79.9	36 Kr krypton 83.8	
37 Rb rubidium 85.5	38 Sr strontium 87.6	39 Y yttrium 88.9	40 Zr zirconium 91.2	41 Nb niobium 92.9	42 Mo molybdenum 95.9	43 Tc technetium —	44 Ru ruthenium 101.1	45 Rh rhodium 102.9	46 Pd palladium 106.4	47 Ag silver 107.9	48 Cd cadmium 112.4	49 In indium 114.8	50 Sn tin 118.7	51 Sb antimony 121.8	52 Te tellurium 127.6	53 I iodine 126.9	54 Xe xenon 131.3	
55 Cs caesium 132.9	56 Ba barium 137.3	lanthanoids										81 Tl thallium 204.4	82 Pb lead 207.2	83 Bi bismuth 209.0	84 Po polonium —	85 At astatine —	86 Rn radon —	
87 Fr francium —	88 Ra radium —	actinoids										113 Nh nihonium —	114 Fl flerovium —	115 Mc moscovium —	116 Lv livermorium —	117 Ts tennessine —	118 Og oganeson —	

lanthanoids

57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.2	61 Pm promethium —	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0
89 Ac actinium —	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium —	94 Pu plutonium —	95 Am americium —	96 Cm curium —	97 Bk berkelium —	98 Cf californium —	99 Es einsteinium —	100 Fm fermium —	101 Md mendelevium —	102 No nobelium —	103 Lr lawrencium —

actinoids

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.





Cambridge International AS & A Level

CANDIDATE
NAME



CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

CHEMISTRY

9701/22

Paper 2 AS Level Structured Questions

October/November 2024

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 60.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **16** pages.



1 Vanadium, niobium and tantalum are metals in the same group of the Periodic Table.

(a) The shorthand electronic configuration of vanadium in the ground state is $[\text{Ar}]3d^34s^2$.

(i) State what is meant by the term ground state.

..... [1]

(ii) Show the electronic configuration of vanadium using electrons in boxes notation.

$[\text{Ar}]$

[1]

(iii) Deduce the total number of electrons in the p sub-shells of a vanadium atom.

..... [1]

(b) Pelopium was the suggested name for a new element discovered in a mineral.

Pelopium was later found to be a mixture of niobium, Nb, and tantalum, Ta.

Only one naturally occurring isotope exists for each of Nb and Ta.

(i) Complete Table 1.1.

Table 1.1

isotope	relative isotopic mass	number of protons	number of neutrons
${}^{93}_{41}\text{Nb}$	92.91		
${}^{181}_{73}\text{Ta}$	180.95		

[2]

(ii) Define relative isotopic mass.

.....

 [2]





- (iii) A sample of pelopium contains 90.9% by mass ${}^{93}_{41}\text{Nb}$ and 9.1% by mass ${}^{181}_{73}\text{Ta}$.

Calculate the theoretical relative atomic mass of pelopium based on these data and Table 1.1.

Give your answer to two decimal places.

Show your working.

theoretical relative atomic mass of pelopium = ... [2]

[Total: 9]





2 Oxygen is a Group 16 element.

(a) (i) Write equations for the following reactions.

- sodium and oxygen

.....

- sulfur and oxygen

.....

[2]

(ii) Draw a dot-and-cross diagram to show the species present in Al_2O_3 .

Draw outer electrons only.

[1]

(iii) The maximum oxidation state of the Period 3 elements in their oxides varies across the period.

State and explain the variation.

.....

.....

.....

[2]



(b) H_2O reacts with both inorganic and organic compounds.


(i) Complete Table 2.1 to give details of the reactions of some Period 3 oxides with H_2O .

Table 2.1




Period 3 oxide	product of reaction with H_2O	pH of solution formed
	$\text{Mg}(\text{OH})_2$	
P_4O_{10}		

[2]

(ii) Write an equation for the reaction of CH_3CN with H_2O in acidic conditions.



(iii) Draw the structures of the two alcohols formed in the reaction shown in equation 1.



[2]

(iv) Explain why alcohols are less acidic than water.

.....

 [2]





(c) Fig. 2.1 shows the boiling points of H_2O and other Group 16 hydrides.




Fig. 2.1

(i) Explain the trend in the boiling points of the Group 16 hydrides H_2S to H_2Te .

.....

.....

.....

..... [2]

(ii) Explain why the boiling point of H_2O is much higher than that of H_2S .

.....

.....

..... [1]

[Total: 15]





3 Nitrogen and phosphorus are elements in Group 15 of the Periodic Table.

(a) Nitrogen is found in inorganic compounds such as nitrogen oxides (NO_x), nitrates and nitric acid.

(i) Identify **one** natural and **one** man-made occurrence of nitrogen oxides in the atmosphere.

natural

man-made

[2]

(ii) Write an equation to describe the role of NO_2 in the direct formation of acid rain.

..... [1]

(iii) Peroxyacetyl nitrate, PAN, is a component of photochemical smog.

Describe how PAN forms from NO_2 .

.....

..... [1]

(iv) Nitric acid reacts with basic oxides to form nitrates.


Write an equation for the reaction of nitric acid with calcium oxide.

..... [1]

(v) Describe what is seen when solid calcium nitrate is heated strongly.

..... [1]

(b) A common test for nitrates is the reaction with NaOH and Al . Equation 1 shows the reaction.



(i) Deduce the oxidation state of nitrogen in NO_3^- .

..... [1]

(ii) Identify the species that is oxidised in equation 1.

..... [1]

(iii) NH_3 is a basic gas.

Describe how NH_3 is able to act as a base.

.....

..... [1]





(iv) Suggest the shape of the $[\text{Al}(\text{OH})_4]^-$ ion.

..... [1]

(c) Fig. 3.1 shows a sketch of some of the ionisation energies of phosphorus, P.




Fig. 3.1

(i) Construct an equation to represent the third ionisation energy of P.

..... [1]

(ii) Complete the graph in Fig. 3.1 to show the third to sixth ionisation energies of P.

[2]



- (d) Complete Table 3.1 to show the properties of nitrogen and phosphorus in their standard states.

Table 3.1

	nitrogen	phosphorus
state and appearance of standard state	colourless gas	white solid
electrical conductivity		poor
type of bonding		
type of structure	simple	

[2]

- (e) A form of solid nitrogen has a lattice structure similar to solid iodine.

Identify the type of lattice structure of solid nitrogen.

..... [1]

- (f) At very high temperatures, phosphorus can form P_2 molecules.

P_2 contains a triple bond, $P \equiv P$.

- (i) Describe the formation of the $P \equiv P$ bond in terms of orbital overlap.

.....

 [2]

- (ii) The bond energy of $P \equiv P$ is 485 kJ mol^{-1} . The bond energy of $N \equiv N$ is 944 kJ mol^{-1} . Compare the reactivity of P_2 and N_2 . Explain your answer.

.....

 [1]

[Total: 19]





- 4 Bromoalkanes are used widely in industry, although there is increasing concern about their environmental impact.

Fig. 4.1 shows a reaction scheme involving 1,2-dibromoethane.




Fig. 4.1

- (a) Complete Fig. 4.2 to show the mechanism for the formation of 1,2-dibromoethane in reaction 1.

Include charges, dipoles, lone pairs of electrons and curly arrows as appropriate.




Fig. 4.2

[3]



- (b) The enthalpy change of reaction 1, $\Delta H_r = -90.0 \text{ kJ mol}^{-1}$.



The enthalpy change of formation of ethene, $\Delta H_f = +52.2 \text{ kJ mol}^{-1}$.

Calculate the enthalpy change of formation of 1,2-dibromoethane.

ΔH_f of 1,2-dibromoethane = ... kJ mol^{-1} [1]

- (c) (i) Complete Fig. 4.1 to:

- draw the structure of compound **A**
- name compound **B**.

[2]

- (ii) Draw the structure of one repeat unit of polymer **C** in the box.

one repeat unit of polymer **C**

[1]

- (iii) In reaction 5, compound **B** reacts with an excess of NaOH dissolved in ethanol. The products are HBr, H_2O and an unsaturated hydrocarbon **D**.

Suggest the identity of **D**.



..... [1]





(d) Compound **E** is the only isomer of 1,2-dibromoethane.

Alkaline hydrolysis of **E** gives compound **F**.



(i) Identify the type of isomerism shown by **E** and 1,2-dibromoethane.

..... [1]

(ii) Name the homologous series that **F** belongs to.

..... [1]

(iii) Complete Table 4.1 to state what is observed when **F** reacts with the reagents listed.

Table 4.1

reagent	observation with F
2,4-dinitrophenylhydrazine (2,4-DNPH reagent)	
Tollens' reagent	
alkaline I ₂ (aq)	

[3]





Question 4 continues on page 14.





(e) Compound **F** reacts with reagent **G** to form compound **H**.



The infrared spectrum of **H** is shown in Fig. 4.3.




Fig. 4.3


Table 4.2

bond	functional groups containing the bond	characteristic infrared absorption range (in wavenumbers)/cm ⁻¹
C–O	hydroxy, ester	1040–1300
C=C	aromatic compound, alkene	1500–1680
C=O	amide carbonyl, carboxyl ester	1640–1690 1670–1740 1710–1750
C≡N	nitrile	2200–2250
C–H	alkane	2850–2950
N–H	amine, amide	3300–3500
O–H	carboxyl hydroxy	2500–3000 3200–3600



H also shows a molecular ion peak at $m/e = 60$ in its mass spectrum.

- (i) Use the information in (e), Fig. 4.3 and Table 4.2 to deduce the structure of **H**. Explain your answer fully.



[3]

- (ii) Suggest the role of reagent **G**.

[1]

[Total: 17]

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 J g ⁻¹ K ⁻¹)





The Periodic Table of Elements

Group																			
1	2													13	14	15	16	17	18
		1 H hydrogen 1.0																	
		Key atomic number atomic symbol name relative atomic mass																	
3	4																		
Li	Be																		
lithium	beryllium																		
6.9	9.0																		
11	12																		
Na	Mg																		
sodium	magnesium																		
23.0	24.3																		
19	20																		
K	Ca																		
potassium	calcium																		
39.1	40.1																		
37	38																		
Rb	Sr																		
rubidium	strontium																		
85.5	87.6																		
55	56																		
Cs	Ba																		
caesium	barium																		
132.9	137.3																		
87	88																		
Fr	Ra																		
francium	radium																		
—	—																		
		3	4	5	6	7	8	9	10	11	12			13	14	15	16	17	18
		5	6	7	8	9	10	11	12			13	14	15	16	17	18		
		B	C	N	O	F			Ne										
		boron	carbon	nitrogen	oxygen	fluorine			neon										
		10.8	12.0	14.0	16.0	19.0			20.2										
		13	14	15	16	17			18										
		Al	Si	P	S	Cl			Ar										
		aluminium	silicon	phosphorus	sulfur	chlorine			argon										
		27.0	28.1	31.0	32.1	35.5			39.9										
		31	32	33	34	35			36										
		Ga	Ge	As	Se	Br			Kr										
		gallium	germanium	arsenic	selenium	bromine			krypton										
		69.7	72.6	74.9	79.0	79.9			83.8										
		49	50	51	52	53			54										
		In	Sn	Sb	Te	I			Xe										
		indium	tin	antimony	tellurium	iodine			xenon										
		114.8	118.7	121.8	127.6	126.9			131.3										
		81	82	83	84	85			86										
		Tl	Pb	Bi	Po	At			Rn										
		thallium	lead	bismuth	polonium	astatine			radon										
		204.4	207.2	209.0	—	—			—										
		113	114	115	116	117			118										
		Nh	Fl	Mc	Lv	Ts			Og										
		nihonium	flerovium	moscovium	livermorium	tennessine			oganesson										
		—	—	—	—	—			—										

lanthanoids

57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.2	61 Pm promethium —	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0
89 Ac actinium —	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium —	94 Pu plutonium —	95 Am americium —	96 Cm curium —	97 Bk berkelium —	98 Cf californium —	99 Es einsteinium —	100 Fm fermium —	101 Md mendelevium —	102 No nobelium —	103 Lr lawrencium —

actinoids

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.





Cambridge International AS & A Level

CANDIDATE
NAME



CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

CHEMISTRY

9701/23

Paper 2 AS Level Structured Questions

October/November 2024

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 60.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **20** pages. Any blank pages are indicated.



1 Cobalt, rhodium and iridium are metals in the same group of the Periodic Table.

(a) The shorthand electronic configuration of cobalt is $[\text{Ar}]3d^74s^2$.

(i) Identify what is meant by $[\text{Ar}]$ by giving its full electronic configuration.

..... [1]

(ii) The lowest-energy electrons in cobalt are in the 1s orbital.

Draw the shape of a 1s orbital.

[1]

(iii) Deduce the number of unpaired electrons in a cobalt atom.

..... [1]

(b) Table 1.1 gives some details of the stable naturally occurring isotopes of rhodium and iridium.

Table 1.1

isotope	number of protons	number of neutrons	total number of electron shells
$^{103}_{45}\text{Rh}$		58	
$^{191}_{77}\text{Ir}$			6
$^{193}_{77}\text{Ir}$			6

Complete Table 1.1.

[3]



- (c) Table 1.2 shows the relative abundances of isotopes in a sample of an alloy containing rhodium and iridium only.

Table 1.2

isotope	relative isotopic mass	relative abundance in alloy
$^{103}_{45}\text{Rh}$	102.91	50.00
$^{191}_{77}\text{Ir}$	190.96	15.18
$^{193}_{77}\text{Ir}$	192.96	34.82

- (i) Define relative isotopic mass.

.....

 [2]

- (ii) Use Table 1.2 to calculate the relative atomic mass, A_r , of iridium in the alloy.

Give your answer to **two** decimal places.

relative atomic mass of iridium = ... [2]





- (d) Hydrated rhodium(III) chloride, $\text{RhCl}_3 \cdot x\text{H}_2\text{O}$, catalyses the conversion of ethene to but-2-ene.

Both stereoisomers of but-2-ene are formed in the reaction.

- (i) Hydrated rhodium(III) chloride contains 20.5% by mass of water of crystallisation.

Deduce the integer value of x in $\text{RhCl}_3 \cdot x\text{H}_2\text{O}$.

Show your working.

$x = \dots\dots\dots$
[2]



- (ii) Define stereoisomers.

.....
..... [1]

- (iii) Explain how the conversion of ethene to but-2-ene can be described as an addition reaction.

.....
..... [1]

- (iv) Draw the two stereoisomers of but-2-ene.



[2]

[Total: 16]





BLANK PAGE

DO NOT WRITE IN THIS MARGIN





2 Chlorine is one of the elements in Group 17 of the Periodic Table.

(a) (i) Describe the colours of the Group 17 elements, chlorine to iodine, at room temperature.

.....
 [1]

(ii) Describe the relative reactivity of the elements chlorine to iodine as oxidising agents.

..... [1]

(iii) State what is observed when chlorine reacts with hydrogen.

..... [1]

(iv) Explain why the thermal stability of the hydrogen halides decreases down the group.

.....
 [1]

(b) The halogenoalkane $\text{CH}_3\text{CH}_2\text{Cl}$ forms when chlorine reacts with C_2H_6 via a free-radical substitution mechanism.

(i) Define free radical.

.....
 [1]

(ii) State the essential condition for chlorine to react with C_2H_6 at room temperature.

..... [1]

(iii) Write **two** equations to show the propagation steps in this reaction.

1
 2 [2]

(c) CHCl_3 is another halogenoalkane. CHCl_3 forms when propanone reacts with NaClO .

NaClO is made from chlorine in a disproportionation reaction.

(i) Identify a reagent and conditions that can be used to convert chlorine to NaClO .

..... [1]

(ii) Define disproportionation.

.....
 [1]





- (iii) Write numbers in the boxes to balance the equation showing the reaction of propanone with NaClO .



[1]

- (iv) Aqueous AgNO_3 dissolved in ethanol reacts with an aqueous solution of CHCl_3 .

State what is observed in this reaction. Explain your answer.

.....

 [2]

[Total: 13]





3 The Group 14 elements show a change from non-metallic to metallic character down the group.

- (a) Table 3.1 shows some properties of two Group 14 elements, C and Sn, in their standard states. The table is incomplete.

Table 3.1

	C (graphite)	Sn
state and appearance in standard state	grey shiny solid	silvery solid
electrical conductivity		good
type of bonding		metallic
type of structure	giant	

- (i) Complete Table 3.1. [3]

- (ii) Identify the lattice structure shown by graphite.

..... [1]

- (iii) Explain why Sn has good electrical conductivity.

.....
 [1]

- (b) Carbon is found in inorganic compounds such as carbonates.

- (i) Write an equation for the reaction of magnesium carbonate with dilute HCl(aq).

..... [1]

- (ii) Describe the thermal stability of the carbonates down Group 2.

..... [1]

- (iii) Ammonium carbonate undergoes an acid–base reaction with NaOH(aq).

Explain this statement.

.....

 [2]



(c) Fig. 3.1 shows a sketch of some of the ionisation energies of silicon, Si.




Fig. 3.1

(i) Complete the graph in Fig. 3.1 to show the third to sixth ionisation energies of Si.

[2]

(ii) Construct an equation to represent the second ionisation energy of Si.

..... [1]





(d) Fig. 3.2 shows the boiling points of the simplest hydrides of the Group 14 elements, C to Pb.




Fig. 3.2

(i) Explain the trend in the boiling points of the Group 14 hydrides shown in Fig. 3.2.

.....

 [2]

(ii) Deduce the shape of a molecule of SiH₄.

..... [1]



(e) Silicon readily reacts with elements of high electronegativity.

(i) Write an equation for the formation of SiCl_4 from its constituent elements.

..... [1]

(ii) Describe what is observed when a small sample of SiCl_4 is added to water.

.....
..... [1]

(iii) SiO_2 is a white solid that melts above 1700°C .

SiCl_4 is a colourless liquid at room temperature.

Explain the difference in the melting points of these two compounds with reference to their structure and bonding.

.....
.....
.....
.....
..... [2]

(f) Tin forms an amphoteric oxide, SnO_2 .

Suggest the formula of the tin compound that forms when SnO_2 reacts with H_2SO_4 in an acid–base reaction.

..... [1]

[Total: 20]





- 4 Propanone, CH_3COCH_3 , is an important organic reagent. Fig. 4.1 shows some reactions of propanone and its derivatives.




Fig. 4.1

- (a) Reaction 1 is a nucleophilic addition reaction.

- (i) Complete Fig. 4.2 to show the mechanism for the formation of **A** from propanone.

Include charges, dipoles, lone pairs of electrons and curly arrows as appropriate.



Fig. 4.2

[3]

- (ii) Explain why **A** does **not** show optical isomerism.

.....
 [1]

- (b) Suggest the reagents and conditions for reaction 2.

..... [1]





(c) Reaction 3 is a reduction reaction.

(i) Construct an equation to represent reaction 3.

Use [H] to represent one atom of hydrogen from the reducing agent.

..... [1]

(ii) Name C.

..... [1]

(d) State what is observed in reaction 4.

..... [1]

(e) Explain why Fehling's reagent does **not** react with propanone.

.....
..... [1]





(f) Compounds **A**, **B** and **C** can be distinguished using infrared spectroscopy.






Fig. 4.3 shows the infrared spectrum of one of the compounds.




Fig. 4.3

Table 4.1

bond	functional groups containing the bond	characteristic infrared absorption range (in wavenumbers)/cm ⁻¹
C–O	hydroxy, ester	1040–1300
C=C	aromatic compound, alkene	1500–1680
C=O	amide carbonyl, carboxyl ester	1640–1690 1670–1740 1710–1750
C≡N	nitrile	2200–2250
C–H	alkane	2850–2950
N–H	amine, amide	3300–3500
O–H	carboxyl hydroxy	2500–3000 3200–3600





- (i) Explain why the absorptions at $2850\text{--}2950\text{ cm}^{-1}$ are **not** useful to help determine which of the compounds **A**, **B** or **C** produces the infrared spectrum in Fig. 4.3.

Use Table 4.1 to answer this question.

.....
 [1]

- (ii) Identify which of compounds **A**, **B** or **C** produces the infrared spectrum in Fig. 4.3. Explain your answer.

compound

explanation

.....
 [1]

[Total: 11]







BLANK PAGE

DO NOT WRITE IN THIS MARGIN





DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN





Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 J g ⁻¹ K ⁻¹)





The Periodic Table of Elements

Group																					
1	2											13	14	15	16	17	18				
Key atomic number atomic symbol name relative atomic mass 1 H hydrogen 1.0 2 He helium 4.0																					
3 Li lithium 6.9	4 Be beryllium 9.0															5 B boron 10.8	6 C carbon 12.0	7 N nitrogen 14.0	8 O oxygen 16.0	9 F fluorine 19.0	10 Ne neon 20.2
11 Na sodium 23.0	12 Mg magnesium 24.3															13 Al aluminium 27.0	14 Si silicon 28.1	15 P phosphorus 31.0	16 S sulfur 32.1	17 Cl chlorine 35.5	18 Ar argon 39.9
19 K potassium 39.1	20 Ca calcium 40.1	21 Sc scandium 45.0	22 Ti titanium 47.9	23 V vanadium 50.9	24 Cr chromium 52.0	25 Mn manganese 54.9	26 Fe iron 55.8	27 Co cobalt 58.9	28 Ni nickel 58.7	29 Cu copper 63.5	30 Zn zinc 65.4	31 Ga gallium 69.7	32 Ge germanium 72.6	33 As arsenic 74.9	34 Se selenium 79.0	35 Br bromine 79.9	36 Kr krypton 83.8				
37 Rb rubidium 85.5	38 Sr strontium 87.6	39 Y yttrium 88.9	40 Zr zirconium 91.2	41 Nb niobium 92.9	42 Mo molybdenum 95.9	43 Tc technetium —	44 Ru ruthenium 101.1	45 Rh rhodium 102.9	46 Pd palladium 106.4	47 Ag silver 107.9	48 Cd cadmium 112.4	49 In indium 114.8	50 Sn tin 118.7	51 Sb antimony 121.8	52 Te tellurium 127.6	53 I iodine 126.9	54 Xe xenon 131.3				
55 Cs caesium 132.9	56 Ba barium 137.3	57–71 lanthanoids		72 Hf hafnium 178.5	73 Ta tantalum 180.9	74 W tungsten 183.8	75 Re rhenium 186.2	76 Os osmium 190.2	77 Ir iridium 192.2	78 Pt platinum 195.1	79 Au gold 197.0	80 Hg mercury 200.6	81 Tl thallium 204.4	82 Pb lead 207.2	83 Bi bismuth 209.0	84 Po polonium —	85 At astatine —	86 Rn radon —			
87 Fr francium —	88 Ra radium —	89–103 actinoids		104 Rf rutherfordium —	105 Db dubnium —	106 Sg seaborgium —	107 Bh bohrium —	108 Hs hassium —	109 Mt meitnerium —	110 Ds darmstadtium —	111 Rg roentgenium —	112 Cn copernicium —	113 Nh nihonium —	114 Fl flerovium —	115 Mc moscovium —	116 Lv livermorium —	117 Ts tennessine —	118 Og oganesson —			

lanthanoids

actinoids

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

