

Cambridge International AS & A Level

MATHEMATICS**9709/11**

Paper 1 Pure Mathematics 1

October/November 2020

MARK SCHEME

Maximum Mark: 75

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2020 series for most Cambridge IGCSE™, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

This document consists of **16** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1: Marks must be awarded in line with: - the specific content of the mark scheme or the generic level descriptors for the question - the specific skills defined in the mark scheme or in the generic level descriptors for the question - the standard of response required by a candidate as exemplified by the standardisation scripts.
GENERIC MARKING PRINCIPLE 2: Marks awarded are always whole marks (not half marks, or other fractions).
GENERIC MARKING PRINCIPLE 3: Marks must be awarded positively: - marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate - marks are awarded when candidates clearly demonstrate what they know and can do - marks are not deducted for errors - marks are not deducted for omissions - answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.
GENERIC MARKING PRINCIPLE 4: Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.
GENERIC MARKING PRINCIPLE 5: Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).
GENERIC MARKING PRINCIPLE 6: Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

PUBLISHED

Mathematics Specific Marking Principles	
1	Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
2	Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
3	Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
4	Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
5	Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread.
6	Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

PUBLISHED**Mark Scheme Notes**

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B** Mark for a correct result or statement independent of method marks.
- DM or DB** When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- FT** Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.
- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
 - For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
 - The total number of marks available for each question is shown at the bottom of the Marks column.
 - Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
 - Square brackets [] around text or numbers show extra information not needed for the mark to be awarded.

Abbreviations

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no ‘follow through’ from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

Question	Answer	Marks	Guidance
1	$2x^2 + 5 = mx - 3 \rightarrow 2x^2 - mx + 8 (=0)$	B1	Form 3-term quadratic
	$m^2 - 64$	M1	Find $b^2 - 4ac$.
	$-8 < m < 8$	A1	Accept (-8, 8) and equality included
		3	

Question	Answer	Marks	Guidance
2	$(y =) \left[-(x-3)^{-1} \right] \left[+\frac{1}{2}x^2 \right] (+c)$	B1 B1	
	$7 = 1 + 2 + c$	M1	Substitute $x = 2, y = 7$ into an integrated expansion (c present). Expect $c = 4$
	$y = -(x-3)^{-1} + \frac{1}{2}x^2 + 4$	A1	OE
		4	

PUBLISHED

Question	Answer	Marks	Guidance
3	(Derivative =) $4\pi r^2 \rightarrow 400\pi$	B1	SOI Award this mark for $\frac{dr}{dt}$
	$\frac{50}{\text{their derivative}}$	M1	Can be in terms of r
	$\frac{1}{8\pi}$ or 0.0398	A1	AWRT
		3	

Question	Answer	Marks	Guidance
4	$(y =) [3] + [2] \left[\cos \frac{1}{2} \theta \right]$	B1 B1 B1	
		3	

Question	Answer	Marks	Guidance
5(a)	$6C2 \times [2(x^2)]^4 \times \left[\frac{a}{(x)} \right]^2$, $6C3 \times [2(x^2)]^3 \times \left[\frac{a}{(x)} \right]^3$	B1 B1	SOI Can be seen in an expansion
	$15 \times 2^4 \times a^2 = 20 \times 2^3 \times a^3$	M1	SOI Terms must be from a correct series
	$a = \frac{15 \times 2^4}{20 \times 2^3} = \frac{3}{2}$	A1	OE
		4	

PUBLISHED

Question	Answer	Marks	Guidance
5(b)	0	B1	
		1	

Question	Answer	Marks	Guidance
6	$\frac{dy}{dx} = \left[\frac{1}{2} (25 - x^2)^{-1/2} \right] \times [-2x]$	B1 B1	
	$\frac{-x}{(25 - x^2)^{1/2}} = \frac{4}{3} \rightarrow \frac{x^2}{25 - x^2} = \frac{16}{9}$	M1	Set = $\frac{4}{3}$ and square both sides
	$16(25 - x^2) = 9x^2 \rightarrow 25x^2 = 400 \rightarrow x = (\pm)4$	A1	
	When $x = -4, y = 5 \rightarrow (-4, 5)$	A1	
		5	

PUBLISHED

Question	Answer	Marks	Guidance
7(a)	$\left(\frac{\sin \theta}{1 - \sin \theta} - \frac{\sin \theta}{1 + \sin \theta} \right) \frac{\sin \theta(1 + \sin \theta) - \sin \theta(1 - \sin \theta)}{1 - \sin^2 \theta}$	*M1	Put over a single common denominator
	$\frac{2\sin^2 \theta}{\cos^2 \theta}$	DM1	Replace $1 - \sin^2 \theta$ by $\cos^2 \theta$ and simplify numerator
	$2\tan^2 \theta$	A1	AG
		3	
7(b)	$2 \tan^2 \theta = 8 \rightarrow \tan \theta = (\pm) 2$	B1	SOI
	$(\theta =) 63.4^\circ, 116.6^\circ$	B1 B1 FT	FT on 180 – 1st solution (with justification)
		3	

Question	Answer	Marks	Guidance
8(a)	$S = \frac{a}{1-r}, \quad 2S = \frac{a}{1-R}$	B1	SOI at least one correct
	$\frac{2a}{1-r} = \frac{a}{1-R}$	M1	SOI
	$2 - 2R = 1 - r \rightarrow r = 2R - 1$	A1	AG
		3	

Question	Answer	Marks	Guidance
8(b)	$ar^2 = aR \rightarrow (a)(2R-1)^2 = R(a)$	*M1	
	$4R^2 - 5R + 1 (=0) \rightarrow (4R-1)(R-1) (=0)$	DM1	Allow use of formula or completing square.
	$R = \frac{1}{4}$	A1	Allow $R = 1$ in addition
	$S = \frac{2a}{3}$	A1	
	Alternative method for question 8(b)		
	$ar^2 = aR \rightarrow (a)r^2 = \frac{1}{2}(r+1)(a)$	*M1	Eliminating 1 variable
	$2r^2 - r - 1 (=0) \rightarrow (2r+1)(r-1) (=0)$	DM1	Allow use of formula or completing square. Must solve a quadratic.
	$r = -\frac{1}{2}$	A1	Allow $r = 1$ in addition
	$S = \frac{2a}{3}$	A1	
		4	

Question	Answer	Marks	Guidance
9(a)	$m_{AB} = \frac{4-2}{-1-3} = -\frac{1}{2}$	B1	
	Equation of tangent is $y-2=2(x-3)$	B1 FT	(3, 2) with <i>their</i> gradient $-\frac{1}{m_{AB}}$
		2	
9(b)	$AB^2 = 4^2 + 2^2 = 20$ or $r^2 = 20$ or $r = \sqrt{20}$ or $AB = \sqrt{20}$	B1	
	Equation of circle centre B is $(x-3)^2 + (y-2)^2 = 20$	M1 A1	FT <i>their</i> 20 for M1
		3	
9(c)	$(x-3)^2 + (2x-6)^2 = \text{their } 20$	M1	Substitute <i>their</i> $y-2=2x-6$ into <i>their</i> circle, centre B
	$5x^2 - 30x + 25 = 0$ or $5(x-3)^2 = 20$	A1	
	$[(5)(x-5)(x-1) \text{ or } x-3 = \pm 2]$ $x = 5, 1$	A1	
		3	

Question	Answer	Marks	Guidance
10(a)	$\left(\sin \theta = \frac{r}{OC} \rightarrow\right) OC = \frac{r}{\sin \theta}$	M1 A1	
	$CD = r + \frac{r}{\sin \theta}$	A1	
		3	
10(b)	Radius of arc $AB = 4 + \frac{4}{\sin \frac{\pi}{6}} = 4 + 8 = 12$	B1	SOI
	(Arc $AB \Rightarrow$ $their\ 12 \times \frac{2\pi}{6}$ or $\left(\frac{1}{2} AB = \right) \left(their\ 12 \times \frac{\pi}{6} \right)$	M1	Expect 4π , must use <i>their</i> CD, not 4
	Perimeter = $24 + 4\pi$	A1	
		3	

Question	Answer	Marks	Guidance
10(c)	Area $FOC = \frac{1}{2} \times 4 \times \text{their } OC \times \sin \frac{\pi}{3}$	M1	
	$8\sqrt{3}$	A1	
	Area sector $FOE = \frac{1}{2} \times \frac{2\pi}{3} \times 4^2 = \frac{16\pi}{3}$	B1	
	Shaded area = $16\sqrt{3} - \frac{16\pi}{3}$	A1	
	Alternative method for question 10(c)		
	$FC = \sqrt{(\text{their } OC)^2 - 4^2}$	M1	$\sqrt{48}$ or $4\sqrt{3}$
	Area $FOC = \frac{1}{2} \times 4 \times 4\sqrt{3} = 8\sqrt{3}$	A1	
	Area of half sector $FOE = \frac{1}{2} \times \frac{\pi}{3} \times 4^2 = \frac{8\pi}{3}$	B1	
	Shaded area = $16\sqrt{3} - \frac{16\pi}{3}$	A1	
		4	

PUBLISHED

Question	Answer	Marks	Guidance
11(a)	$fg(x) = (2x+1)^2 + 3$	B1	OE
		1	
11(b)	$y = (2x+1)^2 + 3 \rightarrow 2x+1 = (\pm)\sqrt{y-3}$	M1	1st two operations. Allow one sign error or x/y interchanged
	$x = (\pm)\frac{1}{2}(\sqrt{y-3} - 1)$	M1	OE 2nd two operations. Allow one sign error or x/y interchanged
	$(fg^{-1}(x) =) \frac{1}{2}(\sqrt{x-3} - 1) \text{ for } (x) > 3$	A1 B1	Allow (3, ∞)
		4	
11(c)	$gf(x) = 2(x^2 + 3) + 1$	B1	SOI
	$(2x+1)^2 + 3 - 3 = 2(x^2 + 3) + 1 \rightarrow 2x^2 + 4x - 6 (= 0)$	*M1	Express as 3-term quadratic
	$(2)(x+3)(x-1) (= 0)$	DM1	Or quadratic formula or completing the square
	$x = 1$	A1	
		4	

Question	Answer	Marks	Guidance
12(a)	$4x^{\frac{1}{2}} - 2x = 3 - x \rightarrow x - 4x^{\frac{1}{2}} + 3 (=0)$	*M1	3-term quadratic. Can be expressed as e.g. $u^2 - 4u + 3 (=0)$
	$\left(x^{\frac{1}{2}} - 1\right)\left(x^{\frac{1}{2}} - 3\right) (=0)$ or $(u-1)(u-3)(=0)$	DM1	Or quadratic formula or completing square
	$x^{\frac{1}{2}} = 1, 3$	A1	SOI
	$x = 1, 9$	A1	
	Alternative method for question 12(a)		
	$\left(4x^{\frac{1}{2}}\right)^2 = (3+x)^2$	*M1	Isolate $x^{\frac{1}{2}}$
	$16x = 9 + 6x + x^2 \rightarrow x^2 - 10x + 9 (=0)$	A1	3-term quadratic
	$(x-1)(x-9) (=0)$	DM1	Or formula or completing square on a quadratic obtained by a correct method
	$x = 1, 9$	A1	
		4	
12(b)	$\frac{dy}{dx} = 2x^{1/2} - 2$	*B1	
	$\frac{dy}{dx}$ or $2x^{1/2} - 2 = 0$ when $x = 1$ hence B is a stationary point	DB1	
		2	

Question	Answer	Marks	Guidance
12(c)	Area of correct triangle = $\frac{1}{2} (9 - 3) \times 6$	M1	or $\int_3^9 (3 - x)(dx) = \left[3x - \frac{1}{2}x^2 \right] \rightarrow -18$
	$\int (4x^{\frac{1}{2}} - 2x)(dx) = \left[\frac{4x^{\frac{3}{2}}}{\frac{3}{2}} - x^2 \right]$	B1 B1	
	$(72 - 81) - \left(\frac{64}{3} - 16 \right)$	M1	Apply limits 4 $\rightarrow$ <i>their</i> 9 to an integrated expression
	$-14\frac{1}{3}$	A1	OE
	Shaded region = $18 - 14\frac{1}{3} = 3\frac{2}{3}$	A1	OE
		6	



Cambridge International AS & A Level

MATHEMATICS

9709/12

Paper 1 Pure Mathematics 1

October/November 2020

MARK SCHEME

Maximum Mark: 75

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2020 series for most Cambridge IGCSE™, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

This document consists of **19** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

PUBLISHED

Mathematics Specific Marking Principles	
1	Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
2	Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
3	Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
4	Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
5	Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread.
6	Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

PUBLISHED**Mark Scheme Notes**

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

DM or DB When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.

FT Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.

- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
- For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
- The total number of marks available for each question is shown at the bottom of the Marks column.
- Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
- Square brackets [] around text or numbers show extra information not needed for the mark to be awarded.

Abbreviations

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no ‘follow through’ from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

PUBLISHED

Question	Answer	Marks	Guidance
1	Coefficient of x^3 in $(1-2x)^5$ is -80	B1	Can be seen in an expansion but must be simplified correctly.
	Coefficient of x^2 in $(1-2x)^5$ is 40	B1	
	Coefficient of x^3 in $(1+kx)(1-2x)^5$ is $40k-80=20$	M1	Uses the relevant two terms to form an equation $=20$ and solves to find k . Condone x^3 appearing in some terms if recovered.
	$(k=)\frac{5}{2}$	A1	
		4	

PUBLISHED

Question	Answer	Marks	Guidance
2	$(-2p)^2 = (2p + 6) \times (p + 2)$ or $\frac{-2p}{2p+6} = \frac{p+2}{-2p}$	M1	OE. Using “ a, b, c then $b^2 = ac$ ” or $a = 2p+6$, $ar = -2p$ and $ar^2 = p+2$ to form a correct relationship in terms of p only
	$(2p^2 - 10p - 12 = 0) p = 6$	A1	
	$a = 18$ and $r = -\frac{2}{3}$	A1	
	$(s_{\infty}) = \text{their } a \div (1 - \text{their } r)$ $\left(= 18 \div \frac{5}{3} \right)$	M1	Correct formula used with their values for a and r , $ r < 1$ Both a & r from the same value of p .
	$(s_{\infty} =)10.8$	A1	OE. A0 if an extra solution given
			SC B2 for $s_{\infty} = \frac{2p+6}{1 - \frac{-2p}{2p+6}}$ or $\frac{2p+6}{1 - \frac{p+2}{-2p}}$ ignore any subsequent algebraic simplification.
		5	

PUBLISHED

Question	Answer	Marks	Guidance
3	$2x^2 + m(2x + 1) - 6x - 4 (= 0)$	*M1	y eliminated and all terms on one side with correct algebraic steps. Condone $\pm$ errors
	Using $b^2 - 4ac$ on $2x^2 + x(2m - 6) + m - 4 (= 0)$	DM1	Any use of discriminant with their a , b and c identified correctly.
	$4m^2 - 32m + 68$ or $2m^2 - 16m + 34$ or $m^2 - 8m + 17$	A1	
	$(2m - 8)^2 + k$ or $(m - 4)^2 + k$ or minimum point $(4, k)$ or finds $b^2 - 4ac$ $(= -4, -16, -64)$	DM1	OE. Any valid method attempted on their 3-term quadratic
	$(m - 4)^2 + 1$ oe + always $> 0 \rightarrow 2$ solutions for all values of m or Minimum point $(4, 1) + (fn)$ always $> 0 \rightarrow 2$ solutions for all values of m or $b^2 - 4ac < 0$ + no solutions $\rightarrow 2$ solutions for the original equation for all values of m	A1	Clear and correct reasoning and conclusion without wrong working.
		5	

PUBLISHED

Question	Answer	Marks	Guidance
4	S_x and S_{x+1}	M1	Using two values of n in the given formula
	$a = 5, d = 2$	A1 A1	
	$a + (n - 1) d > 200 \rightarrow 5 + 2(k - 1) > 200$	M1	Correct formula used with their a and d to form an equation or inequality with 200, condone use of n
	$(k =) 99$	A1	Condone ≥ 99
	Alternative method for question 4		
	$\frac{n}{2}(2a + (n - 1) d) \equiv n^2 + 4n \rightarrow \left(\frac{d}{2} = 1, a - \frac{1}{2}d = 4\right)$	M1	Equating two correct expressions of S_n and equating coefficients of n and n^2
	$d = 2, a = 5$	A1 A1	
	$a + (n - 1) d > 200 \rightarrow 5 + 2(k - 1) > 200$	M1	Correct formula used with their a and d to form an equation or inequality with 200, condone use of n
	$(k =) 99$	A1	Condone ≥ 99
	Alternative method for question 4		
	$sum_k - sum_{k-1} \rightarrow k^2 + 4k - (k - 1)^2 - 4(k - 1)$	M1 A1	Using given formula with consecutive expressions subtracted. Allow $k+1$ and k .
	$2k + 3 > 200$ or $= 200$	M1 A1	Simplifying to a linear equation or inequality
	$(k =) 99$	A1	Condone ≥ 99
		5	

PUBLISHED

Question	Answer	Marks	Guidance
5(a)	0	B1	
		1	
5(b)	$(f^{-1}(x)) = \frac{x+2}{4}, (g^{-1}(x)) = \frac{4-x}{x} \text{ or } \frac{4}{x} - 1$	B1 B1	OE. Sight of correct inverses.
	$x^2 + 6x - 16 (= 0)$	B1	Equating inverses and simplifying.
	$(x+8)$ and $(x-2)$	M1	Correct attempt at solution of <i>their</i> 3-term quadratic-factorising, completing the square or use of formula.
	$(x =) 2 \text{ or } -8$	A1	Do not accept answers obtained with no method shown.
		5	

PUBLISHED

Question	Answer	Marks	Guidance
6(a)	$\left(\frac{1}{\cos x} - \frac{\sin x}{\cos x}\right)\left(\frac{1}{\sin x} + 1\right)$	B1	Uses “ $\tan x = \sin x \div \cos x$ ” throughout
	$\left(\frac{1 - \sin x}{\cos x}\right)\left(\frac{1 + \sin x}{\sin x}\right)$ or $\left(\frac{1 - \sin^2 x}{\cos x \sin x}\right)$	M1	Correct algebra leading to two or four terms
	$\left(\frac{\cos^2 x}{\cos x \sin x}\right)$	A1	OE. A correct expression which can be cancelled directly to $\frac{\cos x}{\sin x}$ e.g. $\frac{\cos x(1 - \sin^2 x)}{\sin x(1 - \sin^2 x)}$
	$\left(\frac{\cos^2 x}{\cos x \sin x}\right) = \left(\frac{\cos x}{\sin x}\right) = \frac{1}{\tan x}$	A1	AG. Must show cancelling. If x is missing throughout their working withhold this mark.
		4	
6(b)	Uses (a) $\rightarrow \frac{1}{\tan x} = 2 \tan^2 x \quad \tan^3 x = \frac{1}{2}$	M1	Reducing to $\tan^3 x = k$.
	$(x =) 38.4^\circ$	A1	AWRT. Ignore extra answers outside the range 0 to 180° but A0 if within.
		2	

PUBLISHED

Question	Answer	Marks	Guidance
7(a)	$f'(4) \left(= \frac{5}{2} \right)$	*M1	Substituting 4 into $f'(x)$
	$\left(\frac{dy}{dt} = \frac{dy}{dx} \times \frac{dx}{dt} \right) \rightarrow \left(\frac{dy}{dt} \right) = \frac{5}{2} \times 0.12$	DM1	Multiplies <i>their</i> $f'(4)$ by 0.12
	$\left(\frac{dy}{dt} = \right) 0.3$	A1	OE
		3	
7(b)	$\frac{6x^{\frac{1}{2}}}{\frac{1}{2}} - \frac{4x^{-\frac{1}{2}}}{-\frac{1}{2}} (+c)$	B1 B1	B1 for each unsimplified integral.
	Uses (4, 7) leading to $c = (-21)$	M1	Uses (4, 7) to find a c value
	y or $f(x) = 12x^{\frac{1}{2}} + 8x^{-\frac{1}{2}} - 21$ or $12\sqrt{x} + \frac{8}{\sqrt{x}} - 21$	A1	Need to see y or $f(x) =$ somewhere in <i>their</i> solution and 12 and 8
		4	

PUBLISHED

Question	Answer	Marks	Guidance
8(a)	Use of correct formula for the area of triangle ABC	M1	Use of $180-2\theta$ scores M0. Condone $2\pi-2\theta$
	$\frac{1}{2}r^2 \sin(\pi-2\theta)$ or $\frac{1}{2}r^2 \sin 2\theta$ or $2 \times \frac{1}{2}r \times r \cos \theta \times \sin \theta$ or $2 \times \frac{1}{2}r \cos \theta \times r \sin \theta$	A1	OE
	[Shaded area = triangle – sector] = <i>their</i> triangle area – $\frac{1}{2}r^2\theta$	B1 FT	FT for <i>their</i> triangle area – $\frac{1}{2}r^2\theta$ (Condone use of 180 degrees for triangle area for B1)
		3	
8(b)	Arc $BD = r\theta = 6$ cm	B1	SOI
	$AC = 2r \cos \theta = (2 \times 10 \cos 0.6 = 20 \cos 0.6 = 16.506)$ or $\sqrt{(2r^2 - 2r^2 \cos(\pi - 2\theta))}$ or $\frac{r \times \sin(\pi - 2\theta)}{\sin \theta}$	*M1	Finding AC or $\frac{1}{2}AC (= 8.25)$
	$DC = 2r \cos \theta - r$ or $\sqrt{(2r^2 - 2r^2 \cos(\pi - 2\theta))} - r (= 6.506)$	DM1	Subtracting r from <i>their</i> AC or $r - r \cos \theta$ from <i>their</i> half AC (8.25-1.75)
	(Perimeter = $10 + 6 + 6.506 =$) 22.5	A1	AWRT
		4	

PUBLISHED

Question	Answer	Marks	Guidance
9(a)	$r = \sqrt{6^2 + 3^2}$ or $r^2 = 45$	B1	Sight of $r = 6.7$ implies B1
	$(x - 5)^2 + (y - 1)^2 = r^2$ or $x^2 - 10x + y^2 - 2y = r^2 - 26$	M1	Using centre given and <i>their</i> radius or r in correct formula
	$(x - 5)^2 + (y - 1)^2 = 45$ or $x^2 - 10x + y^2 - 2y = 19$	A1	Do not allow $(\sqrt{45})^2$ for r^2
		3	
9(b)	C has coordinates (11, 4)	B1	
	0.5	B1	OE, Gradient of AB , BC or AC .
	Grad of $CD = -2$	M1	Calculation of gradient needs to be shown for this M1.
	$(\frac{1}{2} \times -2 = -1)$ then states + perpendicular $\rightarrow$ hence shown or tangent	A1	Clear reasoning needed.
	Alternative method for question 9(b)		
	C has coordinates (11, 4)	B1	
	0.5	B1	OE, Gradient of AB , BC or AC .
	Gradient of the perpendicular is -2 $\rightarrow$ Equation of the perpendicular is $y - 4 = -2(x - 11)$	M1	Use of $m_1 m_2 = -1$ with <i>their</i> gradient of AB , BC or AC and correct method for the equation of the perpendicular. Could use $D(5, 16)$ instead of $C(11, 4)$.
	Checks $D(5, 16)$ or checks gradient of CD and then states D lies on the line or CD has gradient $-2 \rightarrow$ hence shown or tangent	A1	Clear check and reasoning needed. Checks that the other point lies on the line or checks gradient.

PUBLISHED

Question	Answer	Marks	Guidance
9(b)	Alternative method for question 9(b)		
	C has coordinates (11, 4) or Gradient of AB , BC or $AC = 0.5$	B1	Only one of AB , BC or AC needed.
	Equation of the perpendicular is $y - 4 = -2(x - 11)$	B1	Finding equation of CD .
	$(x - 5)^2 + (-2x + 26 - 1)^2 = 45 \rightarrow (x^2 - 22x + 121 = 0)$	M1	Solving simultaneously with the equation of the circle.
	$(x - 11)^2 = 0$ or $b^2 - 4ac = 0 \rightarrow$ repeated root $\rightarrow$ hence shown or tangent	A1	Must state repeated root.
	Alternative method for question 9(b)		
	C has coordinates (11, 4)	B1	
	Finding $CD = \sqrt{180}$ and $BD = \sqrt{225}$	B1	OE. Calculated from the co-ordinates of B , C & D without using r .
	Checking $(\text{their } BD)^2 - (\text{their } CD)^2$ is the same as $(\text{their } r)^2$	M1	
	$\therefore$ Pythagoras valid $\therefore$ perpendicular $\rightarrow$ hence shown or tangent	A1	Triangle ACD could be used instead.
	Alternative method for question 9(b)		
	C has coordinates (11, 4)	B1	
	Finding vectors $\overrightarrow{AC}$ and $\overrightarrow{CD}$ or $\overrightarrow{BC}$ and $\overrightarrow{CD}$ $(= \begin{pmatrix} 6 \\ 3 \end{pmatrix} \text{ and } \begin{pmatrix} -6 \\ 12 \end{pmatrix} \text{ or } \begin{pmatrix} 12 \\ 6 \end{pmatrix} \text{ and } \begin{pmatrix} -6 \\ 12 \end{pmatrix})$	B1	Must be correct pairing.
	Applying the scalar product to one of these pairs of vectors	M1	Accept <i>their</i> $\overrightarrow{AC}$ and $\overrightarrow{CD}$ or <i>their</i> $\overrightarrow{BC}$ and $\overrightarrow{CD}$
	Scalar product = 0 then states $\therefore$ perpendicular $\rightarrow$ hence shown or tangent	A1	
		4	

PUBLISHED


Question	Answer	Marks	Guidance
9(c)	$E(-1, 4)$	B1 B1	WWW B1 for each coordinate Note: Equation of DE which is $y = 2x + 6$ may be used to find E
		2	

Question	Answer	Marks	Guidance
10(a)	$\left(\frac{dy}{dx}\right) = [8] \times [(3-2x)^{-3}] + [-1]$ $\left(= \frac{8}{(3-2x)^3} - 1 \right)$	B2, 1, 0	B2 for all three elements correct, B1 for two elements correct, B0 for only one or no elements correct.
	$\frac{d^2y}{dx^2} = -3 \times 8 \times (3-2x)^{-4} \times (-2)$ $\left(= \frac{48}{(3-2x)^4} \right)$	B1 FT	FT providing <i>their</i> bracket is to a negative power
	$\int y dx = [(3-2x)^{-1}] [2 \div (-1 \times -2)] [-\frac{1}{2}x^2] (+c)$ $\left(= \frac{1}{3-2x} - \frac{1}{2}x^2 + c \right)$	B1 B1 B1	Simplification not needed, B1 for each correct element
		6	

PUBLISHED

Question	Answer	Marks	Guidance
10(b)	$\frac{dy}{dx} = 0 \rightarrow (3 - 2x)^3 = 8 \rightarrow 3 - 2x = k \rightarrow x =$	M1	Setting <i>their</i> 2-term differential to 0 and attempts to solve as far as $x =$
	$\frac{1}{2}$	A1	
	Alternative method for question 10(b)		
	$y = 0 \rightarrow \frac{2}{(3 - 2x)^2} - x = 0 \rightarrow (x - 2)(2x - 1)^2 = 0 \rightarrow x =$	M1	Setting y to 0 and attempts to solve a cubic as far as $x =$ (3 factors needed)
	$\frac{1}{2}$	A1	
		2	
10(c)	Area under curve = <i>their</i> $\left[\frac{1}{3 - 2 \times \left(\frac{1}{2}\right)} - \frac{\left(\frac{1}{2}\right)^2}{2} \right] - \left[\frac{1}{3 - 2 \times 0} - 0 \right]$	M1	Using <i>their</i> integral, <i>their</i> positive x limit from part (b) and 0 correctly.
	$\frac{1}{24}$	A1	
		2	

PUBLISHED

Question	Answer	Marks	Guidance
11(a)	5, -1	B1 B1	Sight of each value
		2	
11(b)		*B1	Needs to be a curve, not straight lines. One complete cycle starting and finishing at <i>their</i> largest value.
		DB1	One complete cycle starting and finishing at $y = 5$ and going down to $y = -1$ and starting to level off at least one end.
		2	
11(c)(i)	0 solution	B1	
		1	
11(c)(ii)	2 solutions	B1	
		1	
11(c)(iii)	1 solution	B1	
		1	

PUBLISHED

Question	Answer	Marks	Guidance
11(d)	Stretch by (scale factor) $\frac{1}{2}$, parallel to x -axis or in x direction (or horizontally)	B1	
	Translation of $\begin{pmatrix} 0 \\ 4 \end{pmatrix}$	B1	Accept translation/shift Accept translation 4 units in positive y -direction.
		2	
11(e)	Translation of $\begin{pmatrix} -\frac{\pi}{2} \\ 0 \end{pmatrix}$	B1	Accept translation/shift Accept translation $-\frac{\pi}{2}$ units in x -direction.
	Stretch by (scale factor) 2 parallel to y -axis (or vertically).	B1	
		2	

Cambridge International AS & A Level

MATHEMATICS**9709/13**

Paper 1 Pure Mathematics 1

October/November 2020

MARK SCHEME

Maximum Mark: 75

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2020 series for most Cambridge IGCSE™, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

This document consists of **14** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

PUBLISHED

Mathematics Specific Marking Principles	
1	Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
2	Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
3	Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
4	Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
5	Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread.
6	Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B** Mark for a correct result or statement independent of method marks.
- DM or DB** When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- FT** Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.
- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
 - For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
 - The total number of marks available for each question is shown at the bottom of the Marks column.
 - Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
 - Square brackets [] around text or numbers show extra information not needed for the mark to be awarded.

Abbreviations

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no ‘follow through’ from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

PUBLISHED

Question	Answer	Marks	Guidance
1(a)	$\left[(x+3)^2 \right] \quad [-4]$	B1 B1	
		2	
1(b)	[Translation or shift] $\begin{pmatrix} -3 \\ -4 \end{pmatrix}$	B1 B1 FT	Accept [translation/shift] $\begin{pmatrix} -their\ a \\ their\ b \end{pmatrix}$ OR translation –3 units in x -direction and (translation) –4 units in y -direction.
		2	

Question	Answer	Marks	Guidance
2(a)	$\frac{-2}{x+2}$	B1	Integrate $f(x)$. Accept $-2(x+2)^{-1}$. Can be unsimplified.
	$0 - \left(-\frac{2}{3} \right) = \frac{2}{3}$	M1 A1	Apply limit(s) to an integrated expansion. CAO for A1
		3	
2(b)	$-1 = -2 + c$	M1	Substitute $x = -1, y = -1$ into <i>their</i> integrated expression (c present)
	$y = \frac{-2}{x+2} + 1$	A1	Accept $y = -2(x+2)^{-1} + 1$. –2 must be resolved.
		2	

PUBLISHED

Question	Answer	Marks	Guidance
3	$3\tan^4\theta + \tan^2\theta - 2 (=0)$	M1	SOI 3-term quartic, condone sign errors for this mark only
	$(3\tan^2\theta - 2)(\tan^2\theta + 1) (=0)$	M1	Attempt to factorise or solve 3-term quadratic in $\tan^2\theta$.
	$\tan\theta = (\pm)\sqrt{\frac{2}{3}}$ or $(\pm)0.816$ or $(\pm)0.817$	A1	SOI Implied by final answer = 39.2° after 1st M1 scored
	$39.2^\circ, 140.8^\circ$	A1 A1 FT	FT for 2nd solution = $180^\circ - 1^\text{st}$ solution
		5	

Question	Answer	Marks	Guidance
4	$3x^2 - 4x + 4 = mx + m - 1 \rightarrow 3x^2 - (4+m)x + (5-m) (=0)$	M1	3-term quadratic
	$b^2 - 4ac = (4+m)^2 - 4 \times 3 \times (5-m)$	M1	Find $b^2 - 4ac$ for <i>their</i> quadratic
	$m^2 + 20m - 44$	A1	
	$(m+22)(m-2)$	A1	Or use of formula or completing square. This step must be seen
	$m > 2$, $m < -22$	A1	Allow $x > 2$, $x < -22$
		5	

PUBLISHED

Question	Answer	Marks	Guidance
5	$[7C1a^6b(x)], [7C2a^5b^2(x^2)], [7C4a^3b^4(x^4)]$	B2, 1, 0	SOI, can be seen in an expansion.
	$\frac{7C2a^5b^2(x^2)}{7C1a^6b(x)} = \frac{7C4a^3b^4(x^4)}{7C2a^5b^2(x^2)} \rightarrow \frac{21a^5b^2}{7a^6b} = \frac{35a^3b^4}{21a^5b^2}$	M1 A1	M1 for a correct relationship OE (Ft from <i>their</i> 3 terms). For A1 binomial coefficients must be correct & evaluated.
	$\frac{a}{b} = \frac{5}{9}$	A1	OE
		5	

Question	Answer	Marks	Guidance
6(a)	$y = \frac{2x}{3x-1} \rightarrow 3xy - y = 2x \rightarrow 3xy - 2x = y$ (or $-y = 2x - 3xy$)	*M1	For 1st two operations. Condone a sign error
	$x(3y-2) = y \rightarrow x = \frac{y}{3y-2}$ (or $x = \frac{-y}{2-3y}$)	DM1	For 2nd two operations. Condone a sign error
	$(f^{-1}(x)) = \frac{x}{3x-2}$	A1	Allow $(f^{-1}(x)) = \frac{-x}{2-3x}$
		3	
6(b)	$\left[\frac{2(3x-1)+2}{3(3x-1)} \right] = \left[\frac{6x}{3(3x-1)} = \frac{2x}{3x-1} \right]$	B1 B1	AG, WWW First B1 is for a correct single unsimplified fraction. An intermediate step needs to be shown. Equivalent methods accepted.
		2	

PUBLISHED

Question	Answer	Marks	Guidance
6(c)	$(f(x)) > \frac{2}{3}$	B1	Allow $(y) > \frac{2}{3}$. Do not allow $x > \frac{2}{3}$
		1	

Question	Answer	Marks	Guidance
7(a)	$(d =) -\frac{\tan^2 \theta}{\cos^2 \theta} - \frac{1}{\cos^2 \theta}$	B1	Allow sign error(s). Award only at form $(d =) \dots$ stage
	$-\frac{\sin^2 \theta}{\cos^4 \theta} - \frac{1}{\cos^2 \theta}$ or $-\frac{\sec^2 \theta}{\cos^2 \theta}$	M1	Allow sign error(s). Can imply B1
	$\frac{-\sin^2 \theta - \cos^2 \theta}{\cos^4 \theta}$ or $-\frac{1}{\cos^2 \theta}$	M1	
	$-\frac{1}{\cos^4 \theta}$	A1	AG, WWW
		4	
7(b)	$a = \frac{4}{3}, d = -\frac{16}{9}$	B1	SOI, both required. Allow $a = \frac{1}{3}, d = -\frac{1}{9}$ $\frac{4}{4} \quad \frac{16}{16}$
	$u_{13} = \frac{1}{\cos^2 \theta} - \frac{12}{\cos^4 \theta} = \frac{4}{3} + 12\left(\frac{-16}{9}\right)$	M1	Use of correct formula with <i>their a</i> and <i>their d</i> . The first 2 steps could be reversed
	-20	A1	WWW
		3	

PUBLISHED

Question	Answer	Marks	Guidance
8(a)	$\frac{dy}{dx} = [2] \quad [-2(2x+1)^{-2}]$	B1 B1	
	$\frac{d^2y}{dx^2} = 8(2x+1)^{-3}$	B1	
		3	
8(b)	Set <i>their</i> $\frac{dy}{dx} = 0$ and attempt solution	M1	
	$(2x+1)^2 = 1 \rightarrow 2x+1 = (\pm) 1$ or $4x^2 + 4x = 0 \rightarrow (4)x(x+1) = 0$	M1	Solving as far as $x = \dots$
	$x = 0$	A1	WWW. Ignore other solution.
	$(0, 2)$	A1	One solution only. Accept $x = 0, y = 2$ only.
	$\frac{d^2y}{dx^2} > 0$ from a solution $x > -\frac{1}{2}$ hence minimum	B1	Ignore other solution. Condone arithmetic slip in value of $\frac{d^2y}{dx^2}$. <i>Their</i> $\frac{d^2y}{dx^2}$ must be of the form $k(2x+1)^{-3}$
		5	

Question	Answer	Marks	Guidance
9(a)	$\cos BAO = \frac{6}{8}$ or $\frac{8^2 + 12^2 - 8^2}{2 \times 8 \times 12}$	M1	Or other correct method
	$BAO = 0.723$	A1	
		2	

Question	Answer	Marks	Guidance
9(b)	Sector $ABC = \frac{1}{2} \times 12^2 \times \text{their } 0.7227$	*M1	Accept 52.1
	Triangle $AOB = \frac{1}{2} \times 8 \times 12 \sin(\text{their } 0.7227)$ or $\frac{1}{2} \times 12 \times \sqrt{28}$	*M1	or $\frac{1}{2} \times 8 \times 8 \sin(\pi - 2 \times \text{their } 0.7227)$. Expect 31.7 or 31.8
	Shaded area = $\text{their } 52.0 - \text{their } 31.7 = 20.3$	DM1 A1	M1 dependent on both previous M marks
		4	
9(c)	Arc $BC = 12 \times \text{their } 0.7227$	*M1	Expect 8.67
	Perimeter = $8 + 4 + \text{their } 8.67 = 20.7$	DM1 A1	
		3	

Question	Answer	Marks	Guidance
10(a)	$\frac{dy}{dx} = \left[\frac{x^{-1/2}}{2k} \right] - \left[\frac{x^{-3/2}}{2} \right] + ([0])$	B2, 1, 0	$([0])$ implies that more than 2 terms counts as an error
	Sub $\frac{dy}{dx} = 3$ when $x = \frac{1}{4}$ Expect $3 = \frac{1}{k} - 4$	M1	
	$k = \frac{1}{7}$ (or 0.143)	A1	
		4	

PUBLISHED

Question	Answer	Marks	Guidance
10(b)	$\int \frac{1}{k} x^{1/2} + x^{-1/2} + \frac{1}{k^2} = \left[\frac{2x^{3/2}}{3k} \right] + \left[2x^{1/2} \right] + \left[\frac{x}{k^2} \right]$	B2, 1, 0	OE
	$\left(\frac{2k^2}{3} + 2k + 1 \right) - \left(\frac{k^2}{12} + k + \frac{1}{4} \right)$	M1	Apply limits $\frac{k^2}{4} \rightarrow k^2$ to an integrated expression. Expect $\frac{7}{12}k^2 + k + \frac{3}{4}$
	$\frac{7}{12}k^2 + k + \frac{3}{4} = \frac{13}{12}$	M1	Equate to $\frac{13}{12}$ and simplify to quadratic. OE, expect $7k^2 + 12k - 4 (= 0)$
	$k = \frac{2}{7}$ only (or 0.286)	A1	Dependent on $(7k - 2)(k + 2) (= 0)$ or formula or completing square.
		5	

PUBLISHED

Question	Answer	Marks	Guidance
11(a)	$(-6-8)^2 + (6-4)^2$	M1	OE
	$= 200$	A1	
	$\sqrt{200} > 10$, hence outside circle	A1	AG ('Shown' not sufficient). Accept equivalents of $\sqrt{200} > 10$
	Alternative method for question 11(a)		
	Radius = 10 and $C = (8, 4)$	B1	
	Min(x) on circle = $8 - 10 = -2$	M1	
	Hence outside circle	A1	AG
		3	
11(b)	angle = $\sin^{-1}\left(\frac{\text{their } 10}{\text{their } 10\sqrt{2}}\right)$	M1	Allow decimals for $10\sqrt{2}$ at this stage. If cosine used, angle ACT or BCT must be identified, or implied by use of $90^\circ - 45^\circ$.
	angle = $\sin^{-1}\left(\frac{1}{\sqrt{2}}\right)$ or $\frac{\sqrt{2}}{2}$ or $\frac{10}{10\sqrt{2}}$ or $\frac{10}{\sqrt{200}}$) = 45°	A1	AG Do not allow decimals
	Alternative method for question 11(b)		
	$(10\sqrt{2})^2 = 10^2 + TA^2$	M1	
	$TA = 10 \rightarrow 45^\circ$	A1	AG
		2	

PUBLISHED

Question	Answer	Marks	Guidance
11(c)	Gradient, m , of $CT = -\frac{1}{7}$	B1	OE
	Attempt to find mid-point (M) of CT	*M1	Expect (1, 5)
	Equation of AB is $y - 5 = 7(x - 1)$	DM1	Through <i>their</i> (1, 5) with gradient $-\frac{1}{m}$
	$y = 7x - 2$	A1	
		4	
11(d)	$(x - 8)^2 + (7x - 2 - 4)^2 = 100$ or equivalent in terms of y	M1	Substitute <i>their</i> equation of AB into equation of circle.
	$50x^2 - 100x (= 0)$	A1	
	$x = 0$ and 2	A1	WWW
	Alternative method for question 11(d)		
	$\mathbf{MC} = \begin{pmatrix} 7 \\ -1 \end{pmatrix}$	M1	
	$\begin{pmatrix} 1 \\ 5 \end{pmatrix} + \begin{pmatrix} -1 \\ -7 \end{pmatrix} = \begin{pmatrix} 0 \\ -2 \end{pmatrix}, \quad \begin{pmatrix} 1 \\ 5 \end{pmatrix} + \begin{pmatrix} 1 \\ 7 \end{pmatrix} = \begin{pmatrix} 2 \\ 12 \end{pmatrix}$	A1	
	$x = 0$ and 2	A1	
		3	