

BIOLOGY



EXAM QUESTION PACK

TOPIC 2: CELLS, VIRUSES AND REPRODUCTION OF LIVING THINGS
SEXUAL REPRODUCTION IN MAMMALS

SCORE

/28

PERCENTAGE

%

STEMUP

These questions belong to Pearson Edexcel, not to us.

The examination questions and mark schemes in this pack were written and published by Pearson Edexcel. They are past assessment materials, reproduced here so that students can revise from them.

What we do not own

STEMUP is not affiliated with Pearson Edexcel and does not own, and makes no claim to own, any of the question or mark scheme material in this pack. Pearson has not endorsed, approved or been involved in the production of it. Pearson, Edexcel and the qualification titles used here belong to their respective owners and appear for identification only.

Where to find the originals


Pearson publishes its past papers and mark schemes free of charge. Always check the current specification at [qualifications.pearson.com](https://www.pearson.com/qualifications) before you revise, because content and wording can change between series.

If you hold the rights to any material in this pack and would like it amended or removed, contact admin@stemup.co.uk and we will act on it.

Q1.

A sea urchin is an invertebrate animal that lives on the seabed.

The photograph shows a sea urchin.



Sea urchin embryos are often used to study embryonic development.

In the early stages of embryonic development, a sea urchin embryo is similar to a mammalian embryo.

(a) A scientist observed a fertilised sea urchin egg during the first few days after fertilisation.

Explain how this fertilised egg became a hollow structure.

(3)

.....

.....

.....

.....

.....

.....

.....

.....

(b) The scientist found that 128 cells were present after a few days.

Calculate the number of rounds of cleavage that had taken place.

(1)

Answer

Q2.

A zygote is formed when gametes fuse at fertilisation.

Describe how the process of fertilisation results in the formation of a zygote from the gametes in humans.

(3)

.....

.....

.....

.....

.....

.....

.....

.....


.....

(Total for question = 3 marks)

Q3.

In vitro fertilisation (IVF) is a technique in which eggs are taken from the ovaries and fertilised with sperm in a laboratory.

The resulting embryos are cultured until the four-cell stage, as shown in the diagram.



Two screening techniques used to identify embryos with chromosomal abnormalities are:

- polar body biopsy in which the polar bodies are removed and the chromosomes analysed

- pre-implantation genetic diagnosis (PGD) in which one cell from the four-cell stage embryo is removed and the chromosomes analysed.

Embryos without chromosome abnormalities are placed into the mother's uterus.

The success rates of both techniques are shown in the table.

Technique	Percentage of embryos that survive screening (%)	Percentage of embryos transferred to the uterus that lead to the birth of baby (%)
Polar body biopsy	87	21
PGD	74	29
Control	IVF with no screening	16

Analyse the data to comment on the effectiveness of the two techniques.

(4)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(Total for question = 4 marks)

Mitochondrial disorders may be caused by mutations in the genes coding for mitochondrial components. Some of these genes are found in mitochondrial DNA (mtDNA) and some are found in nuclear DNA.

Leigh syndrome is an example of a mitochondrial disorder. In this syndrome, a number of different proteins involved in respiration are affected.

These mutations may be inherited or may occur when DNA replicates.

Explain why mutations in nuclear DNA can be inherited from either the mother or the father whereas mutations in mtDNA are only inherited from the mother.

(2)

.....

.....

.....

.....

(Total for question = 2 marks)

Q5.

Leigh syndrome is a genetic disorder inherited from the mother. The mother carries genes for the disorder in her mitochondrial DNA.

Scientists have developed a technique for producing 'three-parent' babies.

This ensures that a mother with Leigh syndrome will not pass on the genes for this disorder to her baby.

The technique involves:

- removing the nucleus from the ovum of the mother
- removing the nucleus from the ovum of a donor female to produce an enucleated ovum
- inserting the nucleus from the ovum of the mother into the enucleated donor ovum
- fertilising this ovum with the sperm of the father to produce a zygote
- implanting the resulting embryo into the uterus of the mother.

(i) The 'three-parent' baby produced by this technique will inherit mitochondrial DNA from the

(1)

☒ **A** donor female

- ☐ **B** donor female and father
- ☐ **C** mother
- ☐ **D** mother and father

(ii) Explain the importance of DNA replication during the development of this zygote into a blastocyst.

(3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

(Total for question = 4 marks)

Q6.

Answer the question with a cross in the box you think is correct ☐. If you change your mind about an answer, put a line through the box ☐ and then mark your new answer with a cross ☐.

(i) The table shows the chromosome number and chromatid number of some of the cells formed during oogenesis in humans.

Cell type	Chromosome number	Chromatid number
ovum	23	23
primary oocyte	46	92
secondary oocyte	23	46

Which of the cells are haploid?

(1)

- ☐ **A** ovum and primary oocyte
- ☐ **B** ovum and secondary oocyte
- ☐ **C** primary oocyte and secondary oocyte
- ☐ **D** ovum alone

(ii) Explain how meiosis produces new combinations of alleles in gametes.

(4)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....


(Total for question = 5 marks)

Q7.


The diagram shows the structure of an animal cell.



The diagram shows the development of a zygote into a blastocyst.



The graph shows how the ratio of the volume of the nucleus to the volume of the cytoplasm of each embryonic cell changes as the blastocyst develops.



(i) The ratio shown in the graph can be calculated using the formula

$$\text{ratio} = \frac{\text{volume of nucleus}}{\text{total volume of cell} - \text{volume of nucleus}}$$

On day 2, the volume of the cell nucleus was $900 \mu\text{m}^3$.

Calculate the total volume of this cell on day 2.

Give your answer to two significant figures.

(3)

Answer μm^3

(ii) Comment on the changes in the ratios as the zygote develops into a blastocyst.

(3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(Total for question = 6 marks)

Mark Scheme

Q1.

Question Number	Acceptable Answer	Additional Guidance	Mark
(a)	An explanation that makes reference to the following points: • increase in number of cells by mitosis without growth (1) • therefore cells are progressively smaller (1) • cells form a thin layer surrounding a fluid-filled cavity (1)		(3)

Question Number	Acceptable Answer	Additional Guidance	Mark
(b)	• correct calculation (1)	<u>Example of calculation</u> ($\log_2 128 = $) 7 Allow mark for correct answer with no working	(1)

Q2.

Question Number	Answer	Additional Guidance	Mark
	A description that makes reference to three of the following: • (contact between sperm and secondary oocyte results in) <u>acrosome</u> reaction (1) • meiosis is completed (1) • <u>cortical</u> reaction takes place (1) • fusion of sperm {nucleus / genetic material} with {nucleus genetic material} of ovum (1)	PENALISE 'egg' once ACCEPT ovum / egg cell / female gamete description e.g. enzymes are released from the <u>acrosome</u> DO NOT ACCEPT if described after cortical reaction or fusion of two cells ACCEPT description e.g. <u>cortical</u> granules are released that {hardens the membrane / forms a fertilisation membrane} ACCEPT egg cell / nuclei of the gametes	(3) EXP

Q3.

Question Number	Answer	Additional Guidance	Mark
	An answer that makes reference to four of the following: {screening / both techniques} are {more effective / produce more births} than control (1) - more embryos survive screening with polar body biopsy than PGD / polar body biopsy is less damaging than PGD (1) - PGD more effective (than polar body biopsy) / PGD produces more births (than polar body biopsy) (1) (because) PGD produces 21% births compared to 18% with polar body (1) - PGD detects abnormalities in both paternal and maternal chromosomes / polar body biopsy only checks for maternal chromosome abnormalities (1)	ACCEPT converse ACCEPT converse ACCEPT converse ACCEPT 21.46 / 21.5 / 18.27 / 18.3 ACCEPT 0.21 / 0.215 / 0.18 / 0.183 / 0.1827 DO NOT ACCEPT just 21% alone ACCEPT ideas that polar body biopsy only screens for abnormalities from mother / PGD screens both parents	(4)

Q4.

Question Number	Answer	Additional Guidance	Mark
	An explanation that makes reference to the following: - because nuclear DNA is present in both the male and female {gametes / sex cells} (1) - because mitochondria are present in the { female gamete / ovum / secondary oocyte / egg cell} and not the sperm head (1)	Accept in male and female nuclei in the context of fertilisation Accept mitochondria are present in sperm {neck / mid piece} / mitochondria (DNA) not released by the sperm	(2)

Q5.

Question Number	Answer	Mark
(i)	The only correct answer is A <i>B is not correct because the father's mitochondria do not enter the ovum on fertilisation</i> <i>C is not correct because only the nucleus was used from the mother</i> <i>D is not correct because only the nucleus was used from the mother and the father's mitochondria do not enter the ovum on fertilisation</i>	(1)

Question Number	Answer	Additional Guidance	Mark
(ii)	An explanation that makes reference to three of the following: - zygote divides by mitosis (several times to form blastocyst) (1) - to make identical copies of the {DNA (molecules) / chromatids} (1) - so that all cells (in the blastocyst) {will be diploid / have two copies of each chromosome} (1) - so that when the mitochondria divide they will have a copy of the DNA (1)	ACCEPT genetically-identical (daughter) cells / same genetic information ACCEPT correct number of chromosome / 46 chromosomes / 23 pairs ACCEPT mitochondrial DNA divides	(3)

Q6.

Question Number	Answer	Additional Guidance	Mark
(i)	B ovum and secondary oocyte A is incorrect because secondary oocyte is diploid C is incorrect because the secondary oocyte is diploid D is incorrect because the primary oocyte is diploid		1
(ii)	<i>An explanation that makes reference to:</i> - crossing over (1) - which swaps {alleles / DNA / genes} between {homologous chromosomes} (1) - independent / random assortment (1) - because it is random movement of homologous chromosomes to poles (1)	Allow random movement of paternal and maternal chromosomes / random combinations of paternal and maternal chromosomes	4

Q7.

Question Number	Answer	Additional Guidance	Mark
(i)	- correct reading from graph (1) - correct calculation of mean total cell volume (1) - correct conversion into two significant figures (1)	Example: 17 or 16 952.94 or 956.25 (ignore dps) 950 or 960 (952.94 or 956.25 gains two marks) (950 or 960 gains three marks) Correct answer with no working gains full marks	3 exp

Question Number	Answer	Additional Guidance	Mark
(ii)	An answer that makes reference to three of the following points: • (ratio) ICM increases up to 3/4 days (at a steady rate) / trophoctoderm cells increase up to 3/4/5 days (1) • (after 3/4 days) ratios increases more steeply for ICM cells / (after 3/4/5 days) levels off for trophoctoderm cells (1) • (as ratio increases) {volume of cytoplasm / volume of cells} decreases (over time) (for ICM cells) (1) • cells are differentiating (1)	Accept ratio increases up to 3 days Accept faster increase for ICM cells Accept ICM divide faster than trophoctoderm cells	3 exp