Please check the examination details below before entering your candidate information

Candidate surname StemUp Model Solution | Other names

Centre Number | Candidate Number

Pearson Edexcel Level 3 GCE

Time 1 hour 30 minutes | **Paper reference** 9FM0/01

Further Mathematics

Advanced

PAPER 1: Core Pure Mathematics 1

You must have:
Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for algebraic manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 10 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶

P71776A

©2022 Pearson Education Ltd.

Q:1/1/1/1/

P 7 1 7 7 6 A 0 1 3 2

Pearson

1. $f(z) = z^3 + az + 52$ where a is a real constant

Given that $2 - 3i$ is a root of the equation $f(z) = 0$

(a) write down the other complex root.

(1)

(b) Hence

(i) solve completely $f(z) = 0$

(ii) determine the value of a

(4)

(c) Show all the roots of the equation $f(z) = 0$ on a single Argand diagram.

(1)

(a) $2 + 3i$

(b)(i) Sum of complex roots $= (2 + 3i) + (2 - 3i) = 4$

Product $= (2 + 3i)(2 - 3i) = 4 + 9 = 13$

so $z^2 - 4z + 13$ is a factor of $f(z)$

$z^3 + az + 52 = (z^2 - 4z + 13)(z + k)$: $13k = 52 \Rightarrow k = 4$

$f(z) = (z^2 - 4z + 13)(z + 4)$


Roots: $z = 2 + 3i$, $z = 2 - 3i$, $z = -4$

(ii) $(z^2 - 4z + 13)(z + 4) = z^3 + 4z^2 - 4z^2 - 16z + 13z + 52$

$= z^3 - 3z + 52$

$\therefore a = -3$

(c)



DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 1 continued

(Total for Question 1 is 6 marks)

2. **In this question you must show all stages of your working.**

Solutions relying entirely on calculator technology are not acceptable.

Determine the values of x for which

$$64\cosh^4 x - 64\cosh^2 x - 9 = 0$$

Give your answers in the form $q\ln 2$ where q is rational and in simplest form.

(4)

Let $c = \cosh^2 x$: $64c^2 - 64c - 9 = 0$

$(8c - 9)(8c + 1) = 0$

$\cosh^2 x = \frac{9}{8}$ $\left(\cosh^2 x = -\frac{1}{8}\right.$ rejected since $\cosh^2 x \geq 1)$

$\cosh x = \frac{3}{2\sqrt{2}} = \frac{3\sqrt{2}}{4}$ $(\cosh x > 0)$

$\frac{1}{2}(e^x + e^{-x}) = \frac{3\sqrt{2}}{4} \Rightarrow e^x + e^{-x} = \frac{3\sqrt{2}}{2}$

$2e^{2x} - 3\sqrt{2}\,e^x + 2 = 0$

$e^x = \frac{3\sqrt{2} \pm \sqrt{(18 - 16)}}{4} = \frac{3\sqrt{2} \pm \sqrt{2}}{4}$

$e^x = \sqrt{2}$ or $e^x = \frac{\sqrt{2}}{2} = \frac{1}{\sqrt{2}}$

$x = \ln\sqrt{2} = \frac{1}{2}\ln 2$ or $x = \ln 2^{-\frac{1}{2}} = -\frac{1}{2}\ln 2$

$\therefore\ x = \pm\frac{1}{2}\ln 2$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 2 continued

(Total for Question 2 is 4 marks)

3. (a) Determine the general solution of the differential equation

$$\cos x \frac{dy}{dx} + y \sin x = e^{2x} \cos^2 x$$

giving your answer in the form $y = f(x)$

(3)

Given that $y = 3$ when $x = 0$

(b) determine the smallest positive value of x for which $y = 0$

(3)

(a) Divide by cos x: $\frac{dy}{dx} + y \tan x = e^{2x} \cos x$

Integrating factor $= e^{\int \tan x \, dx} = e^{\ln \sec x} = \sec x$

$\sec x \frac{dy}{dx} + y \sec x \tan x = e^{2x} \cos x \sec x = e^{2x}$

$\frac{d}{dx}(y \sec x) = e^{2x} \Rightarrow y \sec x = \int e^{2x} \, dx = \frac{1}{2} e^{2x} + c$

$y = \left(\frac{1}{2} e^{2x} + c\right) \cos x$

(b) $x = 0, y = 3$: $3 = \left(\frac{1}{2} + c\right) \times 1 \Rightarrow c = \frac{5}{2}$

$y = \left(\frac{1}{2} e^{2x} + \frac{5}{2}\right) \cos x$

$y = 0$: $\frac{1}{2} e^{2x} + \frac{5}{2} > 0$ for all x, so $\cos x = 0$

Smallest positive value: $x = \frac{\pi}{2}$ $(= 1.57)$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 3 continued

(Total for Question 3 is 6 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

4. (a) Use the method of differences to prove that for $n > 2$

$$\sum_{r=2}^{n} \ln\left(\frac{r+1}{r-1}\right) \equiv \ln\left(\frac{n(n+1)}{2}\right)$$

(4)

(b) Hence find the exact value of

$$\sum_{r=51}^{100} \ln\left(\frac{r+1}{r-1}\right)^{35}$$

Give your answer in the form $a\ln\left(\frac{b}{c}\right)$ where a, b and c are integers to be determined.

(3)

(a) $\ln\left(\frac{r+1}{r-1}\right) = \ln(r+1) - \ln(r-1)$

$\sum_{r=2}^{n} \ln\left(\frac{r+1}{r-1}\right) = \sum_{r=2}^{n} [\ln(r+1) - \ln(r-1)]$

$r = 2$: $\ln 3 - \ln 1$

$r = 3$: $\ln 4 - \ln 2$

$r = 4$: $\ln 5 - \ln 3$

$r = 5$: $\ln 6 - \ln 4$

⋮ ⋮

$r = n-2$: $\ln(n-1) - \ln(n-3)$

$r = n-1$: $\ln n - \ln(n-2)$

$r = n$: $\ln(n+1) - \ln(n-1)$

All other terms cancel, leaving

$\sum_{r=2}^{n} \ln\left(\frac{r+1}{r-1}\right) = \ln n + \ln(n+1) - \ln 1 - \ln 2$

$= \ln n + \ln(n+1) - \ln 2 \quad (\ln 1 = 0)$

$= \ln\left(\frac{n(n+1)}{2}\right)$ as required

(b) $\sum_{r=51}^{100} \ln\left(\frac{r+1}{r-1}\right)^{35} = 35\sum_{r=51}^{100} \ln\left(\frac{r+1}{r-1}\right)$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

$$= 35\left(\sum_{r=2}^{100} \ln\left(\frac{r+1}{r-1}\right) - \sum_{r=2}^{50} \ln\left(\frac{r+1}{r-1}\right)\right)$$

$$= 35\left(\ln\left(\frac{100 \times 101}{2}\right) - \ln\left(\frac{50 \times 51}{2}\right)\right)$$

$$= 35(\ln 5050 - \ln 1275) = 35 \ln\left(\frac{5050}{1275}\right)$$

$$= 35 \ln\left(\frac{202}{51}\right)$$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

(Total for Question 4 is 7 marks)

5.

$$\mathbf{M} = \begin{pmatrix} a & 2 & -3 \\ 2 & 3 & 0 \\ 4 & a & 2 \end{pmatrix} \quad \text{where } a \text{ is a constant}$$

(a) Show that **M** is non-singular for all values of a.

(2)

(b) Determine, in terms of a, $\mathbf{M}^{-1}$

(4)

(a) $\det M = a(3 \times 2 - 0 \times a) - 2(2 \times 2 - 0 \times 4) + (-3)(2a - 3 \times 4)$

$= 6a - 8 - 6a + 36$

$= 28$

$\det M = 28 \neq 0$ for all values of a, so M is non-singular for all values of a.

(b) Matrix of minors: $\begin{pmatrix} 6 & 4 & 2a - 12 \\ 4 + 3a & 2a + 12 & a^2 - 8 \\ 9 & 6 & 3a - 4 \end{pmatrix}$

Matrix of cofactors: $\begin{pmatrix} 6 & -4 & 2a - 12 \\ -4 - 3a & 2a + 12 & 8 - a^2 \\ 9 & -6 & 3a - 4 \end{pmatrix}$

Transpose: $\begin{pmatrix} 6 & -4 - 3a & 9 \\ -4 & 2a + 12 & -6 \\ 2a - 12 & 8 - a^2 & 3a - 4 \end{pmatrix}$

$M^{-1} = \frac{1}{28}\begin{pmatrix} 6 & -4 - 3a & 9 \\ -4 & 2a + 12 & -6 \\ 2a - 12 & 8 - a^2 & 3a - 4 \end{pmatrix}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 5 continued

(Total for Question 5 is 6 marks)

6. (a) Express as partial fractions

$$\frac{2x^2 + 3x + 6}{(x+1)(x^2+4)}$$

(3)

(b) Hence, show that

$$\int_0^2 \frac{2x^2 + 3x + 6}{(x+1)(x^2+4)}\,\mathrm{d}x = \ln\left(a\sqrt{2}\right) + b\pi$$

where a and b are constants to be determined.

(4)

(a) $\dfrac{2x^2 + 3x + 6}{(x+1)(x^2+4)} \equiv \dfrac{A}{x+1} + \dfrac{Bx + C}{x^2+4}$

$2x^2 + 3x + 6 \equiv A(x^2+4) + (Bx + C)(x+1)$

$x = -1$: $\quad 2 - 3 + 6 = 5A \Rightarrow A = 1$

$x = 0$: $\quad 6 = 4A + C \Rightarrow C = 2$

coeff x^2: $\quad 2 = A + B \Rightarrow B = 1$

$\dfrac{2x^2 + 3x + 6}{(x+1)(x^2+4)} \equiv \dfrac{1}{x+1} + \dfrac{x+2}{x^2+4}$

(b) $\displaystyle\int_0^2 \frac{1}{x+1} + \frac{x}{x^2+4} + \frac{2}{x^2+4}\,dx$

$= \left[\ln(x+1) + \frac{1}{2}\ln(x^2+4) + \arctan\left(\frac{x}{2}\right)\right]_0^2$

$= \left[\ln 3 + \frac{1}{2}\ln 8 + \arctan 1\right] - \left[\ln 1 + \frac{1}{2}\ln 4 + \arctan 0\right]$

$= \ln 3 + \frac{1}{2}\ln 8 - \frac{1}{2}\ln 4 + \frac{\pi}{4} = \ln 3 + \frac{1}{2}\ln 2 + \frac{\pi}{4}$

$= \ln(3\sqrt{2}) + \frac{\pi}{4}$

$a = 3, \quad b = \frac{1}{4}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 6 continued

Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 6 continued

(Total for Question 6 is 7 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

7. Given that $z = a + bi$ is a complex number where a and b are real constants,

(a) show that zz^* is a real number.

(2)

Given that

- $zz^* = 18$
- $\dfrac{z}{z^*} = \dfrac{7}{9} + \dfrac{4\sqrt{2}}{9}\,i$

(b) determine the possible complex numbers z

(5)

(a) $z^* = a - bi$

$zz^* = (a + bi)(a - bi) = a^2 - abi + abi - b^2i^2 = a^2 + b^2$

a and b are real, so $a^2 + b^2$ is real $\therefore$ zz^* is a real number

(b) $\dfrac{z}{z^*} = \dfrac{z \times z}{z^* \times z} = \dfrac{z^2}{zz^*} = \dfrac{z^2}{18}$

$\dfrac{z^2}{18} = \dfrac{7}{9} + \dfrac{4\sqrt{2}}{9}i \Rightarrow z^2 = 14 + 8\sqrt{2}\,i$

$(a + bi)^2 = a^2 - b^2 + 2abi$

$a^2 - b^2 = 14$ and $2ab = 8\sqrt{2}$

$zz^* = 18$: $a^2 + b^2 = 18$

Adding: $2a^2 = 32 \Rightarrow a^2 = 16 \Rightarrow a = \pm 4$

$b^2 = 18 - 16 = 2 \Rightarrow b = \pm\sqrt{2}$

$2ab = 8\sqrt{2} > 0$ so a and b have the same sign:

$a = 4, b = \sqrt{2}$ or $a = -4, b = -\sqrt{2}$

$z = 4 + \sqrt{2}\,i$ or $z = -4 - \sqrt{2}\,i$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 7 continued

Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 7 continued

(Total for Question 7 is 7 marks)

8. (a) Given

$$z^n + \frac{1}{z^n} = 2\cos n\theta \qquad n \in \mathbb{N}$$

show that

$$32\cos^6\theta \equiv \cos 6\theta + 6\cos 4\theta + 15\cos 2\theta + 10$$

(5)




Figure 1




Figure 2

Figure 1 shows a solid paperweight with a flat base.

Figure 2 shows the curve with equation

$$y = H\cos^3\left(\frac{x}{4}\right) \qquad -4 \leqslant x \leqslant 4$$

where H is a positive constant and x is in radians.

The region R, shown shaded in Figure 2, is bounded by the curve, the line with equation $x = -4$, the line with equation $x = 4$ and the x-axis.

The paperweight is modelled by the solid of revolution formed when R is rotated **180°** about the x-axis.

Given that the maximum height of the paperweight is 2 cm,

(b) write down the value of H.

(1)

(c) Using algebraic integration and the result in part (a), determine, in cm^3, the volume of the paperweight, according to the model. Give your answer to 2 decimal places.

[*Solutions based entirely on calculator technology are not acceptable.*]

(5)

(d) State a limitation of the model.

(1)

(a) $z = \cos\theta + i\sin\theta$: $\left(z + \frac{1}{z}\right)^6 = (2\cos\theta)^6 = 64\cos^6\theta$

$\left(z + \frac{1}{z}\right)^6 = z^6 + 6z^5\left(\frac{1}{z}\right) + 15z^4\left(\frac{1}{z^2}\right) + 20z^3\left(\frac{1}{z^3}\right) + 15z^2\left(\frac{1}{z^4}\right) + 6z\left(\frac{1}{z^5}\right) + \frac{1}{z^6}$

$= \left(z^6 + \frac{1}{z^6}\right) + 6\left(z^4 + \frac{1}{z^4}\right) + 15\left(z^2 + \frac{1}{z^2}\right) + 20$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 8 continued

$64\cos^6\theta = 2\cos 6\theta + 6(2\cos 4\theta) + 15(2\cos 2\theta) + 20$

$64\cos^6\theta = 2\cos 6\theta + 12\cos 4\theta + 30\cos 2\theta + 20$

$\div 2$: $32\cos^6\theta \equiv \cos 6\theta + 6\cos 4\theta + 15\cos 2\theta + 10$ as required

(b) $H = 2$

(c) Rotating through 180° gives half the solid of revolution:

$$V = \frac{1}{2}\pi\int_{-4}^{4} y^2\,dx = \frac{1}{2}\pi\int_{-4}^{4}\left(2\cos^3\left(\frac{x}{4}\right)\right)^2 dx = 2\pi\int_{-4}^{4}\cos^6\left(\frac{x}{4}\right)dx$$

Using (a) with $\theta = \frac{x}{4}$: $\cos^6\left(\frac{x}{4}\right) = \frac{1}{32}\left(\cos\frac{6x}{4} + 6\cos\frac{4x}{4} + 15\cos\frac{2x}{4} + 10\right)$

$$V = \frac{2\pi}{32}\int_{-4}^{4}\left(\cos\frac{3x}{2} + 6\cos x + 15\cos\frac{x}{2} + 10\right)dx$$

$$= \frac{\pi}{16}\left[\frac{2}{3}\sin\frac{3x}{2} + 6\sin x + 30\sin\frac{x}{2} + 10x\right]_{-4}^{4}$$

$$= \frac{\pi}{16} \times 2\left(\frac{2}{3}\sin 6 + 6\sin 4 + 30\sin 2 + 40\right)$$

$$= \frac{\pi}{8} \times 62.5518\ldots = 24.5640\ldots$$

Volume = 24.56 cm^3

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 8 continued

(d) The curve may not be an exact model of the shape of the paperweight (the measurements may not be accurate / the paperweight may not be smooth).

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 8 continued

(Total for Question 8 is 12 marks)

9. (i) (a) Explain why $\int_0^\infty \cosh x \, dx$ is an improper integral.

(1)

(b) Show that $\int_0^\infty \cosh x \, dx$ is divergent.

(3)

(ii) $4\sinh x = p\cosh x$ where p is a real constant

Given that this equation has real solutions, determine the range of possible values for p

(2)

(i)(a) The interval of integration is unbounded: the upper limit is infinite.

(b) $\int_0^\infty \cosh x \, dx = \lim_{t\to\infty} \int_0^t \cosh x \, dx$

$= \lim_{t\to\infty} [\sinh x]_0^t = \lim_{t\to\infty} (\sinh t - \sinh 0) = \lim_{t\to\infty} \sinh t$

$\sinh t = \frac{1}{2}(e^t - e^{-t})$: as $t \to \infty$, $e^t \to \infty$ and $e^{-t} \to 0$, so $\sinh t \to \infty$

The limit does not exist (finite), therefore the integral is divergent.

(ii) $4\sinh x = p\cosh x \Rightarrow \frac{\sinh x}{\cosh x} = \frac{p}{4} \Rightarrow \tanh x = \frac{p}{4}$

For real solutions: $-1 < \tanh x < 1 \Rightarrow -1 < \frac{p}{4} < 1$

$-4 < p < 4$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 9 continued

(Total for Question 9 is 6 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

10.




Figure 3

The motion of a pendulum, shown in Figure 3, is modelled by the differential equation

$$\frac{d^2\theta}{dt^2} + 9\theta = \frac{1}{2}\cos 3t$$

where θ is the angle, in radians, that the pendulum makes with the downward vertical, t seconds after it begins to move.

(a) (i) Show that a particular solution of the differential equation is

$$\theta = \frac{1}{12}t\sin 3t$$

(4)

(ii) Hence, find the general solution of the differential equation.

(4)

Initially, the pendulum

- makes an angle of $\frac{\pi}{3}$ radians with the downward vertical
- is at rest

Given that, 10 seconds after it begins to move, the pendulum makes an angle of α radians with the downward vertical,

(b) determine, according to the model, the value of α to 3 significant figures.

(4)

Given that the true value of α is 0.62

(c) evaluate the model.

(1)

The differential equation

$$\frac{d^2\theta}{dt^2} + 9\theta = \frac{1}{2}\cos 3t$$

models the motion of the pendulum as moving with forced harmonic motion.

(d) Refine the differential equation so that the motion of the pendulum is simple harmonic motion.

(1)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 10 continued

(a)(i) $\theta = \frac{1}{12} t \sin 3t$

$\frac{d\theta}{dt} = \frac{1}{12} \sin 3t + \frac{1}{4} t \cos 3t$

$\frac{d^2\theta}{dt^2} = \frac{1}{4} \cos 3t + \frac{1}{4} \cos 3t - \frac{3}{4} t \sin 3t = \frac{1}{2} \cos 3t - \frac{3}{4} t \sin 3t$

$\frac{d^2\theta}{dt^2} + 9\theta = \frac{1}{2} \cos 3t - \frac{3}{4} t \sin 3t + 9\left(\frac{1}{12} t \sin 3t\right)$

$= \frac{1}{2} \cos 3t - \frac{3}{4} t \sin 3t + \frac{3}{4} t \sin 3t = \frac{1}{2} \cos 3t$

so $\theta = \frac{1}{12} t \sin 3t$ is a particular solution

(ii) $m^2 + 9 = 0 \Rightarrow m = \pm 3i$

CF: $\theta = A \cos 3t + B \sin 3t$

General solution = CF + PI:

$\theta = A \cos 3t + B \sin 3t + \frac{1}{12} t \sin 3t$

(b) $t = 0, \theta = \frac{\pi}{3}$: $A = \frac{\pi}{3}$

$\frac{d\theta}{dt} = -3A \sin 3t + 3B \cos 3t + \frac{1}{12} \sin 3t + \frac{1}{4} t \cos 3t$

$t = 0, \frac{d\theta}{dt} = 0$ (at rest): $3B = 0 \Rightarrow B = 0$

$\theta = \frac{\pi}{3} \cos 3t + \frac{1}{12} t \sin 3t$

$t = 10$: $\alpha = \frac{\pi}{3} \cos 30 + \frac{10}{12} \sin 30 = 0.16153... - 0.82336...$

$= -0.66183...$

$\alpha = -0.662$ (an angle of 0.662 radians, on the other side of the vertical)

Question 10 continued

(c) The model gives an angle of 0.662 radians, which is close to the true value 0.62 (about 7% too large), so the model is good (at $t = 10$).

(d) $\frac{d^2\theta}{dt^2} + 9\theta = 0$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 10 continued

Question 10 continued

(Total for Question 10 is 14 marks)

TOTAL FOR PAPER IS 75 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA