Please check the examination details below before entering your candidate information

Candidate surname StemUp Model Solution | Other names

Centre Number | Candidate Number

Pearson Edexcel Level 3 GCE

Time 1 hour 30 minutes

Paper reference **9FM0/02**

Further Mathematics

Advanced

PAPER 2: Core Pure Mathematics 2

You must have:
Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
  – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 9 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
  – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶

P71801A

©2022 Pearson Education Ltd.

Q:1/1/1/1/1/

P 7 1 8 0 1 A 0 1 3 2

Pearson

1. A student was asked to answer the following:

For the complex numbers $z_1 = 3 - 3i$ and $z_2 = \sqrt{3} + i$, find the value of $\arg\left(\frac{z_1}{z_2}\right)$

The student's attempt is shown below.

Line 1 → $\arg(z_1) = \tan^{-1}\left(\frac{3}{3}\right) = \frac{\pi}{4}$

Line 2 → $\arg(z_2) = \tan^{-1}\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6}$

Line 3 → $\arg\left(\frac{z_1}{z_2}\right) = \frac{\arg(z_1)}{\arg(z_2)}$

Line 4 → $= \left(\frac{\pi}{4}\right) \Big/ \left(\frac{\pi}{6}\right) = \frac{3}{2}$

The student made errors in line 1 and line 3

Correct the error that the student made in

(a) (i) line 1

(ii) line 3

(2)

(b) Write down the correct value of $\arg\left(\frac{z_1}{z_2}\right)$

(1)

(a)(i) Line 1: $z_1 = 3 - 3i$ is in the 4th quadrant, so $\arg(z_1)$ is negative

$\arg(z_1) = \tan^{-1}\frac{-3}{3} = \tan^{-1}(-1) = -\frac{\pi}{4}$

(ii) Line 3: the arguments should be subtracted, not divided:

$\arg\left(\frac{z_1}{z_2}\right) = \arg(z_1) - \arg(z_2)$

(b) $\arg\left(\frac{z_1}{z_2}\right) = -\frac{\pi}{4} - \frac{\pi}{6} = -\frac{5\pi}{12}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 1 continued

(Total for Question 1 is 3 marks)

2. **In this question you must show all stages of your working.**

A college offers only three courses: Construction, Design and Hospitality.

Each student enrols on just one of these courses.

In 2019, there was a total of 1110 students at this college.

There were 370 more students enrolled on Construction than Hospitality.

In 2020 the number of students enrolled on

- Construction **increased** by 1.25%
- Design **increased** by 2.5%
- Hospitality **decreased** by 2%

In 2020, the total number of students at the college increased by 0.27% to 2 significant figures.

(a) (i) Define, for each course, a variable for the number of students enrolled on that course in 2019.

(ii) Using your variables from part (a)(i), write down **three** equations that model this situation.

(4)

(b) By forming and solving a matrix equation, determine how many students were enrolled on each of the three courses in 2019.

(4)

(a)(i) Let C = number of students enrolled on Construction in 2019

D = number of students enrolled on Design in 2019

H = number of students enrolled on Hospitality in 2019

(ii) Total 2019: $C + D + H = 1110$

370 more on Construction: $C - H = 370$

Total 2020 = $1110 \times 1.0027 = 1112.997 \approx 1113$

$1.0125C + 1.025D + 0.98H = 1113$

(b) $$\begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & -1 \\ 1.0125 & 1.025 & 0.98 \end{pmatrix} \begin{pmatrix} C \\ D \\ H \end{pmatrix} = \begin{pmatrix} 1110 \\ 370 \\ 1113 \end{pmatrix}$$

$$\begin{pmatrix} C \\ D \\ H \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & -1 \\ 1.0125 & 1.025 & 0.98 \end{pmatrix}^{-1} \begin{pmatrix} 1110 \\ 370 \\ 1113 \end{pmatrix} = \begin{pmatrix} 720 \\ 40 \\ 350 \end{pmatrix}$$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 2 continued

So in 2019: 720 students enrolled on Construction,
40 students enrolled on Design
and 350 students enrolled on Hospitality.

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 2 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 2 continued

(Total for Question 2 is 8 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

3. $\mathbf{M} = \begin{pmatrix} 3 & a \\ 0 & 1 \end{pmatrix}$ where a is a constant

(a) Prove by mathematical induction that, for $n \in \mathbb{N}$

$$\mathbf{M}^n = \begin{pmatrix} 3^n & \frac{a}{2}(3^n - 1) \\ 0 & 1 \end{pmatrix}$$

(6)

Triangle T has vertices A, B and C.

Triangle T is transformed to triangle T' by the transformation represented by $\mathbf{M}^n$ where $n \in \mathbb{N}$

Given that

- triangle T has an area of $5\,\text{cm}^2$
- triangle T' has an area of $1215\,\text{cm}^2$
- vertex $A(2, -2)$ is transformed to vertex $A'(123, -2)$

(b) determine

(i) the value of n

(ii) the value of a

(5)

(a) $n = 1$: $M^1 = \begin{pmatrix} 3^1 & \frac{1}{2}a(3^1 - 1) \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 3 & a \\ 0 & 1 \end{pmatrix} = M$ so true for $n = 1$

Assume true for $n = k$: $M^k = \begin{pmatrix} 3^k & \frac{1}{2}a(3^k - 1) \\ 0 & 1 \end{pmatrix}$

Then $M^{k+1} = M^k M = \begin{pmatrix} 3^k & \frac{1}{2}a(3^k - 1) \\ 0 & 1 \end{pmatrix}\begin{pmatrix} 3 & a \\ 0 & 1 \end{pmatrix}$

$= \begin{pmatrix} 3 \times 3^k & a \times 3^k + \frac{1}{2}a(3^k - 1) \\ 0 & 1 \end{pmatrix}$

$= \begin{pmatrix} 3^{k+1} & \frac{1}{2}a(2 \times 3^k + 3^k - 1) \\ 0 & 1 \end{pmatrix}$

$= \begin{pmatrix} 3^{k+1} & \frac{1}{2}a(3 \times 3^k - 1) \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 3^{k+1} & \frac{1}{2}a(3^{k+1} - 1) \\ 0 & 1 \end{pmatrix}$

which is the result with $n = k + 1$.

If the result is true for $n = k$ then it is true for $n = k + 1$, and as it is true for $n = 1$, it is true for all $n \in \mathbb{N}$ by mathematical induction.

(b)(i) $\det(M^n) = 3^n \times 1 - 0 = 3^n$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 3 continued

$5 \times 3^n = 1215 \Rightarrow 3^n = 243 \Rightarrow n = 5$

(ii) $M^5 = \begin{pmatrix} 243 & \frac{1}{2}a(243-1) \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 243 & 121a \\ 0 & 1 \end{pmatrix}$

$\begin{pmatrix} 243 & 121a \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 \\ -2 \end{pmatrix} = \begin{pmatrix} 123 \\ -2 \end{pmatrix}$

$\Rightarrow 486 - 242a = 123$

$242a = 363 \Rightarrow a = 1.5$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 3 continued

(Total for Question 3 is 11 marks)

4. (i) Given that

$$z_1 = 6e^{\frac{\pi}{3}i} \text{ and } z_2 = 6\sqrt{3}e^{\frac{5\pi}{6}i}$$

show that

$$z_1 + z_2 = 12e^{\frac{2\pi}{3}i}$$

(3)

(ii) Given that

$$\arg(z - 5) = \frac{2\pi}{3}$$

determine the least value of $|z|$ as z varies.

(3)

(i) $z_1 = 6(\cos\frac{\pi}{3} + i\sin\frac{\pi}{3}) = 3 + 3\sqrt{3}\,i$

$z_2 = 6\sqrt{3}(\cos\frac{5\pi}{6} + i\sin\frac{5\pi}{6}) = 6\sqrt{3}(-\frac{\sqrt{3}}{2} + \frac{1}{2}i) = -9 + 3\sqrt{3}\,i$

$z_1 + z_2 = (3 + 3\sqrt{3}\,i) + (-9 + 3\sqrt{3}\,i) = -6 + 6\sqrt{3}\,i$

$|z_1 + z_2| = \sqrt{((-6)^2 + (6\sqrt{3})^2)} = \sqrt{(36 + 108)} = \sqrt{144} = 12$

$-6 + 6\sqrt{3}\,i$ is in the 2nd quadrant, so

$\arg(z_1 + z_2) = \pi - \tan^{-1}\frac{6\sqrt{3}}{6} = \pi - \tan^{-1}(\sqrt{3}) = \pi - \frac{\pi}{3} = \frac{2\pi}{3}$

$\therefore\ z_1 + z_2 = 12(\cos\frac{2\pi}{3} + i\sin\frac{2\pi}{3}) = 12e^{2\pi i/3}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

(ii) Locus: half-line from 5 at angle $\frac{2\pi}{3}$ to the positive real axis.


The least $|z|$ is the perpendicular distance from 0 to the half-line (see sketch).

Angle between half-line and real axis

$= \pi - \frac{2\pi}{3} = \frac{\pi}{3}$

$\sin\frac{\pi}{3} = \frac{|z|}{5} \Rightarrow |z| = 5 \times \frac{\sqrt{3}}{2}$

Least value of $|z| = \frac{5\sqrt{3}}{2}$



DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

(Total for Question 4 is 6 marks)

5. (a) Given that

$$y = \arcsin x \qquad -1 \leqslant x \leqslant 1$$

show that

$$\frac{dy}{dx} = \frac{1}{\sqrt{1 - x^2}}$$

(3)

(b) $$f(x) = \arcsin(e^x) \qquad x \leqslant 0$$

Prove that f(x) has no stationary points.

(3)

(a) $y = \arcsin x \Rightarrow \sin y = x$

$\Rightarrow \cos y \frac{dy}{dx} = 1 \Rightarrow \frac{dy}{dx} = \frac{1}{\cos y}$

$\cos^2 y = 1 - \sin^2 y \Rightarrow \cos y = \sqrt{(1 - \sin^2 y)} = \sqrt{(1 - x^2)}$

(positive root as $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ so $\cos y \geq 0$)

$\therefore \frac{dy}{dx} = \frac{1}{\sqrt{(1 - x^2)}}$

(b) $f(x) = \arcsin(e^x) \Rightarrow f'(x) = \frac{1}{\sqrt{(1 - (e^x)^2)}} \times e^x = \frac{e^x}{\sqrt{(1 - e^{2x})}}$

Stationary points need $f'(x) = 0 \Rightarrow e^x = 0$

but $e^x > 0$ for all x, so $e^x = 0$ has no solutions.

Therefore f(x) has no stationary points.

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 5 continued

(Total for Question 5 is 6 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

6. The cubic equation

$$4x^3 + px^2 - 14x + q = 0$$

where p and q are real positive constants, has roots α, β and γ

Given that $\alpha^2 + \beta^2 + \gamma^2 = 16$

(a) show that $p = 12$

(3)

Given that $\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma} = \frac{14}{3}$

(b) determine the value of q

(3)

Without solving the cubic equation,

(c) determine the value of $(\alpha - 1)(\beta - 1)(\gamma - 1)$

(4)

(a) $\alpha + \beta + \gamma = -\frac{p}{4}$ $\quad \alpha\beta + \beta\gamma + \gamma\alpha = \frac{-14}{4} = -\frac{7}{2}$ $\quad \alpha\beta\gamma = -\frac{q}{4}$

$\alpha^2 + \beta^2 + \gamma^2 = (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \beta\gamma + \gamma\alpha)$

$16 = (-\frac{p}{4})^2 - 2(-\frac{7}{2}) = \frac{p^2}{16} + 7$

$\frac{p^2}{16} = 9 \Rightarrow p^2 = 144 \Rightarrow p = 12 \quad (p > 0)$

(b) $\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma} = \frac{\alpha\beta + \beta\gamma + \gamma\alpha}{\alpha\beta\gamma} = \frac{-7/2}{-q/4} = \frac{14}{q}$

$\frac{14}{q} = \frac{14}{3} \Rightarrow q = 3$

(c) $(\alpha - 1)(\beta - 1)(\gamma - 1) = \alpha\beta\gamma - (\alpha\beta + \beta\gamma + \gamma\alpha) + (\alpha + \beta + \gamma) - 1$

$= (-\frac{3}{4}) - (-\frac{7}{2}) + (-\frac{12}{4}) - 1$

$= -\frac{3}{4} + \frac{7}{2} - 3 - 1$

$= -\frac{5}{4}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 6 continued

Question 6 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 6 continued

(Total for Question 6 is 10 marks)

7.




Figure 1

Figure 1 shows a sketch of the curve C with equation

$$r = 1 + \tan\theta \qquad 0 \leqslant \theta < \frac{\pi}{3}$$

Figure 1 also shows the tangent to C at the point A.
This tangent is perpendicular to the initial line.

(a) Use differentiation to prove that the polar coordinates of A are $\left(2, \frac{\pi}{4}\right)$

(4)

The finite region R, shown shaded in Figure 1, is bounded by C, the tangent at A and the initial line.

(b) Use calculus to show that the exact area of R is $\frac{1}{2}(1 - \ln 2)$

(6)

(a) $x = r\cos\theta = (1 + \tan\theta)\cos\theta = \cos\theta + \sin\theta$

$\frac{dx}{d\theta} = -\sin\theta + \cos\theta$

Tangent perpendicular to initial line $\Rightarrow \frac{dx}{d\theta} = 0$

$\Rightarrow \sin\theta = \cos\theta \Rightarrow \tan\theta = 1 \Rightarrow \theta = \frac{\pi}{4}$

$r = 1 + \tan\frac{\pi}{4} = 1 + 1 = 2 \quad \therefore A \text{ is } (2, \frac{\pi}{4})$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 7 continued

(b) Area between curve, OA and initial line:

$$\tfrac{1}{2}\int_0^{\pi/4} (1+\tan\theta)^2\,d\theta = \tfrac{1}{2}\int_0^{\pi/4} (1+2\tan\theta+\tan^2\theta)\,d\theta$$

$$= \tfrac{1}{2}\int_0^{\pi/4} (\sec^2\theta+2\tan\theta)\,d\theta$$

$$= \tfrac{1}{2}\left[\tan\theta + 2\ln|\sec\theta|\right]_0^{\pi/4}$$

$$= \tfrac{1}{2}\left[\left(\tan\tfrac{\pi}{4} + 2\ln\left(\sec\tfrac{\pi}{4}\right)\right) - (\tan 0 + 2\ln(\sec 0))\right]$$

$$= \tfrac{1}{2}\left[(1+2\ln\sqrt{2}) - (0+0)\right] = \tfrac{1}{2}(1+\ln 2) = \tfrac{1}{2} + \tfrac{1}{2}\ln 2$$

A has Cartesian coordinates $\left(2\cos\tfrac{\pi}{4}, 2\sin\tfrac{\pi}{4}\right) = (\sqrt{2}, \sqrt{2})$

Triangle formed by O, A and the foot of the tangent $(\sqrt{2}, 0)$:

area $= \tfrac{1}{2} \times \sqrt{2} \times \sqrt{2} = 1$

Area of R = area of triangle − area between curve and OA

$= 1 - (\tfrac{1}{2} + \tfrac{1}{2}\ln 2)$

$= \tfrac{1}{2} - \tfrac{1}{2}\ln 2 = \tfrac{1}{2}(1 - \ln 2)$

Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 7 continued

(Total for Question 7 is 10 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

8. Two birds are flying towards their nest, which is in a tree.

Relative to a fixed origin, the flight path of each bird is modelled by a straight line.

In the model, the equation for the flight path of the first bird is

$$\mathbf{r}_1 = \begin{pmatrix} -1 \\ 5 \\ 2 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ a \\ 0 \end{pmatrix}$$

and the equation for the flight path of the second bird is

$$\mathbf{r}_2 = \begin{pmatrix} 4 \\ -1 \\ 3 \end{pmatrix} + \mu \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}$$

where λ and μ are scalar parameters and a is a constant.

In the model, the angle between the birds' flight paths is 120°

(a) Determine the value of a.

(4)

(b) Verify that, according to the model, there is a common point on the flight paths of the two birds and find the coordinates of this common point.

(5)

The position of the nest is modelled as being at this common point.

The tree containing the nest is in a park.

The ground level of the park is modelled by the plane with equation

$$2x - 3y + z = 2$$

(c) Hence determine the shortest distance from the nest to the ground level of the park.

(3)

(d) By considering the model, comment on whether your answer to part (c) is reliable, giving a reason for your answer.

(1)

(a) $\begin{pmatrix} 2 \\ a \\ 0 \end{pmatrix} \cdot \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} = 0 + a + 0 = a$ $\quad |d_1| = \sqrt{(4 + a^2)}, \quad |d_2| = \sqrt{2}$

$\cos 120° = \dfrac{a}{\sqrt{(4 + a^2)}\,\sqrt{2}} \Rightarrow \dfrac{a}{\sqrt{(4 + a^2)}\,\sqrt{2}} = -\dfrac{1}{2}$

$2a = -\sqrt{2}\,\sqrt{(4 + a^2)} \Rightarrow 4a^2 = 2(4 + a^2) \Rightarrow 2a^2 = 8 \Rightarrow a^2 = 4$

a must be negative (as $\cos 120° < 0$) $\Rightarrow a = -2$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 8 continued

(b) i: $-1 + 2\lambda = 4 \Rightarrow \lambda = \frac{5}{2}$

k: $2 = 3 - \mu \Rightarrow \mu = 1$

j: LHS $= 5 - 2\lambda = 5 - 2\left(\frac{5}{2}\right) = 0$, RHS $= -1 + \mu = -1 + 1 = 0$

LHS = RHS so all three equations are consistent:
the flight paths have a common point.

$$\mathbf{r}_1 = \begin{pmatrix} -1 \\ 5 \\ 2 \end{pmatrix} + \frac{5}{2}\begin{pmatrix} 2 \\ -2 \\ 0 \end{pmatrix} = \begin{pmatrix} 4 \\ 0 \\ 2 \end{pmatrix}$$

Common point is (4, 0, 2)

(c) Shortest distance from (4, 0, 2) to $2x - 3y + z = 2$:

$$d = \frac{|2(4) - 3(0) + 1(2) - 2|}{\sqrt{(2^2 + (-3)^2 + 1^2)}} = \frac{8}{\sqrt{14}} = \frac{4\sqrt{14}}{7} = 2.138\ldots = 2.14 \text{ (3 s.f.)}$$

(d) Not reliable: the birds will not fly in exactly straight lines,
and the ground of the park is unlikely to be a flat plane.

Question 8 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 8 continued

(Total for Question 8 is 13 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

9. $y = \cosh^n x \qquad n \geqslant 5$

(a) (i) Show that

$$\frac{\mathrm{d}^2 y}{\mathrm{d}x^2} = n^2 \cosh^n x - n(n-1)\cosh^{n-2} x$$

(4)

(ii) Determine an expression for $\dfrac{\mathrm{d}^4 y}{\mathrm{d}x^4}$

(2)

(b) Hence determine the first three non-zero terms of the Maclaurin series for y, giving each coefficient in simplest form.

(2)

(a)(i) $\frac{dy}{dx} = n\cosh^{n-1}x \sinh x$

$\frac{d^2y}{dx^2} = n(n-1)\cosh^{n-2}x \sinh^2 x + n\cosh^{n-1}x \cosh x$

$= n(n-1)\cosh^{n-2}x (\cosh^2 x - 1) + n\cosh^n x$

$= n(n-1)\cosh^n x - n(n-1)\cosh^{n-2}x + n\cosh^n x$

$= (n^2 - n + n)\cosh^n x - n(n-1)\cosh^{n-2}x$

$= n^2\cosh^n x - n(n-1)\cosh^{n-2}x$

(ii) $\frac{d^3y}{dx^3} = n^3\cosh^{n-1}x \sinh x - n(n-1)(n-2)\cosh^{n-3}x \sinh x$

$\frac{d^4y}{dx^4} = n^3(n-1)\cosh^{n-2}x \sinh^2 x + n^3\cosh^n x$

$- n(n-1)(n-2)(n-3)\cosh^{n-4}x \sinh^2 x - n(n-1)(n-2)\cosh^{n-2}x$

(b) At $x = 0$: $\cosh 0 = 1$, $\sinh 0 = 0$

$y(0) = 1$, $y'(0) = 0$, $y''(0) = n^2 - n(n-1) = n$, $y'''(0) = 0$

$y^{(4)}(0) = n^3 - n(n-1)(n-2) = n^3 - (n^3 - 3n^2 + 2n) = 3n^2 - 2n$

$y = y(0) + y'(0)x + \frac{y''(0)}{2!}x^2 + \frac{y'''(0)}{3!}x^3 + \frac{y^{(4)}(0)}{4!}x^4 + \ldots$

$y = 1 + \frac{n}{2}x^2 + \frac{3n^2 - 2n}{24}x^4 + \ldots$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 9 continued

Question 9 continued

(Total for Question 9 is 8 marks)

TOTAL FOR PAPER IS 75 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA