Please check the examination details below before entering your candidate information

Candidate surname StemUp Model Solution | Other names

Centre Number | Candidate Number

Pearson Edexcel Level 3 GCE

Monday 3 June 2024

Afternoon (Time: 1 hour 30 minutes)

Paper reference **9FM0/02**

Further Mathematics

Advanced

PAPER 2: Core Pure Mathematics 2

You must have:
Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for algebraic manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
  – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 9 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
  – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶

P75683A

©2024 Pearson Education Ltd.
F:1/1/1/1/1/

P 7 5 6 8 3 A 0 1 3 2

Pearson

1. (a) Using the definition of $\sinh x$ in terms of exponentials, prove that

$$4\sinh^3 x + 3\sinh x \equiv \sinh 3x$$

(2)

(b) Hence solve the equation

$$\sinh 3x = 19\sinh x$$

giving your answers as simplified natural logarithms where appropriate.

(5)

(a) $LHS = 4\left(\frac{e^x - e^{-x}}{2}\right)^3 + 3\left(\frac{e^x - e^{-x}}{2}\right)$

$= 4 \times \frac{e^{3x} - 3e^x + 3e^{-x} - e^{-3x}}{8} + \frac{3e^x - 3e^{-x}}{2}$

$= \frac{e^{3x} - 3e^x + 3e^{-x} - e^{-3x} + 3e^x - 3e^{-x}}{2}$

$= \frac{e^{3x} - e^{-3x}}{2} = \sinh 3x = RHS \quad \therefore 4\sinh^3 x + 3\sinh x \equiv \sinh 3x$

(b) $\sinh 3x = 19\sinh x \Rightarrow 4\sinh^3 x + 3\sinh x = 19\sinh x$

$\Rightarrow 4\sinh^3 x - 16\sinh x = 0$

$\Rightarrow 4\sinh x(\sinh^2 x - 4) = 0$

$\sinh x = 0 \Rightarrow x = 0$

$\sinh^2 x = 4 \Rightarrow \sinh x = \pm 2$

$x = \operatorname{arsinh}(\pm 2) = \ln(\pm 2 + \sqrt{((\pm 2)^2 + 1)}) = \ln(\pm 2 + \sqrt{5})$

$x = \ln(2 + \sqrt{5})$ or $x = \ln(-2 + \sqrt{5}) = \ln(\sqrt{5} - 2)$

Solutions: $x = 0$, $x = \ln(2 + \sqrt{5})$, $x = \ln(\sqrt{5} - 2)$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 1 continued

(Total for Question 1 is 7 marks)

2. $$f(x) = \tanh^{-1}\left(\frac{3-x}{6+x}\right) \qquad |x| < \frac{3}{2}$$

(a) Show that

$$f'(x) = -\frac{1}{2x+3}$$

(4)

(b) Hence determine $f''(x)$

(1)

(c) Hence show that the Maclaurin series for $f(x)$, up to and including the term in x^2, is

$$\ln p + qx + rx^2$$

where p, q and r are constants to be determined.

(3)

(a) $\frac{d}{dx}\left(\frac{3-x}{6+x}\right) = \frac{-(6+x)-(3-x)}{(6+x)^2} = \frac{-9}{(6+x)^2}$

$f'(x) = \frac{1}{1-((3-x)/(6+x))^2} \times \frac{-9}{(6+x)^2}$

$= \frac{(6+x)^2}{(6+x)^2-(3-x)^2} \times \frac{-9}{(6+x)^2}$

$= \frac{-9}{36+12x+x^2-9+6x-x^2} = \frac{-9}{18x+27}$

$= -\frac{1}{2x+3}$

(b) $f'(x) = -(2x+3)^{-1} \Rightarrow f''(x) = 2(2x+3)^{-2} = \frac{2}{(2x+3)^2}$

(c) $f(0) = \tanh^{-1}\left(\frac{1}{2}\right) = \frac{1}{2}\ln\left(\frac{1+\frac{1}{2}}{1-\frac{1}{2}}\right) = \frac{1}{2}\ln 3 = \ln\sqrt{3}$

$f'(0) = -\frac{1}{3} \qquad f''(0) = \frac{2}{3^2} = \frac{2}{9}$

$f(x) \approx f(0) + f'(0)x + \frac{f''(0)}{2!}x^2$

$= \ln\sqrt{3} - \frac{1}{3}x + \frac{2}{9\times 2}x^2$

$f(x) \approx \ln\sqrt{3} - \frac{1}{3}x + \frac{1}{9}x^2 \qquad p = \sqrt{3},\ q = -\frac{1}{3},\ r = \frac{1}{9}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 2 continued

(Total for Question 2 is 8 marks)

3. (a) Explain why

$$\int_{\frac{4}{3}}^{\infty} \frac{1}{9x^2 + 16} \mathrm{d}x$$

is an improper integral.

(1)

(b) Show that

$$\int_{\frac{4}{3}}^{\infty} \frac{1}{9x^2 + 16} \mathrm{d}x = k\pi$$

where k is a constant to be determined.

(4)

(a) The upper limit of the integral is infinite (∞), so it is an improper integral.

(b) $\int \frac{1}{9x^2 + 16}\,dx = \frac{1}{9}\int \frac{1}{x^2 + 16/9}\,dx = \frac{1}{9} \times \frac{3}{4} \arctan\left(\frac{3x}{4}\right) = \frac{1}{12}\arctan\left(\frac{3x}{4}\right)$

$\int_{4/3}^{\infty} \frac{1}{9x^2 + 16}\,dx = \lim_{t \to \infty} \int_{4/3}^{t} \frac{1}{9x^2 + 16}\,dx = \frac{1}{12} \lim_{t \to \infty} \left[\arctan\left(\frac{3x}{4}\right)\right]_{4/3}^{t}$

$= \frac{1}{12}\left(\lim_{t \to \infty} \arctan\left(\frac{3t}{4}\right) - \arctan(1)\right)$

$= \frac{1}{12}\left(\frac{\pi}{2} - \frac{\pi}{4}\right) = \frac{1}{12} \times \frac{\pi}{4} = \frac{\pi}{48}$

$\therefore \int_{4/3}^{\infty} \frac{1}{9x^2 + 16}\,dx = \frac{1}{48}\pi \qquad k = \frac{1}{48}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 3 continued

(Total for Question 3 is 5 marks)

4. Use the method of differences to show that

$$\sum_{r=1}^{n} \frac{2}{(r+4)(r+6)} = \frac{n(an+b)}{30(n+5)(n+6)}$$

where a and b are integers to be determined.

(6)

$\frac{2}{(r+4)(r+6)} \equiv \frac{A}{r+4} + \frac{B}{r+6} \Rightarrow 2 \equiv A(r+6) + B(r+4)$

$r = -4$: $2 = 2A \Rightarrow A = 1$ $\quad$ $r = -6$: $2 = -2B \Rightarrow B = -1$

$\frac{2}{(r+4)(r+6)} \equiv \frac{1}{r+4} - \frac{1}{r+6}$

$\sum_{r=1}^{n} \frac{2}{(r+4)(r+6)} = \sum_{r=1}^{n} \left(\frac{1}{r+4} - \frac{1}{r+6} \right)$

$r = 1$: $\frac{1}{5} - \frac{1}{7}$

$r = 2$: $\frac{1}{6} - \frac{1}{8}$

$r = 3$: $\frac{1}{7} - \frac{1}{9}$

$\vdots$

$r = n-2$: $\frac{1}{n+2} - \frac{1}{n+4}$

$r = n-1$: $\frac{1}{n+3} - \frac{1}{n+5}$

$r = n$: $\frac{1}{n+4} - \frac{1}{n+6}$

$\text{Sum} = \frac{1}{5} + \frac{1}{6} - \frac{1}{n+5} - \frac{1}{n+6} = \frac{11}{30} - \frac{1}{n+5} - \frac{1}{n+6}$

$= \frac{11(n+5)(n+6) - 30(n+6) - 30(n+5)}{30(n+5)(n+6)}$

$= \frac{11n^2 + 121n + 330 - 30n - 180 - 30n - 150}{30(n+5)(n+6)}$

$= \frac{11n^2 + 61n}{30(n+5)(n+6)} = \frac{n(11n+61)}{30(n+5)(n+6)}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

$\therefore\ a = 11,\ b = 61$

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

(Total for Question 4 is 6 marks)

5. The locus C is given by

$$|z - 4| = 4$$

The locus D is given by

$$\arg z = \frac{\pi}{3}$$

(a) Sketch, on the same Argand diagram, the locus C and the locus D

(4)

The set of points A is defined by

$$A = \{z \in \mathbb{C} : |z - 4| \leqslant 4\} \cap \left\{z \in \mathbb{C} : 0 \leqslant \arg z \leqslant \frac{\pi}{3}\right\}$$


(b) Show, by shading on your Argand diagram, the set of points A

(1)

(c) Find the area of the region defined by A, giving your answer in the form $p\pi + q\sqrt{3}$ where p and q are constants to be determined.

(4)

(a), (b)



C: circle, centre 4, radius 4
(touches the imaginary axis at 0)
D: half-line from 0 at angle π/3
to the positive real axis
A: shaded region

(c) C: $(x - 4)^2 + y^2 = 16$ D: $y = \sqrt{3}x$, $x > 0$

$(x - 4)^2 + 3x^2 = 16 \Rightarrow 4x^2 - 8x = 0 \Rightarrow 4x(x - 2) = 0$

$x = 2$ $(x \neq 0)$, $y = 2\sqrt{3}$ so C and D meet at $P(2, 2\sqrt{3})$

Region A = triangle O P X + sector X P (8, 0), where X(4, 0) is the centre

Triangle: $\frac{1}{2} \times 4 \times 2\sqrt{3} = 4\sqrt{3}$

XP = $(-2, 2\sqrt{3})$ makes angle $\frac{2\pi}{3}$ with the positive real axis

Sector: $\frac{1}{2} \times 4^2 \times \frac{2\pi}{3} = \frac{16\pi}{3}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 5 continued

Area of A $= \frac{16}{3}\pi + 4\sqrt{3}$ $\quad p = \frac{16}{3}$, $q = 4$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 5 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 5 continued

(Total for Question 5 is 9 marks)

6. The motion of a particle P along the x-axis is modelled by the differential equation

$$2\frac{d^2x}{dt^2} + 5\frac{dx}{dt} + 2x = 4t + 12$$

where P is x metres from the origin O at time t seconds, $t \geqslant 0$

(a) Determine the general solution of the differential equation.

(6)

(b) Hence determine the particular solution for which $x = 3$ and $\frac{dx}{dt} = -2$ when $t = 0$

(3)

(c) (i) Show that, according to the model, the minimum distance between O and P is $(2 + \ln 2)$ metres.

(ii) Justify that this distance is a minimum.

(4)

For large values of t the particle is expected to move with constant speed.

(d) Comment on the suitability of the model in light of this information.

(1)

(a) $2m^2 + 5m + 2 = 0 \Rightarrow (2m + 1)(m + 2) = 0 \Rightarrow m = -\frac{1}{2}, -2$

CF: $x = Ae^{-\frac{1}{2}t} + Be^{-2t}$

PI: $x = pt + q \Rightarrow \frac{dx}{dt} = p, \frac{d^2x}{dt^2} = 0$

$2(0) + 5p + 2(pt + q) = 4t + 12$

$2p = 4 \Rightarrow p = 2 \quad 5p + 2q = 12 \Rightarrow 10 + 2q = 12 \Rightarrow q = 1$

GS: $x = Ae^{-\frac{1}{2}t} + Be^{-2t} + 2t + 1$

(b) $t = 0, x = 3$: $3 = A + B + 1 \Rightarrow A + B = 2$

$\frac{dx}{dt} = -\frac{1}{2}Ae^{-\frac{1}{2}t} - 2Be^{-2t} + 2$

$t = 0, \frac{dx}{dt} = -2$: $-2 = -\frac{1}{2}A - 2B + 2 \Rightarrow A + 4B = 8$

$\Rightarrow 3B = 6 \Rightarrow B = 2, A = 0$

$x = 2e^{-2t} + 2t + 1$

(c)(i) $\frac{dx}{dt} = -4e^{-2t} + 2 = 0 \Rightarrow e^{-2t} = \frac{1}{2}$

$\Rightarrow -2t = \ln\frac{1}{2} = -\ln 2 \Rightarrow t = \frac{1}{2}\ln 2$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 6 continued

$x = 2e^{-\ln 2} + 2\left(\frac{1}{2}\ln 2\right) + 1 = 2 \times \frac{1}{2} + \ln 2 + 1 = 2 + \ln 2$

$x > 0$ for all $t \geqslant 0$, so x is the distance OP, and $x(0) = 3 > 2 + \ln 2$

$\therefore$ minimum distance between O and P is $(2 + \ln 2)$ metres

(ii) $\frac{d^2x}{dt^2} = 8e^{-2t} > 0$ for all values of t, so the distance is a minimum

(d) For large t, $e^{-2t} \to 0$ so $x \to 2t + 1$ and $\frac{dx}{dt} \to 2$

so the particle moves with constant speed 2 m s^{-1}: the model is suitable.

Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 6 continued

(Total for Question 6 is 14 marks)

7. (a) Determine the roots of the equation

$$z^6 = 1$$

giving your answers in the form $e^{i\theta}$ where $0 \leqslant \theta < 2\pi$

(2)

(b) Show the roots of the equation in part (a) on a single Argand diagram.

(2)

(c) Show that

$$\left(\sqrt{3} + i\right)^6 = -64$$

(2)

(d) Hence, or otherwise, solve the equation

$$z^6 + 64 = 0$$


giving your answers in the form $re^{i\theta}$ where $0 \leqslant \theta < 2\pi$

(3)

(a) $z^6 = 1 = e^{2k\pi i} \Rightarrow z = e^{2k\pi i/6} = e^{k\pi i/3}$, $k = 0, 1, 2, 3, 4, 5$

$z = e^0, e^{\pi i/3}, e^{2\pi i/3}, e^{\pi i}, e^{4\pi i/3}, e^{5\pi i/3}$

(b)



(c) $|\sqrt{3} + i| = \sqrt{(3 + 1)} = 2$, $\arg(\sqrt{3} + i) = \arctan \frac{1}{\sqrt{3}} = \frac{\pi}{6}$

$(\sqrt{3} + i)^6 = (2e^{\pi i/6})^6 = 2^6 e^{\pi i} = 64(\cos \pi + i \sin \pi) = 64(-1) = -64$

(d) $z^6 = -64 = (\sqrt{3} + i)^6 \Rightarrow z = (\sqrt{3} + i) \times e^{k\pi i/3} = 2e^{\pi i/6} \times e^{k\pi i/3}$

$|z|^6 = 64 \Rightarrow r = 2$

$z = 2e^{i(\pi/6 + k\pi/3)}$, $k = 0, 1, 2, 3, 4, 5$

$z = 2e^{\pi i/6}, 2e^{\pi i/2}, 2e^{5\pi i/6}, 2e^{7\pi i/6}, 2e^{3\pi i/2}, 2e^{11\pi i/6}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 7 continued

Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 7 continued

(Total for Question 7 is 9 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

8. $\mathbf{A} = \begin{pmatrix} 3 & 1 & -1 \\ 1 & 1 & 1 \\ k & 3 & 6 \end{pmatrix} \quad k \neq 0$

(a) Find, in terms of k, $\mathbf{A}^{-1}$

(4)

(b) Determine, in simplest form in terms of k, the coordinates of the point where the following planes intersect.

$$3x + y - z = 3$$
$$x + y + z = 1$$
$$kx + 3y + 6z = 6$$

(3)

(a) $\det A = 3(6-3) - 1(6-k) + (-1)(3-k) = 9 - 6 + k - 3 + k = 2k$

Matrix of cofactors $= \begin{pmatrix} 3 & k-6 & 3-k \\ -9 & 18+k & k-9 \\ 2 & -4 & 2 \end{pmatrix}$

$A^{-1} = \frac{1}{\det A} \times (\text{cofactors})^T = \frac{1}{2k} \begin{pmatrix} 3 & -9 & 2 \\ k-6 & 18+k & -4 \\ 3-k & k-9 & 2 \end{pmatrix}$

(b) $\begin{pmatrix} x \\ y \\ z \end{pmatrix} = A^{-1} \begin{pmatrix} 3 \\ 1 \\ 6 \end{pmatrix} = \frac{1}{2k} \begin{pmatrix} 3 & -9 & 2 \\ k-6 & 18+k & -4 \\ 3-k & k-9 & 2 \end{pmatrix} \begin{pmatrix} 3 \\ 1 \\ 6 \end{pmatrix}$

$= \frac{1}{2k} \begin{pmatrix} 9 - 9 + 12 \\ 3k - 18 + 18 + k - 24 \\ 9 - 3k + k - 9 + 12 \end{pmatrix} = \frac{1}{2k} \begin{pmatrix} 12 \\ 4k - 24 \\ 12 - 2k \end{pmatrix}$

$x = \frac{6}{k}, \quad y = \frac{2k-12}{k}, \quad z = \frac{6-k}{k}$

Point of intersection: $\left(\frac{6}{k}, \frac{2k-12}{k}, \frac{6-k}{k} \right)$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 8 continued

Question 8 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 8 continued

(Total for Question 8 is 7 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

9.




Figure 1




Figure 2

Figure 1 shows the central vertical cross-section *ABCDEFA* of a vase together with measurements that have been taken from the vase.

The horizontal cross-section between *AB* and *FC* is a circle with diameter 4 cm.

The base of the vase *ED* is horizontal and the point *E* is vertically below *F* and the point *D* is vertically below *C*.

Using these measurements, the curve *CD* is modelled by the parametric equations

$$x = a + 3\sin 2t \qquad y = b\cos t \qquad 0 \leqslant t \leqslant \frac{\pi}{2}$$

where *a* and *b* are constants and *O* is the fixed origin, as shown in Figure 2.

(a) Determine the value of *a* and the value of *b* according to the model.

(2)

(b) Using algebraic integration and showing all your working, determine, according to the model, the volume of the vase, giving your answer to the nearest cm^3

(7)

(c) State a limitation of the model.

(1)

(a) At C, $t = 0$: $x = a + 3\sin 0 = a$ and C is 2 cm from the axis $\Rightarrow$ $a = 2$

$y = b\cos 0 = b$ and C is 7 cm above ED $\Rightarrow$ $b = 7$

(b) $V_{curve} = \pi \int x^2\,dy = \pi \int_{\pi/2}^{0} x^2 \frac{dy}{dt}\,dt, \quad \frac{dy}{dt} = -7\sin t$

$= \pi \int_{\pi/2}^{0} (2 + 3\sin 2t)^2(-7\sin t)\,dt = 7\pi \int_{0}^{\pi/2} (4 + 12\sin 2t + 9\sin^2 2t)\sin t\,dt$

Using $\sin 2t = 2\sin t \cos t$:

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 9 continued

$= 7\pi \int_0^{\pi/2} (4\sin t + 24\sin^2 t \cos t + 36\sin^3 t \cos^2 t)\, dt$

$\sin^3 t \cos^2 t = (1 - \cos^2 t)\cos^2 t \sin t = \cos^2 t \sin t - \cos^4 t \sin t$

$= 7\pi \left[-4\cos t + 8\sin^3 t - 12\cos^3 t + \frac{36}{5} \cos^5 t \right]_0^{\pi/2}$

$= 7\pi \left[(0 + 8 - 0 + 0) - (-4 + 0 - 12 + \frac{36}{5}) \right]$

$= 7\pi \left(8 + 16 - \frac{36}{5}\right) = 7\pi \times \frac{84}{5} = \frac{588}{5}\pi$

Cylinder (neck): $\pi \times 2^2 \times 4.5 = 18\pi$

Volume of vase $= \frac{588}{5}\pi + 18\pi = \frac{678}{5}\pi = 425.9999...$

$\approx 426\ cm^3$

(c) The thickness of the sides (walls) of the vase has not been considered.

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 9 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 9 continued

Question 9 continued

(Total for Question 9 is 10 marks)

TOTAL FOR PAPER IS 75 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA