Centres of mass of plane figures 2H

1 We view the square as composed of two rectangles, with the rightmost rectangle having twice the mass of the leftmost one. Now we choose coordinates with the origin at D and DC lying on the x-axis.
Then the centre of mass of the leftmost rectangle is at $(1,2)$ and the coordinates of the rightmost is at $(3,2)$ hence the centre of mass satisfies

$$3\begin{pmatrix}x\\y\end{pmatrix}=\begin{pmatrix}1\\2\end{pmatrix}+2\begin{pmatrix}3\\2\end{pmatrix}$$

Hence

$$3\begin{pmatrix}x\\y\end{pmatrix}=\begin{pmatrix}7\\6\end{pmatrix}$$

Hence

$$\begin{pmatrix}x\\y\end{pmatrix}=\begin{pmatrix}\frac{7}{3}\\2\end{pmatrix}$$

2 We choose coordinates with origin at D and DC lying on the x-axis, then we can view the lamina as composed of a 4 × 6 rectangle and a triangle, in these coordinates the centre of mass of the rectangle is $(8,3)$ and the centre of mass of the triangle can be obtained by averaging the coordinates of the vertices giving

$$\frac{1}{3}\begin{pmatrix}0\\6\end{pmatrix}+\frac{1}{3}\begin{pmatrix}6\\0\end{pmatrix}+\frac{1}{3}\begin{pmatrix}6\\6\end{pmatrix}=\begin{pmatrix}4\\4\end{pmatrix}$$


Hence the centre of mass of the lamina satisfies

$$3\begin{pmatrix}x\\y\end{pmatrix}=2\begin{pmatrix}4\\4\end{pmatrix}+\begin{pmatrix}8\\3\end{pmatrix}=\begin{pmatrix}16\\11\end{pmatrix}$$

Hence

$$\begin{pmatrix}x\\y\end{pmatrix}=\begin{pmatrix}\frac{16}{3}\\\frac{11}{3}\end{pmatrix}$$

3 When the lamina is folded it looks like this



Now using Pythagoras gives

$$x^2=6^2+(10-x)^2$$


which implies that $x=6.8$. Now we choose co-ordinates with the origin at C and EC lying on the x-axis. Now the centre of mass of BCE is

$$\frac{1}{3}\begin{pmatrix}0\\0\end{pmatrix}+\frac{1}{3}\begin{pmatrix}-3.2\\0\end{pmatrix}+\frac{1}{3}\begin{pmatrix}0\\6\end{pmatrix}=\begin{pmatrix}-1.07\\2\end{pmatrix}$$

The centre of mass of FBE is given by

$$\frac{1}{3}\begin{pmatrix}0\\6\end{pmatrix}+\frac{1}{3}\begin{pmatrix}-3.2\\0\end{pmatrix}+\frac{1}{3}\begin{pmatrix}-6.8\\6\end{pmatrix}=\begin{pmatrix}-3.33\\4\end{pmatrix}$$

Finally to find the centre of mass of the last triangle consider the following figure



We have $\sin\theta=\dfrac{3.2}{6.8}$ so that

$y=6\sin\theta=2.82$ and $x=6\cos\theta=5.29$

hence the centre of mass is given by

$$\frac{1}{3}\begin{pmatrix}0\\6\end{pmatrix}+\frac{1}{3}\begin{pmatrix}-6.8\\6\end{pmatrix}+\frac{1}{3}\begin{pmatrix}-5.29\\8.82\end{pmatrix}=\begin{pmatrix}-4.03\\6.94\end{pmatrix}$$

© Pearson Education Ltd 2018. Copying permitted for purchasing institution only. This material is not copyright free.

3 (continued)

Putting this together, the centre of mass of the lamina will satisfy

$$60\begin{pmatrix} x \\ y \end{pmatrix} = 9.6\begin{pmatrix} -1.07 \\ 2 \end{pmatrix} + 40.8\begin{pmatrix} -3.33 \\ 4 \end{pmatrix} + 9.6\begin{pmatrix} -4.03 \\ 6.94 \end{pmatrix}$$

Which gives

$$\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -3.08 \\ 4.15 \end{pmatrix}$$

Hence the angle of suspension satisfies

$$\tan\theta = \frac{5.29 - 3.08}{8.82 - 4.15}$$

Hence $\theta = 25.3°$ (3 s.f.)

4 From the diagram we can see that the centre of mass of AB is $(1,2)$, the centre of mass of BC is $(4,4)$, the centre of mass of CD is $(6,1)$ and the centre of mass of AD is $(3,-1)$ therefore the centre of mass of the system satisfies

$$11\begin{pmatrix} x \\ y \end{pmatrix} = 2\begin{pmatrix} 1 \\ 2 \end{pmatrix} + 3\begin{pmatrix} 4 \\ 4 \end{pmatrix} + \begin{pmatrix} 6 \\ 1 \end{pmatrix} + 5\begin{pmatrix} 3 \\ -1 \end{pmatrix} = \begin{pmatrix} 35 \\ 12 \end{pmatrix}$$

Hence

$$\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \frac{35}{11} \\ \frac{12}{11} \end{pmatrix}$$

5 We first note that a circular wire of twice the thickness has a cross-sectional area of four times the area and so four times the mass.

From the diagram the centre of mass of AB is $(1,4)$ and its length is $\sqrt{4^2+4^2} = 4\sqrt{2}$, the centre of mass of BC is $(4,2)$ and its length is $\sqrt{2^2+8^2} = 2\sqrt{17}$ and the centre of mass of AC is $(2,0)$ and its length is $\sqrt{4^2+6^2} = 2\sqrt{13}$ and hence the centre of mass of the system satisfies

$$\left(4\sqrt{2} + 8\sqrt{13} + 8\sqrt{17}\right)\begin{pmatrix} x \\ y \end{pmatrix}$$

$$= 4\sqrt{2}\begin{pmatrix} 1 \\ 4 \end{pmatrix} + 8\sqrt{17}\begin{pmatrix} 4 \\ 2 \end{pmatrix} + 8\sqrt{13}\begin{pmatrix} 2 \\ 0 \end{pmatrix}$$

Hence

$$\left(4\sqrt{2} + 8\sqrt{13} + 8\sqrt{17}\right)\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 4\sqrt{2} + 16\sqrt{13} + 32\sqrt{17} \\ 16\sqrt{2} + 16\sqrt{17} \end{pmatrix}$$

So

$$\begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{4\sqrt{2} + 8\sqrt{13} + 8\sqrt{17}}\begin{pmatrix} 4\sqrt{2} + 16\sqrt{13} + 32\sqrt{17} \\ 16\sqrt{2} + 16\sqrt{17} \end{pmatrix}$$

$= (2.89, 1.31)$ (3 s.f.)

© Pearson Education Ltd 2018. Copying permitted for purchasing institution only. This material is not copyright free.

6 Firstly we can see that the mass of BC is $0.5M$ since the length of BC is half the length of AD.

Now the centre of mass of AD is $(5,1)$ the centre of mass of BC is $(5,5)$, the centre of mass of AB is $(2,3)$ and the centre of mass of CD is $(8,3)$.

Hence the centre of mass of the system satisfies

$$2.5\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 5 \\ 1 \end{pmatrix} + 0.5\begin{pmatrix} 5 \\ 5 \end{pmatrix} + 0.5\begin{pmatrix} 2 \\ 3 \end{pmatrix} + 0.5\begin{pmatrix} 8 \\ 3 \end{pmatrix}$$

So

$$2.5\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 12.5 \\ 6.5 \end{pmatrix}$$

So

$$\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 5 \\ 2.6 \end{pmatrix}$$

Now B has coordinates (3,5).

Hence the angle with the vertical satisfies

$$\tan\theta = \frac{2.4}{2}$$

Hence $\theta = 50.2^\circ$

7 a We choose coordinates so that the origin is at O and AC lies on the x-axis then nothing that the length of OA is $\sqrt{1.3^2 - 0.5^2} = 1.2$ the coordinates of the centre of mass of the triangle are given by

$$\frac{1}{3}\begin{pmatrix} 0 \\ 0.5 \end{pmatrix} + \frac{1}{3}\begin{pmatrix} 0 \\ -0.5 \end{pmatrix} + \frac{1}{3}\begin{pmatrix} -1.2 \\ 0 \end{pmatrix} = \begin{pmatrix} -0.4 \\ 0 \end{pmatrix}$$

The centre of mass of the semi-circular lamina is given by

$$\begin{pmatrix} \frac{2\times 0.5}{3} & \frac{\sin\frac{\pi}{2}}{\frac{\pi}{2}} \\ 0 \end{pmatrix} = \begin{pmatrix} \frac{2}{3\pi} \\ 0 \end{pmatrix}$$

Hence the centre of mass of the lamina satisfies

$$20\begin{pmatrix} x \\ y \end{pmatrix} = 4\begin{pmatrix} -0.4 \\ 0 \end{pmatrix} + 16\begin{pmatrix} \frac{2}{3\pi} \\ 0 \end{pmatrix} = \begin{pmatrix} \frac{32}{3\pi} - 1.6 \\ 0 \end{pmatrix}$$

Hence

$$\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \frac{8}{15\pi} - 0.08 \\ 0 \end{pmatrix}$$

So the distance of the centre of mass from BD is simply

$$\frac{8}{15\pi} - 0.08 = 0.09$$

b The other wire should be attached at C so that the centre of mass lies between the two wires.

c Let T_1, T_2 be the tension in the wires at B and C respectively, taking moments about B gives

$$0.5T_2 = 0.09\times 20g$$

So

$$T_2 = 3.6g$$

Taking moments about C gives

$$0.5T_1 = 0.41\times 20g$$

So

$$T_1 = 16.4g$$

© Pearson Education Ltd 2018. Copying permitted for purchasing institution only. This material is not copyright free.

8 a We need to find the coordinates of the centre of mass of the lamina, we choose coordinates such that the origin is the midpoint of BD and BD lies on the x-axis, by symmetry the x component of the centre of mass is zero. The centre of mass of the square is then given by $(0,-4)$.

The centre of mass of the triangle is given by

$$\frac{1}{3}\begin{pmatrix}4\\0\end{pmatrix}+\frac{1}{3}\begin{pmatrix}-4\\0\end{pmatrix}+\frac{1}{3}\begin{pmatrix}0\\3\end{pmatrix}=\begin{pmatrix}0\\1\end{pmatrix}$$

So the centre of mass of the lamina satisfies

$$(60\times12+20\times64)\begin{pmatrix}x\\y\end{pmatrix}=60\times12\begin{pmatrix}0\\1\end{pmatrix}$$

$$+20\times64\begin{pmatrix}0\\-4\end{pmatrix}$$

Simplifying gives

$y=-2.2$

$x=0$

So the centre of mass is $(0,-2.2)$

And coordinates of B are clearly $(-4,0)$

So the angle that AB makes with the vertical satisfies

$$\tan\theta=4\times\frac{1}{2.2}=\frac{4}{2.2}$$

So $\theta=61.2^\circ$

If we add a mass at the point A the new centre of mass will satisfy

$$(720+1280+500)\begin{pmatrix}x\\y\end{pmatrix}=720\begin{pmatrix}0\\1\end{pmatrix}$$

$$+1280\begin{pmatrix}0\\-4\end{pmatrix}+500\begin{pmatrix}-4\\-8\end{pmatrix}$$

Which simplifies to

$$\begin{pmatrix}x\\y\end{pmatrix}=\begin{pmatrix}-0.8\\-3.36\end{pmatrix}$$

Hence the angle satisfies.

b $\tan\theta=\dfrac{3.2}{3.36}$

So

$\theta=43.6^\circ$

Challenge

a Let the width of the rectangle be x then since we fold twice, the angle of the triangle at G is 45° hence the height of the rectangle is

$$y=\frac{\frac{x}{2}}{\tan22.5^\circ}=\frac{x}{2(\sqrt{2}-1)}=\frac{\sqrt{2}+1}{2}x$$

Which is the desired ratio.

b Split the shape into areas of ×4, ×3, ×2, ×1 layers of paper.

Using com of triangle $=\left(0,\dfrac{y_1+y_2+y_3}{3}\right)$

Find:

com of ×4 layer lies $0.667x$ below G

com of ×3 layer lies $1.138x$ below G

com of ×2 layer lies$1.61x$ below G

com of ×1 layer triangle lies $2.081x$ below G

area ×4 = $0.414x \times x = 0.414x^2$

area ×3 = $0.414x \times 0.414x = 0.171x^2$

area ×2 = = $0.585x \times 1.414x = 0.827x^2$

area ×1 triangle = $x \times x = x^2$

Let y be the position of the centre of mass below G then

$4.828x^2y = 4(0.667x \times 0.414x^2)$
$+ 3(1.138x \times 0.171x^2) + 2(1.61x \times 0.827x^2)$
$+ (2.081x \times x^2)$

$y = 1.33x$

so the centre of mass lies $1.33x$ vertically below G

© Pearson Education Ltd 2018. Copying permitted for purchasing institution only. This material is not copyright free.