Vectors 11E

1 $\overrightarrow{XY} = \overrightarrow{XW} + \overrightarrow{WY} = \mathbf{b} - \mathbf{a}$

$\overrightarrow{YZ} = \overrightarrow{YW} + \overrightarrow{WZ} = \mathbf{c} - \mathbf{b}$

Since $\overrightarrow{XY} = \overrightarrow{YZ}$:

$\mathbf{b} - \mathbf{a} = \mathbf{c} - \mathbf{b}$

$\mathbf{b} + \mathbf{b} = \mathbf{a} + \mathbf{c}$

$\mathbf{a} + \mathbf{c} = 2\mathbf{b}$

2 a i $\overrightarrow{OB} = 2\overrightarrow{OR}$

$= 2\mathbf{r}$

ii $\overrightarrow{PQ} = \overrightarrow{PO} + \overrightarrow{OQ}$ (addition of vectors)

$= -\overrightarrow{OP} + \overrightarrow{OQ}$

$\overrightarrow{OQ} = \overrightarrow{OA} + \overrightarrow{AQ}$ (addition of vectors)

$\overrightarrow{AQ} = \frac{1}{2}\overrightarrow{AB}$

$\overrightarrow{AB} = \overrightarrow{AO} + \overrightarrow{OB}$ (addition of vectors)

$= -\overrightarrow{OA} + \overrightarrow{OB}$

$= -2\mathbf{p} + 2\mathbf{r}$

$\therefore \overrightarrow{AQ} = \frac{1}{2}(-2\mathbf{p} + 2\mathbf{r})$

$= -\mathbf{p} + \mathbf{r}$

$\therefore \overrightarrow{OQ} = 2\mathbf{p} + (-\mathbf{p} + \mathbf{r})$

$= \mathbf{p} + \mathbf{r}$

$\therefore \overrightarrow{PQ} = -\mathbf{p} + (\mathbf{p} + \mathbf{r})$

$= \mathbf{r}$

b $\overrightarrow{OB} = 2\mathbf{r}$ and $\overrightarrow{PQ} = \mathbf{r}$

$\Rightarrow \overrightarrow{OB}$ and $\overrightarrow{PQ}$ are parallel.

$\Rightarrow \angle AOB = \angle APQ$ and $\angle ABO = \angle AQP$ (corresponding angles, parallel lines).

Angle A is common to both triangles.

$\Rightarrow$ $\triangle PAQ$ and $\triangle OAB$ are similar (three equal angles)

3 a M divides OA in the ratio $2:1$.

$\Rightarrow \overrightarrow{OM} = \frac{2}{3}\mathbf{a}$

Using vector addition:

$\overrightarrow{ON} = \overrightarrow{OA} + \overrightarrow{AN}$

$\overrightarrow{AN} = \lambda\overrightarrow{AB}$ (N lies on AB, so $\overrightarrow{AN} = \lambda\overrightarrow{AB}$)

$\overrightarrow{AB} = \mathbf{b} - \mathbf{a}$

$\overrightarrow{ON} = \mathbf{a} + \lambda(\mathbf{b} - \mathbf{a})$

3 b $\overrightarrow{ON} = \overrightarrow{OM} + \overrightarrow{MN}$

$= \overrightarrow{OM} + \mu\overrightarrow{OB}$ ($\overrightarrow{MN}$ is parallel to $\overrightarrow{OB}$)

$= \frac{2}{3}\mathbf{a} + \mu\mathbf{b}$

But $\overrightarrow{ON} = \mathbf{a} + \lambda(\mathbf{b} - \mathbf{a})$

So:

$\mathbf{a} + \lambda(\mathbf{b} - \mathbf{a}) = \frac{2}{3}\mathbf{a} + \mu\mathbf{b}$

$\mathbf{a}(1 - \lambda) + \lambda\mathbf{b} = \frac{2}{3}\mathbf{a} + \mu\mathbf{b}$

$\Rightarrow$ (comparing coefficients of **a** and **b**):

$1 - \lambda = \frac{2}{3}$ and $\lambda = \mu$

so $\lambda = \mu = \frac{1}{3}$ and $\overrightarrow{ON} = \frac{2}{3}\mathbf{a} + \frac{1}{3}\mathbf{b}$

$\overrightarrow{AN} = \frac{1}{3}(\mathbf{b} - \mathbf{a})$

$\Rightarrow \overrightarrow{NB} = \frac{2}{3}(\mathbf{b} - \mathbf{a})$

$\Rightarrow AN : NB = 1:2$

4 M is the mid-point of OA, so:

$\overrightarrow{OM} = \frac{1}{2}\overrightarrow{OA}$

$= \frac{1}{2}\mathbf{a}$

Using vector addition:

$\overrightarrow{MQ} = \overrightarrow{MA} + \overrightarrow{AB} + \overrightarrow{BQ}$

$= \overrightarrow{MA} + \overrightarrow{AB} + \frac{1}{4}\overrightarrow{BC}$

$= \frac{1}{2}\mathbf{a} + \mathbf{c} - \frac{1}{4}\mathbf{a}$

$= \frac{1}{4}\mathbf{a} + \mathbf{c}$

and:

$\overrightarrow{AC} = \overrightarrow{AO} + \overrightarrow{OC}$

$= -\mathbf{a} + \mathbf{c}$

$= \mathbf{c} - \mathbf{a}$

P lies on AC and MQ, so:

$\overrightarrow{OP} = \overrightarrow{OM} + \lambda\overrightarrow{MQ}$

$= \frac{1}{2}\mathbf{a} + \lambda(\frac{1}{4}\mathbf{a} + \mathbf{c})$

and $\overrightarrow{OP} = \overrightarrow{OA} + \mu\overrightarrow{AC}$

$= \mathbf{a} + \mu(\mathbf{c} - \mathbf{a})$

Comparing coefficients of **a** and **c**:

$\frac{1}{2} + \frac{1}{4}\lambda = 1 - \mu$ and $\lambda = \mu$

$\Rightarrow \frac{5}{4}\lambda = \frac{1}{2}$

$\lambda = \mu = \frac{2}{5}$

$\overrightarrow{OP} = \frac{3}{5}\mathbf{a} + \frac{2}{5}\mathbf{c}$

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

5 a $\overrightarrow{AB} = -\overrightarrow{OA} + \overrightarrow{OB}$

$= -\begin{pmatrix} 5 \\ 8 \end{pmatrix} + \begin{pmatrix} 4 \\ 3 \end{pmatrix}$

$= \begin{pmatrix} -1 \\ -5 \end{pmatrix}$

$|\overrightarrow{AB}| = \sqrt{(-1)^2 + (-5)^2}$

$= \sqrt{26}$

b $\overrightarrow{AC} = -\overrightarrow{OA} + \overrightarrow{OC}$

$= -\begin{pmatrix} 5 \\ 8 \end{pmatrix} + \begin{pmatrix} 7 \\ 6 \end{pmatrix}$

$= \begin{pmatrix} 2 \\ -2 \end{pmatrix}$

$|\overrightarrow{AC}| = \sqrt{2^2 + (-2)^2}$

$= \sqrt{8}$

$= 2\sqrt{2}$

c $\overrightarrow{BC} = -\overrightarrow{OB} + \overrightarrow{OC}$

$= -\begin{pmatrix} 4 \\ 3 \end{pmatrix} + \begin{pmatrix} 7 \\ 6 \end{pmatrix}$

$= \begin{pmatrix} 3 \\ 3 \end{pmatrix}$

$|\overrightarrow{BC}| = \sqrt{3^2 + 3^2}$

$= \sqrt{18}$

$= 3\sqrt{2}$

d Using the cosine rule:

$\cos A = \dfrac{b^2 + c^2 - a^2}{2bc}$

$\cos A = \dfrac{(2\sqrt{2})^2 + (\sqrt{26})^2 - (3\sqrt{2})^2}{2(2\sqrt{2})(\sqrt{26})}$

$\cos A = \dfrac{8 + 26 - 18}{4\sqrt{52}}$

$\cos A = \dfrac{16}{8\sqrt{13}}$

$\cos A = \dfrac{2}{\sqrt{13}}$

5 d $A = 56.3...°$

Using the sine rule:

$\dfrac{\sin B}{b} = \dfrac{\sin A}{a}$

$\dfrac{\sin B}{2\sqrt{2}} = \dfrac{\sin 56.3°}{3\sqrt{2}}$

$\sin B = \dfrac{2\sqrt{2}\sin 56.3°}{3\sqrt{2}}$

$B = 33.68...°$

$C = 180° - 56° - 34° = 90°$

The angles are 56°, 34° and 90°.

6 a $\overrightarrow{OS} = \overrightarrow{OP} + \overrightarrow{PR} + \overrightarrow{RS}$

$\overrightarrow{OP} = \mathbf{a}$

$\overrightarrow{PR} = \frac{1}{3}(-\mathbf{a} + \mathbf{b})$

$\overrightarrow{RS} = 2\overrightarrow{OR} = 2\left(\overrightarrow{OP} + \overrightarrow{PR}\right)$

$= 2\left(\mathbf{a} + \frac{1}{3}\left(-\mathbf{a} + \mathbf{b}\right)\right)$

$= 2\mathbf{a} - \frac{2}{3}\mathbf{a} + \frac{2}{3}\mathbf{b}$

$= \frac{4}{3}\mathbf{a} + \frac{2}{3}\mathbf{b}$

So $\overrightarrow{OS} = \mathbf{a} + \frac{1}{3}\left(-\mathbf{a} + \mathbf{b}\right) + \frac{4}{3}\mathbf{a} + \frac{2}{3}\mathbf{b}$

$= 2\mathbf{a} + \mathbf{b}$

b $\overrightarrow{TP} = \overrightarrow{TO} + \overrightarrow{OP}$

$= \mathbf{b} + \mathbf{a} = \mathbf{a} + \mathbf{b}$

$\overrightarrow{PS} = \overrightarrow{PR} + \overrightarrow{RS}$

$= \frac{1}{3}(-\mathbf{a} + \mathbf{b}) + \frac{4}{3}\mathbf{a} + \frac{2}{3}\mathbf{b} = \mathbf{a} + \mathbf{b}$

$\overrightarrow{TP}$ is parallel and equal to $\overrightarrow{PS}$ and point P is common to both lines, so T, P and S lie on a straight line.

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

Challenge

$\overrightarrow{PX} = j\overrightarrow{PR}$

$\overrightarrow{PR} = \overrightarrow{PO} + \overrightarrow{OR}$

$= -\mathbf{a} + \mathbf{b}$

$\overrightarrow{PX} = j(-\mathbf{a} + \mathbf{b})$

$= -j\mathbf{a} + j\mathbf{b}$

$\overrightarrow{PX} = \overrightarrow{PO} + \overrightarrow{OX}$

$\overrightarrow{OX} = k\overrightarrow{ON}$

$\overrightarrow{ON} = \mathbf{a} + \frac{1}{2}\mathbf{b}$

$\overrightarrow{PX} = -\mathbf{a} + k\left(\mathbf{a} + \frac{1}{2}\mathbf{b}\right)$

$= (k-1)\mathbf{a} + \frac{1}{2}k\mathbf{b}$

As $\overrightarrow{PX} = -j\mathbf{a} + j\mathbf{b}$

and $\overrightarrow{PX} = (k-1)\mathbf{a} + \frac{1}{2}k\mathbf{b}$

$-j\mathbf{a} + j\mathbf{b} = (k-1)\mathbf{a} + \frac{1}{2}k\mathbf{b}$

The coefficients of **a** and **b** must be the same, so $k - 1 = -j$ and $\frac{1}{2}k = j$.

Solving the equation simultaneously and using substitution:

$k - 1 = -\frac{1}{2}k$

$k = \frac{2}{3}$

$j = \frac{1}{3}$

$\overrightarrow{PX} = -\frac{1}{3}\mathbf{a} + \frac{1}{3}\mathbf{b}$

As *OPQR* is a parallelogram, $\overrightarrow{YR} = \overrightarrow{PX}$.

Therefore $\overrightarrow{PX} = \overrightarrow{XY} = \overrightarrow{YR}$, so the line *PR* is divided into three equal parts.

Therefore, the lines *ON* and *OM* divide the diagonal *PR* into three equal parts.

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.