Integration 11B

1 a $\int \sin(2x+1)\mathrm{d}x = -\frac{1}{2}\cos(2x+1)+c$

b $\int 3\mathrm{e}^{2x}\mathrm{d}x = \frac{3}{2}\mathrm{e}^{2x}+c$

c $\int 4\mathrm{e}^{x+5}\mathrm{d}x = 4\mathrm{e}^{x+5}+c$

d $\int \cos(1-2x)\mathrm{d}x = -\frac{1}{2}\sin(1-2x)+c$

OR Let $y = \sin(1-2x)$

then $\frac{\mathrm{d}y}{\mathrm{d}x} = \cos(1-2x)\times(-2)$ (by chain rule)

$\therefore \int \cos(1-2x)\mathrm{d}x = -\frac{1}{2}\sin(1-2x)+c$

e $\int \operatorname{cosec}^2 3x\,\mathrm{d}x = -\frac{1}{3}\cot 3x+c$

f $\int \sec 4x\tan 4x\,\mathrm{d}x = \frac{1}{4}\sec 4x+c$

g $\int 3\sin\left(\frac{1}{2}x+1\right)\mathrm{d}x = -6\cos\left(\frac{1}{2}x+1\right)+c$

h $\int \sec^2(2-x)\mathrm{d}x = -\tan(2-x)+c$

OR Let $y = \tan(2-x)$

then $\frac{\mathrm{d}y}{\mathrm{d}x} = \sec^2(2-x)\times(-1)$ (by chain rule)

$\therefore \int \sec^2(2-x)\mathrm{d}x = -\tan(2-x)+c$

i $\int \operatorname{cosec} 2x\cot 2x\,\mathrm{d}x = -\frac{1}{2}\operatorname{cosec} 2x+c$

j $\int(\cos 3x-\sin 3x)\mathrm{d}x$

$= \frac{1}{3}\sin 3x+\frac{1}{3}\cos 3x+c$

$= \frac{1}{3}(\sin 3x+\cos 3x)+c$

2 a $\int(\mathrm{e}^{2x}-\frac{1}{2}\sin(2x-1))\mathrm{d}x$

$= \frac{1}{2}\mathrm{e}^{2x}+\frac{1}{4}\cos(2x-1)+c$

b $\int(\mathrm{e}^x+1)^2\mathrm{d}x$

$= \int(\mathrm{e}^{2x}+2\mathrm{e}^x+1)\mathrm{d}x$

$= \frac{1}{2}\mathrm{e}^{2x}+2\mathrm{e}^x+x+c$

c $\int \sec^2 2x(1+\sin 2x)\mathrm{d}x$

$= \int(\sec^2 2x+\sec^2 2x\sin 2x)\mathrm{d}x$

$= \int(\sec^2 2x+\sec 2x\tan 2x)\mathrm{d}x$

$= \frac{1}{2}\tan 2x+\frac{1}{2}\sec 2x+c$

d $\int\frac{3-2\cos\left(\frac{1}{2}x\right)}{\sin^2\left(\frac{1}{2}x\right)}\mathrm{d}x$

$= \int\left(3\operatorname{cosec}^2\frac{1}{2}x-2\operatorname{cosec}\frac{1}{2}x\cot\frac{1}{2}x\right)\mathrm{d}x$

$= -6\cot\left(\frac{1}{2}x\right)+4\operatorname{cosec}\left(\frac{1}{2}x\right)+c$

2 e $\int(\mathrm{e}^{3-x}+\sin(3-x)+\cos(3-x))\mathrm{d}x$

$= -\mathrm{e}^{3-x}+\cos(3-x)-\sin(3-x)+c$

Note: extra minus signs from $-x$ terms and chain rule.

3 a $\int\frac{1}{2x+1}\mathrm{d}x = \frac{1}{2}\ln|2x+1|+c$

b $\int\frac{1}{(2x+1)^2}\mathrm{d}x$

$= \int(2x+1)^{-2}\mathrm{d}x$

$= \frac{(2x+1)^{-1}}{-1}\times\frac{1}{2}+c$

$= -\frac{1}{2(2x+1)}+c$

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

c $\int (2x+1)^2 \,\mathrm{d}x$

$= \dfrac{(2x+1)^3}{3} \times \dfrac{1}{2} + c$

$= \dfrac{(2x+1)^3}{6} + c$

d $\int \dfrac{3}{4x-1} \,\mathrm{d}x = \dfrac{3}{4} \ln|4x-1| + c$

e $\int \dfrac{3}{1-4x} \,\mathrm{d}x$

$= -\int \dfrac{3}{4x-1} \,\mathrm{d}x$

$= -\dfrac{3}{4} \ln|4x-1| + c$

OR Let $y = \ln|1-4x|$

then $\dfrac{\mathrm{d}y}{\mathrm{d}x} = \dfrac{1}{1-4x} \times (-4)$ (by chain rule)

$\therefore \int \dfrac{3}{1-4x} \,\mathrm{d}x = -\dfrac{3}{4} \ln|1-4x| + c$

Note: $\ln|1-4x| = \ln|4x-1|$ because of $|\ |$ sign.

f $\int \dfrac{3}{(1-4x)^2} \,\mathrm{d}x$

$= \int 3(1-4x)^{-2} \,\mathrm{d}x$

$= \dfrac{3}{-4} \times \dfrac{(1-4x)^{-1}}{-1} + c$

$= \dfrac{3}{4(1-4x)} + c$

g $\int (3x+2)^5 \,\mathrm{d}x = \dfrac{(3x+2)^6}{18} + c$

h $\int \dfrac{3}{(1-2x)^3} \,\mathrm{d}x = \dfrac{3}{-2} \times \dfrac{(1-2x)^{-2}}{-2} + c$

$= \dfrac{3}{4(1-2x)^2} + c$

OR Let $y = (1-2x)^{-2}$

then $\dfrac{\mathrm{d}y}{\mathrm{d}x} = -2(1-2x)^{-3} \times (-2)$
(by chain rule)

$\therefore \int \dfrac{3}{(1-2x)^3} \,\mathrm{d}x = \dfrac{3}{4}(1-2x)^{-2} + c$

4 a $\int \left(3\sin(2x+1) + \dfrac{4}{2x+1}\right) \mathrm{d}x$

$= -\dfrac{3}{2}\cos(2x+1) + \dfrac{4}{2} \ln|2x+1| + c$

$= -\dfrac{3}{2}\cos(2x+1) + 2\ln|2x+1| + c$

b $\int \left(\mathrm{e}^{5x} + (1-x)^5\right) \mathrm{d}x = \int \mathrm{e}^{5x} \mathrm{d}x + \int (1-x)^5 \mathrm{d}x$

$= \dfrac{1}{5}\mathrm{e}^{5x} - \dfrac{1}{6}(1-x)^6 + c$

OR Let $y = (1-x)^6$

then $\dfrac{\mathrm{d}y}{\mathrm{d}x} = 6(1-x)^5 \times (-1)$ (by chain rule)

$\therefore \int (1-x)^5 \,\mathrm{d}x = -\dfrac{1}{6}(1-x)^6 + c$

c $\int \left(\dfrac{1}{\sin^2 2x} + \dfrac{1}{1+2x} + \dfrac{1}{(1+2x)^2}\right) \mathrm{d}x$

$= \int \left(\operatorname{cosec}^2 2x + \dfrac{1}{1+2x} + (1+2x)^{-2}\right) \mathrm{d}x$

$= -\dfrac{1}{2}\cot 2x + \dfrac{1}{2}\ln|1+2x|$

$+ \dfrac{(1+2x)^{-1}}{-1} \times \dfrac{1}{2} + c$

$= -\dfrac{1}{2}\cot 2x + \dfrac{1}{2}\ln|1+2x| - \dfrac{1}{2(1+2x)} + c$

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

d $\int\left((3x+2)^2+\dfrac{1}{(3x+2)^2}\right)\mathrm{d}x$

$=\int\left((3x+2)^2+(3x+2)^{-2}\right)\mathrm{d}x$

$=\dfrac{(3x+2)^3}{9}-\dfrac{(3x+2)^{-1}}{3}+c$

$=\dfrac{(3x+2)^3}{9}-\dfrac{1}{3(3x+2)}+c$

5 a $\int_{\frac{\pi}{4}}^{\frac{3\pi}{4}}\cos(\pi-2x)\mathrm{d}x=\left[-\dfrac{1}{2}\sin(\pi-2x)\right]_{\frac{\pi}{4}}^{\frac{3\pi}{4}}$

$=\left(-\dfrac{1}{2}\sin\left(-\dfrac{\pi}{2}\right)\right)-\left(-\dfrac{1}{2}\sin\dfrac{\pi}{2}\right)$

$=\dfrac{1}{2}+\dfrac{1}{2}=1$

b $\int_{\frac{1}{2}}^{1}\dfrac{12}{(3-2x)^4}\mathrm{d}x$

Consider $y=\dfrac{1}{(3-2x)^3}$

$\dfrac{\mathrm{d}y}{\mathrm{d}x}=\dfrac{6}{(3-2x)^4}$

So $\int_{\frac{1}{2}}^{1}\dfrac{12}{(3-2x)^4}\mathrm{d}x=\left[\dfrac{2}{(3-2x)^3}\right]_{\frac{1}{2}}^{1}$

$=2-\dfrac{1}{4}=\dfrac{7}{4}$

c $\int_{\frac{2\pi}{9}}^{\frac{5\pi}{18}}\sec^2(\pi-3x)\mathrm{d}x=\left[-\dfrac{1}{3}\tan(\pi-3x)\right]_{\frac{2\pi}{9}}^{\frac{5\pi}{18}}$

$=\left(-\dfrac{1}{3}\tan\left(\pi-\dfrac{15\pi}{18}\right)\right)-\left(-\dfrac{1}{3}\tan\left(\pi-\dfrac{6\pi}{9}\right)\right)$

$=\left(-\dfrac{1}{3}\tan\dfrac{\pi}{6}\right)-\left(-\dfrac{1}{3}\tan\dfrac{\pi}{3}\right)$

$=\left(-\dfrac{1}{3}\times\dfrac{1}{\sqrt{3}}\right)-\left(-\dfrac{1}{3}\times\sqrt{3}\right)$

$=-\dfrac{\sqrt{3}}{9}+\dfrac{\sqrt{3}}{3}=\dfrac{2\sqrt{3}}{9}$

d $\int_2^3\dfrac{5}{7-2x}\mathrm{d}x=\left[-\dfrac{5}{2}\ln|7-2x|\right]_2^3$

$=\left(-\dfrac{5}{2}\ln 1\right)-\left(-\dfrac{5}{2}\ln 3\right)$

$=\dfrac{5}{2}\ln 3$

6 $\int_3^b(2x-6)^2\mathrm{d}x=\int_3^b(4x^2-24x+36)\mathrm{d}x$

$\left[\dfrac{4x^3}{3}-12x^2+36x\right]_3^b=36$

$\left(\dfrac{4b^3}{3}-12b^2+36b\right)-(36-108+108)=36$

$\dfrac{4b^3}{3}-12b^2+36b-72=0$

$b^3-9b^2+27b-54=0$

$(b-6)(b^2-3b+9)=0$

$b=6$ since $b^2-3b+9>0$.

7 $\int_{\mathrm{e}^2}^{\mathrm{e}^8}\dfrac{1}{kx}\mathrm{d}x=\left[\dfrac{1}{k}\ln x\right]_{\mathrm{e}^2}^{\mathrm{e}^8}=\dfrac{1}{4}$

$\dfrac{8}{k}-\dfrac{2}{k}=\dfrac{1}{4}$

$k=32-8=24$

8 $\int_{\frac{\pi}{4k}}^{\frac{\pi}{3k}}(1-\pi\sin kx)\mathrm{d}x=\left[x+\dfrac{\pi}{k}\cos kx\right]_{\frac{\pi}{4k}}^{\frac{\pi}{3k}}$

$=\left(\dfrac{\pi}{3k}+\dfrac{\pi}{k}\cos\dfrac{\pi}{3}\right)-\left(\dfrac{\pi}{4k}+\dfrac{\pi}{k}\cos\dfrac{\pi}{4}\right)$

$=\left(\dfrac{\pi}{3k}+\dfrac{\pi}{2k}\right)-\left(\dfrac{\pi}{4k}+\dfrac{\pi}{\sqrt{2}k}\right)$

$=\dfrac{\pi}{k}\left(\dfrac{1}{3}+\dfrac{1}{2}\right)-\left(\dfrac{1}{4}+\dfrac{1}{\sqrt{2}}\right)=\dfrac{\pi}{k}\left(\dfrac{7}{12}-\dfrac{\sqrt{2}}{2}\right)$

$\dfrac{\pi}{k}\left(\dfrac{7}{12}-\dfrac{\sqrt{2}}{2}\right)=\pi(7-6\sqrt{2})$

$\dfrac{\pi}{k}\left(\dfrac{7-6\sqrt{2}}{12}\right)=\pi(7-6\sqrt{2})$

$k=\dfrac{1}{12}$

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

Challenge

$$\int_5^{11} \frac{1}{ax+b}\,\mathrm{d}x = \left[\frac{1}{a}\ln|ax+b| + \frac{1}{a}\ln k\right]_5^{11}$$

where $\frac{1}{a}\ln k$ is a constant

$$= \frac{1}{a}\left[\ln k|ax+b|\right]_5^{11}$$

$$= \frac{1}{a}(\ln k|11a+b| - \ln k|5a+b|)$$

$$= \frac{1}{a}(\ln k|11a+b| - k\ln|5a+b|)$$

So $\ln k|11a+b| - \ln k|5a+b| = \ln\left(\frac{41}{17}\right)$

$$\ln\left|\frac{11a+b}{5a+b}\right| = \ln\left(\frac{41}{17}\right)$$

$$\frac{11a+b}{5a+b} = \pm\frac{41}{17}$$

Case 1:

$$\frac{11a+b}{5a+b} = \frac{41}{17}$$

$$187a+17b = 205a+41b$$

$$18a = -24b$$

$$3a = -4b$$

So a must be a multiple of 4 between 0 and 10.

$a = 4 \Rightarrow b = -3$

$a = 8 \Rightarrow b = -6$

Case 2:

$$\frac{11a+b}{5a+b} = -\frac{41}{17}$$

$$187a+17b = -205a-41b$$

$$392a = -58b$$

$$b = -\frac{196}{29}a$$

But this cannot be an integer, since $a < 29$, so case 2 gives no possible solutions.

Therefore the only two possible solutions are $a = 4$, $b = -3$ and $a = 8$, $b = -6$.

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.