Functions and graphs 2D

1 a i $y \in \mathbb{R}$

ii Let $y = f(x)$

$y = 2x + 3$


$x = \dfrac{y-3}{2}$

$f^{-1}(x) = \dfrac{x-3}{2}$

iii The domain of $f^{-1}(x)$ is $x \in \mathbb{R}$

The range of $f^{-1}(x)$ is $y \in \mathbb{R}$

iv



b i $y \in \mathbb{R}$

ii Let $y = f(x)$

$y = \dfrac{x+5}{2}$


$x = 2y - 5$

$f^{-1}(x) = 2x - 5$

iii The domain of $f^{-1}(x)$ is $x \in \mathbb{R}$

The range of $f^{-1}(x)$ is $y \in \mathbb{R}$

iv



c i $y \in \mathbb{R}$

ii Let $y = f(x)$

$y = 4 - 3x$


$x = \dfrac{4-y}{3}$

$f^{-1}(x) = \dfrac{4-x}{3}$

iii The domain of $f^{-1}(x)$ is $x \in \mathbb{R}$

The range of $f^{-1}(x)$ is $y \in \mathbb{R}$

iv



d i $y \in \mathbb{R}$

ii Let $y = f(x)$

$y = x^3 - 7$


$x = \sqrt[3]{y+7}$

$f^{-1}(x) = \sqrt[3]{x+7}$

iii The domain of $f^{-1}(x)$ is $x \in \mathbb{R}$

The range of $f^{-1}(x)$ is $y \in \mathbb{R}$

iv



© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

2 a Range of f is $f(x) \in \mathbb{R}$

Let $y = f(x)$

$y = 10 - x$

$x = 10 - y$

$f^{-1}(x) = 10 - x, \ \{x \in \mathbb{R}\}$

b Range of f is $f(x) \in \mathbb{R}$

Let $y = g(x)$

$y = \frac{x}{5}$

$x = 5y$

$g^{-1}(x) = 5x, \ \{x \in \mathbb{R}\}$

c Range of f is $f(x) \neq 0$

Let $y = h(x)$

$y = \frac{3}{x}$

$x = \frac{3}{y}$

$h^{-1}(x) = \frac{3}{x}, \ \{x \neq 0\}$

d Range of f is $f(x) \in \mathbb{R}$


Let $y = k(x)$

$y = x - 8$

$x = y + 8$

$k^{-1}(x) = y + 8, \ \{x \in \mathbb{R}\}$

3



$g: x \mapsto 4 - x, \ \{x \in \mathbb{R}, \ x > 0\}$

g has range $\{g(x) \in \mathbb{R}, \ g(x) < 4\}$

The inverse function is $g^{-1}(x) = 4 - x$

Now {Range g} = {Domain g^{-1}}

and {Domain g} = {Range g}

Hence, $g^{-1}(x) = 4 - x, \ \{x \in \mathbb{R}, x < 4\}$

Although $g(x)$ and $g^{-1}(x)$ have identical equations, their domains and hence ranges are different, and so are not identical.

4 a i Maximum value of g when $x = \frac{1}{3}$

Hence $\left\{g(x) \in \mathbb{R}, \ 0 < g(x) \leq \frac{1}{3}\right\}$

ii $g^{-1}(x) = \frac{1}{x}$


iii Domain g^{-1} = Range g

$\Rightarrow$ Domain $g^{-1}: \left\{x \in \mathbb{R}, 0 < x \leq \frac{1}{3}\right\}$

Range g^{-1} = Domain g

$\Rightarrow$ Range $g^{-1}(x): \left\{g^{-1}(x) \in \mathbb{R}, \ g^{-1}(x) \geq 3\right\}$

iv




© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

4 b i Minimum value of $g(x) = -1$
when $x = 0$
Hence $\{g(x) \in \mathbb{R},\ g(x) \geq -1\}$

ii Letting $y = 2x - 1 \Rightarrow x = \dfrac{y+1}{2}$
Hence $g^{-1}(x) = \dfrac{x+1}{2}$

iii Domain g^{-1} = Range g
$\Rightarrow$ Domain $g^{-1}: \{x \in \mathbb{R},\ x \geq -1\}$
Range g^{-1} = Domain g
$\Rightarrow$ Range $g^{-1}(x): \begin{Bmatrix} g^{-1}(x) \in \mathbb{R}, \\ g^{-1}(x) \geq 0 \end{Bmatrix}$

iv




c i $g(x) \to +\infty$ as $x \to 2$
Hence $\{g(x) \in \mathbb{R},\ g(x) > 0\}$

ii Letting $y = \dfrac{3}{x-2} \Rightarrow x = \dfrac{2y+3}{y}$
Hence $g^{-1}(x) = \dfrac{2x+3}{x}$

iii Domain g^{-1} = Range g
$\Rightarrow$ Domain $g^{-1}: \{x \in \mathbb{R},\ x > 0\}$
Range g^{-1} = Domain g
$\Rightarrow$ Range $g^{-1}(x): \{g^{-1}(x) \in \mathbb{R},\ g^{-1}(x) > 2\}$

iv




d i Minimum value of $g(x) = 2$
when $x = 7$
Hence $\{g(x) \in \mathbb{R},\ g(x) \geq 2\}$

ii Letting $y = \sqrt{x-3} \Rightarrow x = y^2 + 3$
Hence $g^{-1}(x) = x^2 + 3$

iii Domain g^{-1} = Range g
$\Rightarrow$ Domain $g^{-1}: \{x \in \mathbb{R},\ x \geq 2\}$
Range g^{-1} = Domain g
$\Rightarrow$ Range $g^{-1}(x): \{g^{-1}(x) \in \mathbb{R},\ g^{-1}(x) \geq 7\}$

iv



© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

4 e i $2^2 + 2 = 6$

Hence $\{g(x) \in \mathbb{R},\ g(x) > 6\}$

ii Letting $y = x^2 + 2$

$y - 2 = x^2$

$x = \sqrt{y-2}$

Hence $g^{-1}(x) = \sqrt{x-2}$


iii Domain g^{-1} = Range g

$\Rightarrow$ Domain $g^{-1}: \{x \in \mathbb{R},\ x > 6\}$

Range g^{-1} = Domain g

$\Rightarrow$ Range $g^{-1}(x): \begin{cases} g^{-1}(x) \in \mathbb{R}, \\ g^{-1}(x) > 2 \end{cases}$

iv



f i Minimum value of $g(x) = 0$

when $x = 2$

Hence $\{g(x) \in \mathbb{R},\ g(x) \geq 0\}$

ii Letting $y = x^3 - 8 \Rightarrow x = \sqrt[3]{y+8}$

Hence $g^{-1}(x) = \sqrt[3]{x+8}$


iii Domain g^{-1} = Range g

$\Rightarrow$ Domain $g^{-1}: \{x \in \mathbb{R},\ x \geq 0\}$

Range g^{-1} = Domain g

$\Rightarrow$ Range $g^{-1}(x): \begin{cases} g^{-1}(x) \in \mathbb{R}, \\ g^{-1}(x) \geq 2 \end{cases}$

iv



5 $t(x) = x^2 - 6x + 5,\ \{x \in \mathbb{R},\ x \geq 5\}$

Let $y = x^2 - 6x + 5$

$y = (x-3)^2 - 9 + 5$ (completing the square)

$y = (x-3)^2 - 4$

This has a minimum point at $(3, -4)$

For the domain $x \geq 5$, $t(x)$ is a one-to-one function so we can find an inverse function.

Make y the subject:

$y = (x-3)^2 - 4$

$y + 4 = (x-3)^2$

$\sqrt{y+4} = x - 3$

$\sqrt{y+4} + 3 = x$


© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

5 (continued)

Domain t^{-1} = Range t

$\Rightarrow$ Domain $g^{-1}: \{x \in \mathbb{R}, x \geq 0\}$

Hence, $t^{-1}(x) = \sqrt{x+4} + 3, \quad \{x \in \mathbb{R}, x \geq 0\}$




6 a $m(x) = x^2 + 4x + 9, \quad \{x \in \mathbb{R}, \ x > a\}$

Let $y = x^2 + 4x + 9$

$y = (x+2)^2 - 4 + 9$

$y = (x+2)^2 + 5$

This has a minimum value of $(-2, 5)$



For $m(x)$ to have an inverse it must be one-to-one. Hence the least value of a is -2

b Changing the subject of the formula:

$y = (x+2)^2 + 5$

$y - 5 = (x+2)^2$

$\sqrt{y-5} = x + 2$

$\sqrt{y-5} - 2 = x$


Hence $m^{-1}(x) = \sqrt{x-5} - 2$

c Domain of $m^{-1}(x)$: $\{x \in \mathbb{R}, \ x > 5\}$

7 a As $x \to 2$, $h(x) \to \frac{5}{0}$

and hence $h(x) \to \infty$

b To find $h^{-1}(3)$ we can find what element of the domain gets mapped to 3



Suppose $h(a) = 3$ for some a such that $a \neq 2$

Then $\frac{2a+1}{a-2} = 3$

$2a + 1 = 3a - 6$

$7 = a$

So $h^{-1}(3) = 7$

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

7 c Let $y = \dfrac{2x+1}{x-2}$ and find x as a function of y

$$y(x-2) = 2x+1$$
$$yx-2y = 2x+1$$
$$yx-2x = 2y+1$$
$$x(y-2) = 2y+1$$
$$x = \frac{2y+1}{y-2}$$

So $\mathrm{h}^{-1}(x) = \dfrac{2x+1}{x-2}, \quad \{x \in \mathbb{R},\ x \neq 2\}$

d If an element b is mapped to itself, then $\mathrm{h}(b) = b$

$$\frac{2b+1}{b-2} = b$$
$$2b+1 = b(b-2)$$
$$2b+1 = b^2-2b$$
$$0 = b^2-4b-1$$
$$b = \frac{4 \pm \sqrt{16+4}}{2} = \frac{4 \pm \sqrt{20}}{2}$$
$$= \frac{4 \pm 2\sqrt{5}}{2} = 2 \pm \sqrt{5}$$

The elements $2+\sqrt{5}$ and $2-\sqrt{5}$ get mapped to themselves by the function.

8 a $\mathrm{nm}(x) = \mathrm{n}(2x+3)$

$$= \frac{2x+3-3}{2}$$
$$= x$$

b $\mathrm{mn}(x) = \mathrm{m}\left(\dfrac{x-3}{2}\right)$

$$= 2\left(\frac{x-3}{2}\right)+3$$
$$= x$$

The functions m(x) and n(x) are the inverse of each other as mn(x) = nm(x) = x.

9 $\mathrm{st}(x) = \mathrm{s}\left(\dfrac{3-x}{x}\right)$

$$= \frac{3}{\left(\frac{3-x}{x}+1\right)}$$
$$= \frac{3}{\left(\frac{3-x+x}{x}\right)}$$
$$= x$$

$\mathrm{st}(x) = \mathrm{t}\left(\dfrac{3}{x+1}\right)$

$$= \frac{\left(3-\frac{3}{x+1}\right)}{\left(\frac{3}{x+1}\right)}$$
$$= \frac{\left(\frac{3x+3-3}{x+1}\right)}{\left(\frac{3}{x+1}\right)}$$
$$= x$$

The functions s(x) and t(x) are the inverse of each other as st(x) = ts(x) = x

10 a Let $y = 2x^2 - 3$

The domain of $\mathrm{f}^{-1}(x)$ is the range of $\mathrm{f}(x)$.

$\mathrm{f}(x) = 2x^2-3, \ \{x \in \mathbb{R},\ x < 0\}$ has range $\mathrm{f}(x) > -3$

Letting $y = 2x^2-3 \Rightarrow x = \pm\sqrt{\dfrac{x+3}{2}}$

We need to consider the domain of $\mathrm{f}(x)$ to determine if *either*

$\mathrm{f}^{-1}(x) = +\sqrt{\dfrac{x+3}{2}}$ *or* $\mathrm{f}^{-1}(x) = -\sqrt{\dfrac{x+3}{2}}$


$\mathrm{f}(x) = 2x^2-3$ has domain $\{x \in \mathbb{R},\ x < 0\}$

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

10 a (continued)

Hence $f^{-1}(x)$ must be the negative square root

$$f^{-1}(x) = -\sqrt{\frac{x+3}{2}}, \quad \{x \in \mathbb{R},\ x > -3\}$$



b If $f(a) = f^{-1}(a)$ then a is negative (see graph).

Solve $f(a) = a$

$$2a^2 - 3 = a$$

$$2a^2 - a - 3 = 0$$

$$(2a - 3)(a + 1) = 0$$

$$a = \frac{3}{2}, -1$$

Therefore $a = -1$

11 a Range of f(x) is f(x) > −5

b Let $y = f(x)$

$y = e^x - 5$

$e^x = y + 5$


$x = \ln(y + 5)$

$f^{-1}(x) = \ln(x + 5)$

Range of f(x) is f(x) > −5,

so domain of $f^{-1}(x)$ is $\{x \in \mathbb{R}, x > -5\}$

c



d Let $y = g(x)$

$y = \ln(x - 4)$

$e^y = x - 4$

$x = e^y + 4$

$g^{-1}(x) = e^x + 4$

Range of g(x) is $g(x) \in \mathbb{R}$,

so domain of $g^{-1}(x)$ is $\{x \in \mathbb{R}\}$

e $g^{-1}(x) = 11$

$e^x + 4 = 11$

$e^x = 7$

$x = \ln 7$

$x = 1.95$

12 a

$$f(x) = \frac{3(x+2)}{x^2 + x - 20} - \frac{2}{x-4}$$

$$= \frac{3(x+2)}{(x+5)(x-4)} - \frac{2}{x-4}$$

$$= \frac{3(x+2)}{(x+5)(x-4)} - \frac{2(x+5)}{(x+5)(x-4)}$$

$$= \frac{3x + 6 - 2x - 10}{(x+5)(x-4)}$$

$$= \frac{x-4}{(x+5)(x-4)}$$

$$= \frac{1}{x+5}, \quad x > 4$$

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

12 b The range of f is

$$\left\{ \mathrm{f}(x) \in \mathbb{R},\ \mathrm{f}(x) < \frac{1}{9} \right\}$$

c Let $y = \mathrm{f}(x)$

$$y = \frac{1}{x+5}$$

$$yx + 5y = 1$$

$$yx = 1 - 5y$$

$$x = \frac{1-5y}{y}$$

$$x = \frac{1}{y} - 5$$

$$\mathrm{f}^{-1}(x) = \frac{1}{x} - 5$$

The domain of $\mathrm{f}^{-1}(x)$ is

$$\left\{ x \in \mathbb{R}, x < \frac{1}{9} \text{ and } x \neq 0 \right\}$$

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.