Trigonometry and Modelling Mixed Exercise

1 a i $\sin 40° \cos 10° - \cos 40° \sin 10°$

$= \sin(40° - 10°) = \sin 30° = \frac{1}{2}$

ii $\frac{1}{\sqrt{2}} \cos 15° - \frac{1}{\sqrt{2}} \sin 15°$

$= \cos 45° \cos 15° - \sin 45° \sin 15°$

$= \cos(45° + 15°) = \cos 60° = \frac{1}{2}$

iii $\frac{1 - \tan 15°}{1 + \tan 15°} = \frac{\tan 45° - \tan 15°}{1 + \tan 45° \tan 15°}$


$= \tan(45° - 15°) = \tan 30° = \frac{\sqrt{3}}{3}$

2 As $\cos(x - y) = \sin y$

$\cos x \cos y + \sin x \sin y = \sin y \qquad (1)$

Draw a right-angled triangle,

where $\sin x = \frac{1}{\sqrt{5}}$



Using Pythagoras' theorem,

$a^2 = \left(\sqrt{5}\right)^2 - 1 = 4 \Rightarrow a = 2$

So $\cos x = \frac{2}{\sqrt{5}}$

Substitute into (1):

$\frac{2}{\sqrt{5}} \cos y + \frac{1}{\sqrt{5}} \sin y = \sin y$

$\Rightarrow 2\cos y + \sin y = \sqrt{5} \sin y$

$\Rightarrow 2\cos y = \sin y\left(\sqrt{5} - 1\right)$

$\Rightarrow \frac{2}{\sqrt{5} - 1} = \tan y \qquad \left(\tan y = \frac{\sin y}{\cos y}\right)$

$\Rightarrow \tan y = \frac{2\left(\sqrt{5} + 1\right)}{\left(\sqrt{5} - 1\right)\left(\sqrt{5} + 1\right)}$

$= \frac{2\left(\sqrt{5} + 1\right)}{4} = \frac{\sqrt{5} + 1}{2}$

3 a $\tan A = 2, \ \tan B = \frac{1}{3}$ since $y = \frac{1}{3}x - \frac{1}{3}$

b The angle required is $(A - B)$.


Using $\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$

$= \frac{2 - \frac{1}{3}}{1 + 2 \times \frac{1}{3}} = \frac{\frac{5}{3}}{\frac{5}{3}} = 1$

$\therefore A - B = 45°$

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

4



Using $\frac{\sin B}{b} = \frac{\sin C}{c}$

$\Rightarrow \frac{\sin(\theta - 30°)}{4} = \frac{\sin(\theta + 30°)}{5}$

$\Rightarrow 5\sin(\theta - 30°) = 4\sin(\theta + 30°)$

$\Rightarrow 5(\sin\theta\cos 30° - \cos\theta\sin 30°)$

$= 4(\sin\theta\cos 30° + \cos\theta\sin 30°)$

$\Rightarrow \sin\theta\cos 30° = 9\cos\theta\sin 30°$

$\Rightarrow \frac{\sin\theta}{\cos\theta} = 9\frac{\sin 30°}{\cos 30°} = 9\tan 30°$

$\Rightarrow \tan\theta = 9 \times \frac{\sqrt{3}}{3} = 3\sqrt{3}$

5 As the three values are consecutive terms of an arithmetic progression,

$\sin(\theta - 30°) - \sqrt{3}\cos\theta = \sin\theta - \sin(\theta - 30°)$

$\Rightarrow 2\sin(\theta - 30°) = \sin\theta + \sqrt{3}\cos\theta$

$\Rightarrow 2(\sin\theta\cos 30° - \cos\theta\sin 30°)$

$= \sin\theta + \sqrt{3}\cos\theta$

$\Rightarrow \sqrt{3}\sin\theta - \cos\theta = \sin\theta + \sqrt{3}\cos\theta$


$\Rightarrow \sin\theta\left(\sqrt{3} - 1\right) = \cos\theta\left(\sqrt{3} + 1\right)$

$\Rightarrow \tan\theta = \frac{\sqrt{3}+1}{\sqrt{3}-1}$

Calculator value is $\theta = \tan^{-1}\frac{\sqrt{3}+1}{\sqrt{3}-1} = 75°$

No other values as θ is acute.

6 a



$\sin A = \frac{3}{5}, \cos A = \frac{4}{5}$ $\quad \sin B = \frac{5}{13}, \cos B = \frac{12}{13}$

i $\sin(A + B) = \sin A\cos B + \cos A\sin B$

$= \frac{3}{5} \times \frac{12}{13} + \frac{4}{5} \times \frac{5}{13} = \frac{56}{65}$

ii $\tan 2B = \frac{2\tan B}{1 - \tan^2 B}$

$= \frac{2 \times \frac{5}{12}}{1 - \left(\frac{5}{12}\right)^2} = \frac{\frac{5}{6}}{\frac{119}{144}}$

$= \frac{5}{6} \times \frac{144}{119} = \frac{120}{119}$

b $\cos C = \cos\left(180° - (A + B)\right)$

$= -\cos(A + B)$

$= -(\cos A\cos B - \sin A\sin B)$

$= -\left(\frac{4}{5} \times \frac{12}{13} - \frac{3}{5} \times \frac{5}{13}\right)$

$= -\frac{33}{65}$

7 a $\cos 2x \equiv 1 - 2\sin^2 x$


$= 1 - 2\left(\frac{2}{\sqrt{5}}\right)^2 = 1 - \frac{8}{5} = -\frac{3}{5}$

b $\cos 2y \equiv 2\cos^2 y - 1$

$= 2\left(\frac{3}{\sqrt{10}}\right)^2 - 1 = 2\left(\frac{9}{10}\right) - 1 = \frac{4}{5}$

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

7 c



i $\tan(x+y)=\dfrac{\tan x+\tan y}{1-\tan x\tan y}$

$=\dfrac{2+\frac{1}{3}}{1-\frac{2}{3}}=\dfrac{\frac{7}{3}}{\frac{1}{3}}=7$

ii $\tan(x-y)=\dfrac{\tan x-\tan y}{1+\tan x\tan y}=\dfrac{\frac{5}{3}}{\frac{5}{3}}=1$

As x and y are acute, and $x>y$,

$x-y$ is acute

So $x-y=\dfrac{\pi}{4}$ $\left(\text{it cannot be } \dfrac{5\pi}{4}\right)$

8 a $\sin(x+y)\equiv\sin x\cos y+\cos x\sin y$

$=\dfrac{1}{2}+\dfrac{1}{3}=\dfrac{5}{6}$

$5\sin(x-y)\equiv 5(\sin x\cos y-\cos x\sin y)$

$=5\left(\dfrac{1}{2}-\dfrac{1}{3}\right)=5\times\dfrac{1}{6}=\dfrac{5}{6}$

b $\dfrac{\sin x\cos y}{\cos x\sin y}=\dfrac{\frac{1}{2}}{\frac{1}{3}}=\dfrac{3}{2}$

$\Rightarrow \dfrac{\tan x}{\tan y}=\dfrac{3}{2}$

so $\tan x=\dfrac{3\tan y}{2}=\dfrac{3k}{2}$

c $\tan 2x=\dfrac{2\tan x}{1-\tan^2 x}=\dfrac{3k}{1-\frac{9}{4}k^2}$

$=\dfrac{12k}{4-9k^2}$

9 a $\sqrt{3}\sin 2\theta+2\sin^2\theta=1$

$\sqrt{3}\sin 2\theta=1-2\sin^2\theta=\cos 2\theta$

$\dfrac{\sin 2\theta}{\cos 2\theta}=\dfrac{1}{\sqrt{3}}\Rightarrow \tan 2\theta=\dfrac{1}{\sqrt{3}}$

b $\tan 2\theta=\dfrac{1}{\sqrt{3}}$, for $0\leq 2\theta\leq 2\pi$

$2\theta=\dfrac{\pi}{6},\dfrac{7\pi}{6}\Rightarrow \theta=\dfrac{\pi}{12},\dfrac{7\pi}{12}$

10 a $\cos 2\theta=5\sin\theta$

$\Rightarrow \cos 2\theta-5\sin\theta=0$

$\Rightarrow 1-2\sin^2\theta-5\sin\theta=0$

$\Rightarrow 2\sin^2\theta+5\sin\theta-1=0$

$a=2$, $b=5$ and $c=-1$

b $2\sin^2\theta+5\sin\theta-1=0$

Using the quadratic formula

$\sin\theta=\dfrac{-5\pm\sqrt{5^2-4(2)(-1)}}{2(2)}$

$=\dfrac{-5\pm\sqrt{33}}{4}$

$\sin\theta=0.1861$, for $-\pi\leq\theta\leq\pi$

$\sin\theta$ is positive so solutions in the first and second quadrants

$\theta=\sin^{-1}0.1861,\ \pi-\sin^{-1}0.1861$

$\theta=0.187,\ 2.954$ (3 d.p.)

11 a $\cos(x-60°)=\cos x\cos 60°+\sin x\sin 60°$

$=\dfrac{1}{2}\cos x+\dfrac{\sqrt{3}}{2}\sin x$

So $2\sin x=\dfrac{1}{2}\cos x+\dfrac{\sqrt{3}}{2}\sin x$

$\Rightarrow\left(2-\dfrac{\sqrt{3}}{2}\right)\sin x=\dfrac{1}{2}\cos x$

$\Rightarrow \tan x=\dfrac{\frac{1}{2}}{2-\frac{\sqrt{3}}{2}}=\dfrac{1}{2}\times\dfrac{2}{4-\sqrt{3}}=\dfrac{1}{4-\sqrt{3}}$

b $\tan x=\dfrac{1}{4-\sqrt{3}}=0.44$ (2 d.p.), in the interval $0°\leq\theta\leq 360°$

$\tan\theta$ is positive so solutions in the first and third quadrants

$x=23.8°,\ 203.8°$ (1 d.p.)

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

12 a $\cos(x+20°) = \sin(90° - 20° - x)$
$= \sin(70° - x)$
$= \sin 70°\cos x - \cos 70°\sin x \quad (1)$

$4\sin(70° + x) = 4\sin 70°\cos x + 4\cos 70°\sin x \quad (2)$

As (1) = (2)

$4\sin 70°\cos x + 4\cos 70°\sin x = \sin 70°\cos x - \cos 70°\sin x$

$5\sin x\cos 70° = -3\sin 70°\cos x$

$\tan x = -\frac{3}{5}\tan 70°$

b $\tan x = -\frac{3}{5}\tan 70°$, for $0° \leq q \leq 180°$


$\tan q$ is negative so the solution is in the second quadrant

$x = 180° + \tan^{-1}\left(-\frac{3}{5}\tan 70°\right)$

$x = 180° - \tan^{-1}(-1.648)$

$x = 180° - (-58.8°) = 121.2°$ (1 d.p.)

13 a Draw a right-angled triangle and find $\sin a$ and $\cos a$.



$\Rightarrow \sin a = \frac{3}{5}, \cos a = \frac{4}{5}$

$3\sin(q+a) + 4\cos(q+a)$
$\equiv 3(\sin q\cos a + \cos q\sin a) + 4(\cos q\cos a - \sin q\sin a)$
$\equiv 3\left(\frac{4}{5}\sin q + \frac{3}{5}\cos q\right) + 4\left(\frac{4}{5}\cos q - \frac{3}{5}\sin q\right)$
$\equiv \frac{12}{5}\sin q + \frac{9}{5}\cos q + \frac{16}{5}\cos q - \frac{12}{5}\sin q$
$\equiv \frac{25}{5}\cos q \equiv 5\cos q$

b $\cos(x+270°)$
$= \cos x°\cos 270° - \sin x°\sin 270°$
$= (-0.8)(0) - (0.6)(-1)$
$= 0 + 0.6 = 0.6$

$\cos(x+540°)$
$= \cos x°\cos 540° - \sin x°\sin 540°$
$= (-0.8)(-1) - (0.6)(0)$
$= 0.8 - 0 = 0.8$

14 a One example is sufficient to disprove a statement. Let $A = 60°$, $B = 0°$

$\sec(A+B) = \sec(60° + 0°)$
$= \sec 60° = \frac{1}{\cos 60°} = 2$

$\sec A = \sec 60° = \frac{1}{\cos 60°} = 2$

$\sec B = \sec 0° = \frac{1}{\cos 0°} = 1$

So $\sec A + \sec B = 2 + 1 = 3$

So $\sec(60° + 0°) \neq \sec 60° + \sec 0°$

$\therefore \sin(A+B) \neq \sec A + \sec B$ is not true for all values of A, B.

b LHS $\equiv \tan q + \cot q$
$\equiv \frac{\sin q}{\cos q} + \frac{\cos q}{\sin q}$
$\equiv \frac{\sin^2 q + \cos^2 q}{\sin q\cos q}$
$\equiv \frac{1}{\frac{1}{2}\sin 2q}$

Using $\sin^2 q + \cos^2 q \equiv 1$, and $\sin 2q \equiv 2\sin q\cos q$

So LHS $\equiv \frac{2}{\sin 2q}$
$\equiv 2\operatorname{cosec} 2q$
$\equiv$ RHS

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

15 a Using $\tan 2\theta = \dfrac{2\tan\theta}{1 - \tan^2\theta}$ with $\theta = \dfrac{\pi}{8}$

$\Rightarrow \quad \tan\dfrac{\pi}{4} = \dfrac{2\tan\frac{\pi}{8}}{1 - \tan^2\frac{\pi}{8}}$

Let $t = \tan\dfrac{\pi}{8}$

So $1 = \dfrac{2t}{1 - t^2}$

$\Rightarrow \quad 1 - t^2 = 2t$

$\Rightarrow \quad t^2 + 2t - 1 = 0$

$\Rightarrow \quad t = \dfrac{-2 \pm \sqrt{8}}{2} = \dfrac{-2 \pm 2\sqrt{2}}{2}$

$= -1 \pm \sqrt{2}$

As $\dfrac{\pi}{8}$ is acute, $\tan\dfrac{\pi}{8}$ is positive, so $\tan\dfrac{\pi}{8} = \sqrt{2} - 1$

b $\tan\dfrac{3\pi}{8} = \tan\left(\dfrac{\pi}{4} + \dfrac{\pi}{8}\right) = \dfrac{\tan\frac{\pi}{4} + \tan\frac{\pi}{8}}{1 - \tan\frac{\pi}{4}\tan\frac{\pi}{8}}$

$= \dfrac{1 + \left(\sqrt{2} - 1\right)}{1 - \left(\sqrt{2} - 1\right)} = \dfrac{\sqrt{2}}{2 - \sqrt{2}}$

$= \dfrac{\sqrt{2}\left(2 + \sqrt{2}\right)}{\left(2 - \sqrt{2}\right)\left(2 + \sqrt{2}\right)}$

$= \dfrac{\sqrt{2}}{2}\left(2 + \sqrt{2}\right) = \sqrt{2} + 1$

16 a Let $\sin x - \sqrt{3}\cos x \equiv R\sin(x - \alpha)$

$\equiv R\sin x\cos\alpha - R\cos x\sin\alpha$

$R > 0,\ 0 < \alpha < 90°$

Compare $\sin x$: $R\cos\alpha = 1$ (1)

Compare $\cos x$: $R\sin\alpha = \sqrt{3}$ (2)


Divide (2) by (1): $\tan\alpha = \sqrt{3}$

$\Rightarrow \quad \alpha = 60°$

$R^2 = \left(\sqrt{3}\right)^2 + 1^2 = 4 \quad \Rightarrow \quad R = 2$

So $\sin x - \sqrt{3}\cos x \equiv 2\sin(x - 60°)$

b Sketch $y = 2\sin(x - 60°)$ by first translating $y = \sin x$ by $60°$ to the right and then stretching the result in the y direction by scale factor 2.



Graph meets y-axis when $x = 0$,

i.e. $y = 2\sin(-60°) = -\sqrt{3}$, at $\left(0, -\sqrt{3}\right)$

Graph meets x-axis when $y = 0$,

i.e. $(-300°,\ 0)$, $(-120°,\ 0)$, $(60°,\ 0)$, $240°,\ 0)$

17 a Let $7\cos 2\theta + 24\sin 2\theta \equiv R\cos(2\theta - \alpha)$

$\equiv R\cos 2\theta\cos\alpha + R\sin 2\theta\sin\alpha$

$R > 0,\ 0 < \alpha < \dfrac{\pi}{2}$

Compare $\cos 2\theta$: $R\cos\alpha = 7$ (1)

Compare $\sin 2\theta$: $R\sin\alpha = 24$ (2)

Divide (2) by (1): $\tan\alpha = \dfrac{24}{7}$

$\Rightarrow \quad \alpha = 1.29$ (2 d.p.)

$R^2 = 24^2 + 7^2 \quad \Rightarrow \quad R = 25$

So $7\cos 2\theta + 24\sin 2\theta \equiv 25\cos(2\theta - 1.29)$

b $14\cos 2\theta + 48\sin\theta\cos\theta$

$\equiv 14\left(\dfrac{1 + \cos 2\theta}{2}\right) + 24(2\sin\theta\cos\theta)$

$\equiv 7(1 + \cos 2\theta) + 24\sin 2\theta$

$\equiv 7 + 7\cos 2\theta + 24\sin 2\theta$

The maximum value of $7\cos 2\theta + 24\sin 2\theta$ is 25

(using (a) with $\cos(2\theta - 1.29) = 1$)

So maximum value of

$7 + 7\cos 2\theta + 24\sin 2\theta = 7 + 25 = 32$

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

17 c Using the answer to part a:
Solve $25\cos(2\theta - 1.29) = 12.5$

$\cos(2\theta - 1.29) = \frac{1}{2}$

$2\theta - 1.29 = -\frac{\pi}{3}, \frac{\pi}{3}$

$\theta = 0.119902..., 1.167099...$

$\theta = 6.9°, 66.9°$

18 a Let $1.5\sin 2x + 2\cos 2x \equiv R\sin(2x + \alpha)$
$\equiv R\sin 2x\cos\alpha + R\cos 2x\sin\alpha$

$R > 0, 0 < \alpha < \frac{\pi}{2}$

Compare $\sin 2x: R\cos\alpha = 1.5$ (1)

Compare $\cos 2x: R\sin\alpha = 2$ (2)

Divide (2) by (1): $\tan\alpha = \frac{4}{3}$

$\therefore \alpha = 0.927$ (3 d.p.)

$R^2 = 2^2 + 1.5^2 \therefore R = 2.5$

b $3\sin x\cos x + 4\cos^2 x$

$\equiv \frac{3}{2}(2\sin x\cos x) + 4\left(\frac{1+\cos 2x}{2}\right)$

$\equiv \frac{3}{2}\sin 2x + 2 + 2\cos 2x$

$\equiv \frac{3}{2}\sin 2x + 2\cos 2x + 2$

c From part (a) $1.5\sin 2x + 2\cos 2x$
$\equiv 2.5\sin(2x + 0.927)$

So maximum value of
$1.5\sin 2x + 2\cos 2x = 2.5 \times 1 = 2.5$

So maximum value of
$3\sin x\cos x + 4\cos^2 x = 2.5 + 2 = 4.5$

19 a $\sin^2\frac{\theta}{2} = 2\sin\theta$

$\frac{1 - \cos\theta}{2} = 2\sin\theta$

$1 - \cos\theta = 4\sin\theta$

$4\sin\theta + \cos\theta = 1$

Let $4\sin\theta + \cos\theta = R\sin(\theta + \alpha)$
$= R\sin\theta\cos\alpha + R\cos\theta\sin\alpha$

So $R\cos\alpha = 4$ and $R\sin\alpha = 1$

$\frac{R\sin\alpha}{R\cos\alpha} = \tan\alpha = \frac{1}{4}$

$\alpha = \tan^{-1}\left(\frac{1}{4}\right) = \tan^{-1}0.25 = 14.04$ (2 d.p.)

$R^2 = 4^2 + 1^2 = \sqrt{17}$

$4\sin\theta + \cos\theta = \sqrt{17}\sin(\theta + 14.04°) = 1$

b $\sqrt{17}\sin(\theta + 14.04°) = 1$, for $0° \leq \theta \leq 360°$

$\sin(\theta + 14.04°) = \frac{1}{\sqrt{17}} = 0.24$ (2 d.p.)

$\theta + 14.04° = \sin^{-1}0.24 = 14.04°$, for
$14.04° \leq \theta + 14.04° \leq 374.04°$

$\theta + 14.04° = 14.04°, 165.96°, 374.04°$

$\theta = 0°, 151.9°, 360°$

20 a $2\cos\theta = 1 + 3\sin\theta$
So $2\cos\theta - 3\sin\theta = 1$
Let $2\cos\theta - 3\sin\theta = R\cos(\theta + \alpha)$
$= R\cos\theta\cos\alpha - R\sin\theta\sin\alpha$

So $R\cos\alpha = 2$ and $R\sin\alpha = 3$

$\frac{R\sin\alpha}{R\cos\alpha} = \tan\alpha = \frac{3}{2}$

$\alpha = \tan^{-1}\left(\frac{3}{2}\right) = 56.3°$ (1 d.p.)

$R^2 = 2^2 + 3^2 = 13$

$R = \sqrt{13}$

So $2\cos\theta - 3\sin\theta = \sqrt{13}\cos(\theta + 56.3°) = 1$

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

20 b $\sqrt{13}\cos(\theta + 56.3°) = 1$, for $0° \leq \theta \leq 360°$

$\cos(\theta + 56.3°) = \dfrac{1}{\sqrt{13}}$,

for $56.3° \leq \theta + 56.3° \leq 416.3°$

$\theta + 56.3° = 73.9°, 286.1°$ (1 d.p.)

$\theta = 17.6°, 229.8°$ (1 d.p.)

21 a LHS $\equiv \dfrac{1}{\cos\theta} \times \dfrac{1}{\sin\theta} \equiv \dfrac{1}{\frac{1}{2}\sin 2\theta}$

$\equiv \dfrac{2}{\sin 2\theta} \equiv 2\operatorname{cosec} 2\theta \equiv$ RHS

b LHS $\equiv \dfrac{\tan\frac{\pi}{4} + \tan x}{1 - \tan\frac{\pi}{4}\tan x} - \dfrac{\tan\frac{\pi}{4} - \tan x}{1 + \tan\frac{\pi}{4}\tan x}$

$\equiv \dfrac{1 + \tan x}{1 - \tan x} - \dfrac{1 - \tan x}{1 + \tan x}$

$\equiv \dfrac{(1 + \tan x)^2 - (1 - \tan x)^2}{(1 - \tan x)(1 + \tan x)}$

$\equiv \dfrac{(1 + 2\tan x + \tan^2 x)}{1 - \tan^2 x} - \dfrac{(1 - 2\tan x + \tan^2 x)}{1 - \tan^2 x}$

$\equiv \dfrac{4\tan x}{1 - \tan^2 x}$

$\equiv 2\left(\dfrac{2\tan x}{1 - \tan^2 x}\right)$

$\equiv 2\tan 2x \equiv$ RHS

c LHS $\equiv (\sin x\cos y + \cos x\sin y) \times (\sin x\cos y - \cos x\sin y)$

$\equiv \sin^2 x\cos^2 y - \cos^2 x\sin^2 y$

$\equiv (1 - \cos^2 x)\cos^2 y - \cos^2 x(1 - \cos^2 y)$

$\equiv \cos^2 y - \cos^2 x\cos^2 y - \cos^2 x + \cos^2 x\cos^2 y$

$\equiv \cos^2 y - \cos^2 x =$ RHS

d LHS $\equiv 1 + 2\cos 2\theta + (2\cos^2 2\theta - 1)$

$\equiv 2\cos 2\theta + 2\cos^2 2\theta$

$\equiv 2\cos 2\theta(1 + \cos 2\theta)$

$\equiv 2\cos 2\theta(2\cos^2\theta)$

$\equiv 4\cos^2\theta\cos 2\theta =$ RHS

22 a LHS $\equiv \dfrac{1 - \cos 2x}{1 + \cos 2x} \equiv \dfrac{1 - (1 - 2\sin^2 x)}{1 + (2\cos^2 x - 1)}$

$\equiv \dfrac{2\sin^2 x}{2\cos^2 x} \equiv \tan^2 x =$ RHS

b $\tan^2 x = 3$

$\tan x = \pm\sqrt{3}$, for $-\pi \leq x \leq \pi$

$\tan x = \sqrt{3} \Rightarrow x = \dfrac{\pi}{3}, -\dfrac{2\pi}{3}$

$\tan x = -\sqrt{3} \Rightarrow x = -\dfrac{\pi}{3}, \dfrac{2\pi}{3}$

$x = -\dfrac{2\pi}{3}, -\dfrac{\pi}{3}, \dfrac{\pi}{3}, \dfrac{2\pi}{3}$

23 a LHS $\equiv \cos^4 2\theta - \sin^4 2\theta$

$\equiv (\cos^2 2\theta - \sin^2 2\theta)(\cos^2 2\theta + \sin^2 2\theta)$

$\equiv (\cos^2 2\theta - \sin^2 2\theta)(1)$

$\equiv \cos 4\theta \equiv$ RHS

b $\cos 4\theta = \dfrac{1}{2}$, for $0° \leq 4\theta \leq 720°$

$4\theta = 60°, 300°, 420°, 660°$

$\theta = 15°, 75°, 105°, 165°$

24 a LHS $\equiv \dfrac{1 - (1 - 2\sin^2\theta)}{2\sin\theta\cos\theta}$

$\equiv \dfrac{2\sin^2\theta}{2\sin\theta\cos\theta}$

$\equiv \dfrac{\sin\theta}{\cos\theta} \equiv \tan\theta =$ RHS

b When $\theta = 180°$, $\sin 2\theta = \sin 360° = 0$ and $2 - 2\cos 360° = 2 - 2 = 0$ therefore $\theta = 180°$ is a solution of the equation $\sin 2\theta = 2 - 2\cos 2\theta$

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

24 c Rearrange $\sin 2\theta = 2 - 2\cos 2\theta$ to give

$$\frac{2(1-\cos 2\theta)}{\sin 2\theta} = 1$$

Using the identity in part (a) gives

$2\tan\theta = 1$

$\Rightarrow \tan q = \frac{1}{2}$, for $0 < q < 360°$

$q = 26.6°, 206.6°$ (1 d.p.)

25 a Set $2\cos x - \sqrt{5}\sin x \equiv R\cos(x+\alpha)$

$\equiv R\cos x\cos a - R\sin x\sin a$

So $R\cos a = 2$ and $R\sin a = \sqrt{5}$

$$\frac{R\sin a}{R\cos a} = \tan a = \frac{\sqrt{5}}{2}$$

$$a = \tan^{-1}\left(\frac{\sqrt{5}}{2}\right) = 0.841 \text{ (3 d.p.)}$$

$$R^2 = 2^2 + \left(\sqrt{5}\right)^2 = 9$$

$R = 3$

$2\cos x - \sqrt{5}\sin x \equiv 3\cos(x + 0.841)$

b $3\cos(x + 0.841) = -1$,

for $0.841 \leq x + 0.841 < 2\pi + 0.841$

$$\cos(x + 0.841) = -\frac{1}{3}$$

$x + 0.841 = 1.911, 4.372$

$x = 1.07, 3.53$ (2 d.p.)

26 a Set $1.4\sin\theta - 5.6\cos\theta \equiv R\sin(\theta - \alpha)$

$\equiv R\sin\theta\cos a - R\cos\theta\sin a$

So $R\cos\alpha = 1.4$ and $R\sin\alpha = 5.6$

$$\frac{R\sin a}{R\cos a} = \tan a = \frac{5.6}{1.4}$$

$a = \tan^{-1} 4 == 75.964°$ (3 d.p.)

$R^2 = 1.4^2 + 5.6^2 = 33.32$

$R = 5.772$ (3 d.p.)

b The maximum value of $5.772\sin(\theta - 75.964)°$ is when $\sin(\theta - 75.964)° = 1$. So the maximum value is 5.772 and it occurs when $\theta - 75.964° = 90°$, $\theta = 165.964°$

c $$12 - 5.6\cos\left(\frac{360t}{365}\right)° + 1.4\sin\left(\frac{360t}{365}\right)°$$

$$\equiv 12 + 5.772\sin\left(\frac{360t}{365} - 75.964\right)°$$

The minimum number of daylight hours is when $\sin\left(\frac{360t}{365} - 75.964\right)° = -1$

So minimum is $12 - 5.772 = 6.228$ hours

d $\sin\left(\frac{360t}{365} - 75.964\right)° = -1$

$$\frac{360t}{365} - 75.964 = 270°$$

$t = 351$ days

27 a Let $12\sin x + 5\cos x \equiv R\sin(x + a)$

$\equiv R\sin x\cos a + R\cos x\sin a$

So $R\cos\alpha = 12$ and $R\sin\alpha = 5$

$$\frac{R\sin a}{R\cos a} = \tan a = \frac{5}{12}$$

$$a = \tan^{-1}\left(\frac{5}{12}\right) = 22.6° \text{ (1 d.p.)}$$

$R^2 = 12^2 + 5^2 = 169$

$R = 13$

So $12\sin x + 5\cos x = 13\sin(x + 22.6°)$

b $$v(x) = \frac{50}{12\sin\left(\frac{2x}{5}\right)° + 5\cos\left(\frac{2x}{5}\right)°}$$

$$= \frac{50}{13\sin\left(\frac{2x}{5} + 22.6°\right)}$$

The minimum value of v is when $\sin\left(\frac{2x}{5} + 22.6\right)° = 1$

So $\frac{50}{13} = 3.85$ m/s (2 d.p.)

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

27 c $\sin\left(\frac{2x}{5}+22.6°\right)=1$, for

$22.6° \leq \frac{2x}{5}+22.6° \leq 166.6°$

$\frac{2x}{5}+22.6°=90°$

$x=168.5$ minutes

Challenge

1 a Using $\cos P+\cos Q$

$$\equiv 2\cos\left(\frac{P+Q}{2}\right)\cos\left(\frac{P-Q}{2}\right)$$

and $\sin P-\sin Q$

$$\equiv 2\cos\left(\frac{P+Q}{2}\right)\sin\left(\frac{P-Q}{2}\right)$$

LHS $\equiv \frac{\cos 2q+\cos 4q}{\sin 2q-\sin 4q}$

$$\equiv \frac{2\cos\left(\frac{6q}{2}\right)\cos\left(\frac{2q}{2}\right)}{2\cos\left(\frac{6q}{2}\right)\sin\left(\frac{-2q}{2}\right)}$$

$$\equiv \frac{2\cos 3q\cos q}{2\cos 3q\sin(-q)}$$

$$\equiv \frac{\cos q}{\sin(-q)} \equiv -\cot q$$

b LHS $\equiv \cos x+2\cos 3x+\cos 5x$

$\equiv \cos 5x+\cos x+2\cos 3x$

$\equiv 2\cos\left(\frac{6x}{2}\right)\cos\left(\frac{4x}{2}\right)+2\cos 3x$


$\equiv 2\cos 3x\cos 2x+2\cos 3x$

$\equiv 2\cos 3x(\cos 2x+1)$

$\equiv 2\cos 3x(2\cos^2 x)$

$\equiv 4\cos^2 x\cos 3x \equiv$ RHS

2 a As $\angle OAB=\angle OBA \Rightarrow \angle AOB=\pi-2q$, so $\angle BOD=2q$



$OB=1$

$OD=\cos 2q$

$BD=\sin 2q$

$AB=2\cos q$

$\sin q=\frac{BD}{AB}=\frac{BD}{2\cos q}$

So $BD=2\sin q\cos q$

But $BD=\sin 2q$

So $\sin 2q \equiv 2\sin q\cos q$

b $AB=2\cos q$

$AD=(2\cos q)\cos q=2\cos^2 q$

$OD=2\cos^2 q-1$

From part (a) $OD=\cos 2q$

So $\cos 2q \equiv 2\cos^2 q-1$

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.