The normal distribution 3F

1 a i Yes, since $n = 120$ is large (> 50) and $p = 0.6$ is close to 0.5.

ii $\mu = np = 120 \times 0.6 = 72$ and $\sigma = \sqrt{np(1-p)} = \sqrt{72 \times 0.4} = \sqrt{28.8} = 5.37$ (3 s.f.)
$X \sim \text{B}(72, 5.37^2)$

b i No, $n = 20$ is not large enough (< 50).

c i Yes, since $n = 250$ is large (> 50) and $p = 0.52$ is close to 0.5.

ii $\mu = np = 250 \times 0.52 = 130$ and $\sigma = \sqrt{np(1-p)} = \sqrt{130 \times 0.48} = \sqrt{62.4} = 7.90$ (3 s.f.)
$X \sim \text{B}(130, 7.90^2)$

d i No, $p = 0.85$ is too far from 0.5.

e i Yes, since $n = 400$ is large (> 50) and $p = 0.48$ is close to 0.5.

ii $\mu = np = 400 \times 0.48 = 192$ and $\sigma = \sqrt{np(1-p)} = \sqrt{192 \times 0.52} = \sqrt{99.84} = 9.99$ (3 s.f.)
$X \sim \text{B}(192, 9.99^2)$

f i Yes, since $n = 1000$ is large (> 50) and $p = 0.58$ is close to 0.5.

ii $\mu = np = 1000 \times 0.58 = 580$ and $\sigma = \sqrt{np(1-p)} = \sqrt{580 \times 0.42} = \sqrt{243.6} = 15.6$ (3 s.f.)
$X \sim \text{B}(580, 15.6^2)$

2 A normal approximation is valid since $n = 150$ is large (> 50) and $p = 0.45$ is close to 0.5.
$\mu = np = 150 \times 0.45 = 67.5$ and $\sigma = \sqrt{np(1-p)} = \sqrt{67.5 \times 0.55} = \sqrt{37.125} = 6.093$ (4 s.f.)

a $\text{P}(X \leqslant 60) \approx \text{P}(Y < 60.5) = 0.1253$ (4 s.f.)

b $\text{P}(X > 75) \approx \text{P}(Y > 75.5) = 0.0946$ (4 s.f.)

c $\text{P}(65 \leqslant X \leqslant 80) \approx \text{P}(64.5 < Y < 80.5) = 0.6723$ (4 s.f.)

3 A normal approximation is valid since $n = 200$ is large (> 50) and $p = 0.53$ is close to 0.5.
$\mu = np = 200 \times 0.53 = 106$ and $\sigma = \sqrt{np(1-p)} = \sqrt{106 \times 0.47} = \sqrt{49.82} = 7.058$ (4 s.f.)

a $\text{P}(X < 90) \approx \text{P}(Y < 89.5) = 0.0097$ (4 s.f.)

b $\text{P}(100 \leqslant X < 110) \approx \text{P}(99.5 < Y < 109.5) = 0.5115$ (4 s.f.)

c $\text{P}(X = 105) \approx \text{P}(104.5 < Y < 105.5) = 0.0559$ (4 s.f.)

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

4 A normal approximation is valid since $n = 100$ is large (> 50) and $p = 0.6$ is close to 0.5.
$\mu = np = 100 \times 0.6 = 60$ and $\sigma = \sqrt{np(1-p)} = \sqrt{60 \times 0.4} = \sqrt{24} = 4.899$ (4 s.f.)

a $P(X > 58) \approx P(Y > 58.5) = 0.6203$ (4 s.f.)

b $P(60 < X \leqslant 72) \approx P(60.5 < Y < 72.5) = 0.4540$ (4 s.f.)

c $P(X = 70) \approx P(69.5 < Y < 70.5) = 0.0102$ (4 s.f.)

5 Let X = number of heads in 70 tosses of a fair coin, so $X \sim B(70, 0.5)$.
Since $p = 0.5$ and 70 is large, X can be approximated by the normal distribution $Y \sim N(\mu, \sigma^2)$,
where $\mu = 70 \times 0.5 = 35$ and $\sigma = \sqrt{70 \times 0.5 \times 0.5} = \sqrt{17.5}$
So $Y \sim N(35, 17.5)$
$P(X > 45) \approx P(Y \geqslant 45.5) = 0.0060$

6 A normal approximation is valid since $n = 1200$ is large and p is close to 0.5.
$\mu = np = 1200 \times \frac{50}{101} = 594.059$
and $\sigma = \sqrt{np(1-p)} = \sqrt{594.059 \times \frac{51}{101}} = \sqrt{299.97059\ldots} = 17.32$ (4 s.f.)
So $Y \sim N(594.059, 299.97...)$
$P(X \geqslant 600) \approx P(Y > 599.5) = 0.3767$ (4 s.f.)

7 **a** The number of trials, n, must be large (> 50), and the success probability, p, must be close to 0.5.

b Using the binomial distribution, $P(X = 10) = \binom{20}{10} \times 0.45^{10} \times 0.55^{10} = 0.1593$ (4 s.f.)

c A normal approximation is valid since $n = 240$ is large and $p = 0.45$ is close to 0.5.
$\mu = np = 240 \times 0.45 = 108$ and $\sigma = \sqrt{np(1-p)} = \sqrt{108 \times 0.55} = \sqrt{59.4} = 7.707$ (4 s.f.)
So $Y \sim N(108, 59.4)$
$P(X < 110) \approx P(Y < 109.5) = 0.5772$ (4 s.f.)

d $P(X \geqslant q) = 0.2 \Rightarrow P(Y > (q - 0.5)) = 0.2$
Using the inverse normal function,
$P(Y > (q - 0.5)) = 0.2 \Rightarrow q - 0.5 = 114.485 \Rightarrow q = 114.985$
So $q = 115$

8 **a** Using the cumulative binomial function with $N = 30$ and $p = 0.52$,
$P(X < 17) = P(X \leqslant 16) = 0.6277$ (4 s.f.)

b A normal approximation is valid since $n = 600$ is large and $p = 0.52$ is close to 0.5.
$\mu = np = 600 \times 0.52 = 312$ and $\sigma = \sqrt{np(1-p)} = \sqrt{312 \times 0.48} = \sqrt{149.76} = 12.24$ (4 s.f.)
So $Y \sim N(312, 149.76)$
$P(300 \leqslant X \leqslant 350) \approx P(299.5 < Y < 350.5) = 0.8456$ (4 s.f.)

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.

9 a Using the binomial distribution, $P(X = 55) = \binom{100}{55} \times 0.56^{55} \times 0.44^{45} = 0.07838$ (4 s.f.)

b A normal approximation is valid since $n = 100$ is large and $p = 0.56$ is close to 0.5.

$\mu = np = 100 \times 0.56 = 56$ and $\sigma = \sqrt{np(1-p)} = \sqrt{56 \times 0.44} = \sqrt{24.64} = 4.964$ (4 s.f.)

So $Y \sim B(56, 24.64)$

$P(X = 55) \approx P(54.5 < Y < 55.5) = 0.07863$ (4 s.f.)

Percentage error = $\dfrac{0.07838... - 0.07863...}{0.07838...} \times 100 = -0.31\%$ (2 d.p.)

© Pearson Education Ltd 2017. Copying permitted for purchasing institution only. This material is not copyright free.