Please check the examination details below before entering your candidate information

Candidate surname StemUp Model Solution | Other names

Pearson Edexcel Level 3 GCE

Centre Number | Candidate Number

Thursday 16 May 2019

Afternoon | Paper Reference **8FM0-21**

Further Mathematics

Advanced Subsidiary
Further Mathematics options
21: Further Pure Mathematics 1
(Part of options A, B, C and D)

You must have:
Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- The total mark for this part of the examination is 40. There are 5 questions.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶

P61862A

©2019 Pearson Education Ltd.

1/1/1/1

P 6 1 8 6 2 A 0 1 1 6

Pearson

stemup.co.uk

1. (a) Write down the t-formula for $\sin x$.

(1)

(b) Use the answer to part (a)

(i) to find the exact value of $\sin x$ when

$$\tan\left(\frac{x}{2}\right) = \sqrt{2}$$

(ii) to show that

$$\cos x = \frac{1 - t^2}{1 + t^2}$$

(4)

(c) Use the t-formulae to solve for $0 < \theta \leqslant 360°$

$$7\sin\theta + 9\cos\theta + 3 = 0$$

giving your answers to one decimal place.

(4)

(a) $\sin x = \frac{2t}{1 + t^2}$ where $t = \tan\left(\frac{x}{2}\right)$

(b)(i) $\tan\left(\frac{x}{2}\right) = \sqrt{2} \Rightarrow t = \sqrt{2}$

$\sin x = \frac{2(\sqrt{2})}{1 + (\sqrt{2})^2} = \frac{2\sqrt{2}}{1 + 2} = \frac{2\sqrt{2}}{3}$

(b)(ii) $\tan x = \frac{2t}{1 - t^2}$ (double angle formula with $t = \tan\left(\frac{x}{2}\right)$)

$\cos x = \frac{\sin x}{\tan x} = \frac{2t}{1 + t^2} \times \frac{1 - t^2}{2t}$

$\therefore \cos x = \frac{1 - t^2}{1 + t^2}$ as required

(c) $t = \tan\left(\frac{\theta}{2}\right)$: $7\left(\frac{2t}{1 + t^2}\right) + 9\left(\frac{1 - t^2}{1 + t^2}\right) + 3 = 0$

$\times (1 + t^2)$: $14t + 9 - 9t^2 + 3 + 3t^2 = 0$

$-6t^2 + 14t + 12 = 0 \Rightarrow 3t^2 - 7t - 6 = 0$

$(t - 3)(3t + 2) = 0 \Rightarrow t = 3$ or $t = -2/3$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

stemup.co.uk

Question 1 continued

$0 < \theta \leqslant 360° \Rightarrow 0 < \frac{\theta}{2} \leqslant 180°$

$t = 3: \quad \frac{\theta}{2} = \arctan 3 = 71.5650...° \Rightarrow \theta = 143.1301...°$

$t = -\frac{2}{3}: \quad \frac{\theta}{2} = 180° + \arctan\left(-\frac{2}{3}\right) = 146.3099...°$

$\Rightarrow \theta = 292.6198...°$

$\theta = 143.1°, \quad \theta = 292.6°$

(Total for Question 1 is 9 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

2. A student was set the following problem.

Use algebra to find the set of values of x for which

$$\frac{x}{x-24} > \frac{1}{x+11}$$

The student's attempt at a solution is written below.

$$x(x-24)(x+11)^2 > (x+11)(x-24)^2$$

$$x(x-24)(x+11)^2 - (x+11)(x-24)^2 > 0$$

$$(x-24)(x+11)\left[x(x+11) - x - 24\right] > 0 \qquad \text{Line 3}$$

$$(x-24)(x+11)\left[x^2 + 10x - 24\right] > 0$$

$$(x-24)(x+11)(x+12)(x-2) > 0$$

$$x = 24,\ x = -11,\ x = -12,\ x = 2$$

$$\{x \in \mathbb{R} : -12 < x < -11\} \cup \{x \in \mathbb{R} : 2 < x < 24\} \qquad \text{Line 7}$$

There are errors in the student's solution.

(a) Identify the error made

(i) in line 3

(ii) in line 7

(2)

(b) Find a correct solution to this problem.

(4)

(a)(i) Line 3: bracketing error. The second term should be $x(x + 11) - (x - 24)$, i.e. $-x + 24$, not $-x - 24$.

(a)(ii) Line 7: they have found the regions where the expression is < 0. It should be $x < -12$ or $-11 < x < 2$ or $x > 24$.

(b) Multiply both sides by $(x - 24)^2(x + 11)^2$ (positive):

$x(x - 24)(x + 11)^2 > (x + 11)(x - 24)^2$

$x(x - 24)(x + 11)^2 - (x + 11)(x - 24)^2 > 0$

$(x - 24)(x + 11)[x(x + 11) - (x - 24)] > 0$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

stemup.co.uk

Question 2 continued

$(x - 24)(x + 11)[x^2 + 10x + 24] > 0$

$(x - 24)(x + 11)(x + 6)(x + 4) > 0$

Critical values: $x = -11$, $x = -6$, $x = -4$, $x = 24$

Quartic with positive x^4 coefficient is positive on the two outside intervals and the middle interval:

$\{x \in \mathbb{R} : x < -11\} \cup \{x \in \mathbb{R} : -6 < x < -4\} \cup \{x \in \mathbb{R} : x > 24\}$

(Total for Question 2 is 6 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

3. Julie decides to start a business breeding rabbits to sell as pets.

Initially she buys 20 rabbits. After t years the number of rabbits, R, is modelled by the differential equation

$$\frac{dR}{dt} = 2R + 4\sin t \qquad t > 0$$

Julie needs to have at least 40 rabbits before she can start to sell them.

Use two iterations of the approximation formula

$$\left(\frac{dy}{dx}\right)_n \approx \frac{y_{n+1} - y_n}{h}$$

to find out if, according to the model, Julie will be able to start selling rabbits after 4 months.

(7)

4 months = $\frac{1}{3}$ year, two iterations $\Rightarrow$ $h = \frac{1}{6}$

$R_{n+1} \approx R_n + h\left(\frac{dR}{dt}\right)_n$

$t_0 = 0$, $R_0 = 20$: $\left(\frac{dR}{dt}\right)_0 = 2(20) + 4\sin 0 = 40$

$R_1 = 20 + \frac{1}{6}(40) = \frac{80}{3} = 26.666\ldots$

$t_1 = \frac{1}{6}$: $\left(\frac{dR}{dt}\right)_1 = 2\left(\frac{80}{3}\right) + 4\sin\left(\frac{1}{6}\right) = 53.99691\ldots$

$R_2 = \frac{80}{3} + \frac{1}{6}(53.99691\ldots) = 35.66615\ldots$

$R_2 \approx 35.7$, i.e. about 35 or 36 rabbits after 4 months

35.666... < 40, so according to the model Julie will not be able to start selling rabbits after 4 months.

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

stemup.co.uk

Question 3 continued

(Total for Question 3 is 7 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

4.




Figure 1

Figure 1 shows a sketch of a solid doorstop made of wood. The doorstop is modelled as a tetrahedron.

Relative to a fixed origin O, the vertices of the tetrahedron are A (2, 1, 4), B (6, 1, 2), C (4, 10, 3) and D (5, 8, d), where d is a positive constant and the units are in centimetres.

(a) Find the area of the triangle ABC.

(4)

Given that the volume of the doorstop is 21 cm^3

(b) find the value of the constant d.

(4)

(a) $AB = \begin{pmatrix} 6-2 \\ 1-1 \\ 2-4 \end{pmatrix} = \begin{pmatrix} 4 \\ 0 \\ -2 \end{pmatrix}$, $AC = \begin{pmatrix} 4-2 \\ 10-1 \\ 3-4 \end{pmatrix} = \begin{pmatrix} 2 \\ 9 \\ -1 \end{pmatrix}$

$AB \times AC = \begin{pmatrix} (0)(-1)-(-2)(9) \\ (-2)(2)-(4)(-1) \\ (4)(9)-(0)(2) \end{pmatrix} = \begin{pmatrix} 18 \\ 0 \\ 36 \end{pmatrix}$

Area $ABC = \frac{1}{2}|AB \times AC| = \frac{1}{2}\sqrt{(18^2 + 0^2 + 36^2)}$

$= \frac{1}{2}\sqrt{1620} = \frac{1}{2}(18\sqrt{5}) = 9\sqrt{5} = 20.1246...$

Area of triangle $ABC = 9\sqrt{5}\ cm^2$ $(\approx 20.1\ cm^2)$

(b) $AD = \begin{pmatrix} 5-2 \\ 8-1 \\ d-4 \end{pmatrix} = \begin{pmatrix} 3 \\ 7 \\ d-4 \end{pmatrix}$

Volume $= \frac{1}{6}|AD \cdot (AB \times AC)|$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

P 6 1 8 6 2 A 0 8 1 6
stemup.co.uk

Question 4 continued

$$AD \cdot (AB \times AC) = \begin{pmatrix} 3 \\ 7 \\ d-4 \end{pmatrix} \cdot \begin{pmatrix} 18 \\ 0 \\ 36 \end{pmatrix} = 54 + 0 + 36(d-4) = 36d - 90$$

$$\frac{1}{6}|36d - 90| = 21 \Rightarrow |36d - 90| = 126$$

$36d - 90 = 126 \Rightarrow d = 6$ ($36d - 90 = -126$ gives $d = -1$, rejected as $d > 0$)

$d = 6$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

stemup.co.uk

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

(Total for Question 4 is 8 marks)

stemup.co.uk

5.




Figure 2

Figure 2 shows a sketch of part of the rectangular hyperbola H with equation

$$xy = c^2 \qquad x > 0$$

where c is a positive constant.

The point $P\left(ct, \frac{c}{t}\right)$ lies on H.

The line l is the tangent to H at the point P.

The line l crosses the x-axis at the point A and crosses the y-axis at the point B.

The region R, shown shaded in Figure 2, is bounded by the x-axis, the y-axis and the line l.

Given that the length OB is twice the length of OA, where O is the origin, and that the area of R is 32, find the exact coordinates of the point P.

(10)

$y = c^2x^{-1} \Rightarrow \frac{dy}{dx} = -c^2x^{-2} = -\frac{c^2}{x^2}$

At $P\left(ct, \frac{c}{t}\right)$: $m = -\frac{c^2}{c^2t^2} = -\frac{1}{t^2}$

Tangent l: $y - \frac{c}{t} = -\frac{1}{t^2}(x - ct)$

$\Rightarrow y = -\frac{1}{t^2}x + \frac{2c}{t}$ (or $x + t^2y = 2ct$)

$y = 0$: $x = 2ct \Rightarrow A(2ct, 0)$; $x = 0$: $y = \frac{2c}{t} \Rightarrow B\left(0, \frac{2c}{t}\right)$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

stemup.co.uk

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 5 continued

$OB = 2OA$: $\frac{2c}{t} = 2(2ct) \Rightarrow 2c = 4ct^2 \Rightarrow t^2 = \frac{1}{2}$

$t > 0$ (P has $x > 0$): $t = \frac{1}{\sqrt{2}} = \frac{\sqrt{2}}{2}$

Area $R = \frac{1}{2}(OA)(OB) = \frac{1}{2}(2ct)\left(\frac{2c}{t}\right) = 2c^2 = 32$

$c^2 = 16 \Rightarrow c = 4 \quad (c > 0)$

$x_P = ct = 4 \times \frac{1}{\sqrt{2}} = 2\sqrt{2}, \qquad y_P = \frac{c}{t} = 4\sqrt{2}$

$P(2\sqrt{2}, 4\sqrt{2})$

Question 5 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

stemup.co.uk

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 5 continued

Question 5 continued

(Total for Question 5 is 10 marks)

TOTAL FOR FURTHER PURE MATHEMATICS 1 IS 40 MARKS

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

stemup.co.uk