Please check the examination details below before entering your candidate information

Candidate surname StemUp Model Solution | Other names

Centre Number | Candidate Number

Pearson Edexcel Level 3 GCE

Friday 19 May 2023

Afternoon

Paper reference **8FM0/23**

Further Mathematics

Advanced Subsidiary
Further Mathematics options
23: Further Statistics 1
(Part of options B, E, F and G)

You must have:
Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
  – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Values from statistical tables should be quoted in full. If a calculator is used instead of tables the value should be given to an equivalent degree of accuracy.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- The total mark for this part of the examination is 40. There are 4 questions.
- The marks for **each** question are shown in brackets
  – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶

P72809A

©2023 Pearson Education Ltd.
N:1/1/1/1/

P 7 2 8 0 9 A 0 1 1 2

Pearson

1. The discrete random variable X has the following distribution

x	0	1	2	3	4
$P(X = x)$	r	k	$\frac{k}{2}$	$\frac{k}{3}$	$\frac{k}{4}$

where r and k are positive constants.

The standard deviation of X equals the mean of X

Find the exact value of r

(6)

$E(X) = 0 \times r + 1 \times k + 2 \times \frac{k}{2} + 3 \times \frac{k}{3} + 4 \times \frac{k}{4} = k + k + k + k = 4k$

$E(X^2) = 1^2 \times k + 2^2 \times \frac{k}{2} + 3^2 \times \frac{k}{3} + 4^2 \times \frac{k}{4} = k + 2k + 3k + 4k = 10k$

$Var(X) = E(X^2) - [E(X)]^2 = 10k - 16k^2$

$\sqrt{Var(X)} = E(X) \Rightarrow Var(X) = [E(X)]^2$

$10k - 16k^2 = (4k)^2 = 16k^2$

$10k = 32k^2 \Rightarrow k = \frac{10}{32} = \frac{5}{16}$ $(k \neq 0$ since $k > 0)$

$\Sigma P(X = x) = 1: \quad r + k + \frac{k}{2} + \frac{k}{3} + \frac{k}{4} = 1$

$r + \frac{1}{12}(12k + 6k + 4k + 3k) = 1 \Rightarrow r + \frac{25}{12}k = 1$

$r = 1 - \frac{25}{12} \times \frac{5}{16} = 1 - \frac{125}{192}$

$r = \frac{67}{192}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 1 continued

(Total for Question 1 is 6 marks)

2. A bag contains a large number of balls, all of the same size and weight. The balls are coloured Red, Blue or Yellow.

Jasmine asks each child in a group of 150 children to close their eyes, select a ball from the bag and show it to her. The child then replaces the ball and repeats the process a second time.

If both balls are the same colour the child receives a prize.

The results are given in the table below.

1st colour / 2nd colour	Red	Blue	Yellow	Total
Red	31	11	18	60
Blue	8	10	9	27
Yellow	21	9	33	63
Total	60	30	60	150

Jasmine carries out a test, at the 5% level of significance, to see whether or not the colour of the 2nd ball is independent of the colour of the 1st ball.

(a) Calculate the expected frequencies for the cases where both balls are the same colour.

(2)

The test statistic Jasmine obtained was 12.712 to three decimal places.

(b) Use this value to complete the test, stating the critical value and conclusion clearly.

(3)

With reference to your calculations in part (a) and the nature of the experiment,

(c) give a plausible reason why Jasmine may have obtained her conclusion in part (b).

(1)

(a) $E(RR) = \frac{60 \times 60}{150} = 24$ $E(BB) = \frac{30 \times 27}{150} = 5.4$

$E(YY) = \frac{60 \times 63}{150} = 25.2$

(b) All $E_i > 5$ (smallest is $E(BB) = 5.4$), so no need to pool

$\nu = (3 - 1)(3 - 1) = 4$

Critical value: $\chi^2_4(5\%) = 9.488$

$12.712 > 9.488$, so the result is significant: reject H_0

There is evidence that the colour of the 2nd ball is not independent of the colour of the 1st ball.

(c) $O_i > E_i$ for all pairings of the same colour (31 > 24, 10 > 5.4,

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 2 continued

33 > 25.2), so same-colour pairs happen more often than expected.

Some children may have been cheating (looking / peeking) when they chose the 2nd ball so as to win a prize, so the choice of the 2nd ball is not independent of the 1st.

(Total for Question 2 is 6 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

3. A machine produces cloth. Faults occur randomly in the cloth at a rate of 0.4 per square metre.

The machine is used to produce tablecloths, each of area A square metres. One of these tablecloths is taken at random.

The probability that this tablecloth has no faults is 0.0907

(a) Find the value of A

(3)

The tablecloths are sold in packets of 20

A randomly selected packet is taken.

(b) Find the probability that more than 1 of the tablecloths in this packet has no faults.

(3)

A hotel places an order for 100 tablecloths each of area A square metres.

The random variable X represents the number of these tablecloths that have no faults.

(c) Find

(i) E(X)

(ii) Var(X)

(3)

(d) Use a Poisson approximation to estimate P($X = 10$)

(2)

It is claimed that a new machine produces cloth with a rate of faults that is less than 0.4 per square metre.

A piece of cloth produced by this new machine is taken at random.
The piece of cloth has area 30 square metres and is found to have 6 faults.

(e) Stating your hypotheses clearly, use a suitable test to assess the claim made for the new machine. Use a 5% level of significance.

(4)

(f) Write down the p-value for the test used in part (e).

(1)

(a) F = number of faults in A m^2, $F \sim Po(0.4A)$

$P(F = 0) = e^{-0.4A} = 0.0907$

$-0.4A = \ln 0.0907 \Rightarrow A = 6.0004...$

$A = 6$

(b) T = number of tablecloths in the packet with no faults

$T \sim B(20, 0.0907)$

$P(T > 1) = 1 - P(T \leq 1)$

$= 1 - (0.9093^{20} + 20 \times 0.0907 \times 0.9093^{19})$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 3 continued

$= 0.55276\ldots = 0.553$

(c) $X \sim B(100, 0.0907)$

(i) $E(X) = 100 \times 0.0907 = 9.07$

(ii) $\text{Var}(X) = 100 \times 0.0907 \times (1 - 0.0907) = 8.247351$

$= 8.25$

(d) X can be approximated by $Po(9.07)$

$$P(X = 10) \approx \frac{e^{-9.07} \times 9.07^{10}}{10!} = 0.119470\ldots$$

$P(X = 10) \approx 0.119$

(e) λ = mean number of faults per square metre (new machine)

H_0: $\lambda = 0.4$ H_1: $\lambda < 0.4$

Y = number of faults in 30 m^2. Under H_0: $Y \sim Po(30 \times 0.4) = Po(12)$

$P(Y \leqslant 6) = 0.04582\ldots$

$0.04582 < 0.05$, so the result is significant: reject H_0

There is evidence to support the claim that the new machine produces cloth with a rate of faults less than 0.4 per square metre.

(f) p-value $= P(Y \leqslant 6) = 0.04582\ldots = 0.0458$

Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 3 continued

(Total for Question 3 is 16 marks)

4. Table 1 below shows the number of car breakdowns in the *Snoreap* district in each of 60 months.

Number of car breakdowns	0	1	2	3	4	5
Frequency	12	11	19	14	3	1

Table 1

Anja believes that the number of car breakdowns per month in *Snoreap* can be modelled by a Poisson distribution. Table 2 below shows the results of some of her calculations.

Number of car breakdowns	0	1	2	3	4	$\geqslant 5$
Observed frequency (O_i)	12	11	19	14	3	1
Expected frequency (E_i)	9.92	17.85	16.07	9.64	4.34	2.18

Table 2

(a) State suitable hypotheses for a test to investigate Anja's belief.

(1)

(b) Explain why Anja has changed the label of the final column to $\geqslant 5$

(1)

(c) Showing your working clearly, complete Table 2

(4)

(d) Find the value of $\dfrac{(O_i - E_i)^2}{E_i}$ when the number of car breakdowns is

(i) 1

(ii) 3

(2)

(e) Explain why Anja used 3 degrees of freedom for her test.

(2)

The test statistic for Anja's test is 6.54 to 2 decimal places.

(f) Stating the critical value and using a 5% level of significance, complete Anja's test.

(2)

(a) H_0: The number of car breakdowns per month can be modelled by a Poisson distribution

H_1: The number of car breakdowns per month cannot be modelled by a Poisson distribution

(b) A Poisson distribution gives a (non-zero) probability to all values

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

greater than 5 (it can take any value 0, 1, 2, ...), so the last class must include all of these values: ≥ 5

(c) $\lambda = \frac{0 \times 12 + 1 \times 11 + 2 \times 19 + 3 \times 14 + 4 \times 3 + 5 \times 1}{12 + 11 + 19 + 14 + 3 + 1} = \frac{108}{60} = 1.8$

Under H_0: $X \sim Po(1.8)$

$E_1 = 60 \times P(X = 1) = 60 \times \frac{e^{-1.8} \times 1.8}{1!} = 17.852... = 17.85$

$E_2 = 60 \times P(X = 2) = 60 \times \frac{e^{-1.8} \times 1.8^2}{2!} = 16.067... = 16.07$

$E_{\geq 5} = 60 - (9.92 + 17.85 + 16.07 + 9.64 + 4.34) = 2.18$

(or $60 \times (1 - P(X \leq 4)) = 2.1843...$)

(d) (i) Breakdowns = 1:

$\frac{(O_i - E_i)^2}{E_i} = \frac{(11 - 17.852...)^2}{17.852...} = 2.630... = 2.63$

(ii) Breakdowns = 3:

$\frac{(O_i - E_i)^2}{E_i} = \frac{(14 - 9.64)^2}{9.64} = 1.97195... = 1.97$

(e) The expected frequencies for 4 and ≥ 5 are both less than 5 (4.34 and 2.18), so these last two columns are combined, leaving 5 classes. The mean λ was estimated from the observed data, so there are 2 constraints: $\nu = 5 - 2 = 3$

(f) Critical value: $\chi^2_3(5\%) = 7.815$

6.54 < 7.815, so the result is not significant: do not reject H_0

There is insufficient evidence to reject Anja's belief: the number of car breakdowns per month can be modelled by a Poisson distribution.

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

(Total for Question 4 is 12 marks)

TOTAL FOR FURTHER STATISTICS 1 IS 40 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA