Please check the examination details below before entering your candidate information

Candidate surname StemUp Model Solution | Other names

Centre Number | Candidate Number

Pearson Edexcel Level 3 GCE

Monday 13 May 2024

Afternoon (Time: 1 hour 40 minutes)

Paper reference **8FM0/01**

Further Mathematics

Advanced Subsidiary

PAPER 1: Core Pure Mathematics

You must have:
Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
  – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
  – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶

P74069A

©2024 Pearson Education Ltd.
F:1/1/1/1/1/1/1/

P 7 4 0 6 9 A 0 1 3 2

Pearson

1. The cubic equation

$$2x^3 - 3x^2 + 5x + 7 = 0$$

has roots α, β and γ.

Without solving the equation, determine the exact value of

(i) $\alpha^2 + \beta^2 + \gamma^2$

(3)

(ii) $\frac{3}{\alpha} + \frac{3}{\beta} + \frac{3}{\gamma}$

(3)

(iii) $(5 - \alpha)(5 - \beta)(5 - \gamma)$

(3)

$\alpha + \beta + \gamma = \frac{3}{2}$, $\quad \alpha\beta + \beta\gamma + \gamma\alpha = \frac{5}{2}$, $\quad \alpha\beta\gamma = -\frac{7}{2}$

(i) $\alpha^2 + \beta^2 + \gamma^2 = (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \beta\gamma + \gamma\alpha) = \left(\frac{3}{2}\right)^2 - 2\left(\frac{5}{2}\right)$

$= \frac{9}{4} - 5 = -\frac{11}{4}$

(ii) $\frac{3}{\alpha} + \frac{3}{\beta} + \frac{3}{\gamma} = \frac{3(\alpha\beta + \beta\gamma + \gamma\alpha)}{\alpha\beta\gamma} = \frac{3 \times 5/2}{-7/2} = -\frac{15}{7}$

(iii) $(5 - \alpha)(5 - \beta)(5 - \gamma) = 125 - 25(\alpha + \beta + \gamma) + 5(\alpha\beta + \beta\gamma + \gamma\alpha) - \alpha\beta\gamma$

$= 125 - 25\left(\frac{3}{2}\right) + 5\left(\frac{5}{2}\right) + \frac{7}{2}$

$= 125 - \frac{75}{2} + \frac{25}{2} + \frac{7}{2} = \frac{207}{2} \quad (= 103.5)$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 1 continued

Question 1 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 1 continued

(Total for Question 1 is 9 marks)

2. With respect to the **right-hand rule**, a rotation through $\theta°$ anticlockwise about the z-axis is represented by the matrix

$$\begin{pmatrix} \cos\theta & -\sin\theta & 0 \\ \sin\theta & \cos\theta & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Given that the matrix **M**, where

$$\mathbf{M} = \begin{pmatrix} -\frac{\sqrt{3}}{2} & \frac{1}{2} & 0 \\ -\frac{1}{2} & -\frac{\sqrt{3}}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

represents a rotation through $\alpha°$ anticlockwise about the z-axis with respect to the **right-hand rule**,

(a) determine the value of α.

(1)

(b) Hence determine the smallest possible positive integer value of k for which $\mathbf{M}^k = \mathbf{I}$

(2)

The 3×3 matrix **N** represents a reflection in the plane with equation $y = 0$

(c) Write down the matrix **N**.

(1)

The point A has coordinates $(-2, 4, 3)$

The point B is the image of the point A under the transformation represented by matrix **M** followed by the transformation represented by matrix **N**.

(d) Show that the coordinates of B are $(2 + \sqrt{3}, 2\sqrt{3} - 1, 3)$

(2)

Given that O is the origin,

(e) show that, to 3 significant figures, the size of angle AOB is 66.9°

(2)

(f) Hence determine the area of triangle AOB, giving your answer to 3 significant figures.

(2)

(a) $\cos\alpha° = -\frac{\sqrt{3}}{2}$, $\sin\alpha° = -\frac{1}{2}$ $\Rightarrow$ $\alpha = 210$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 2 continued

(b) Need $210k$ to be a multiple of 360: $210k = 360m \Rightarrow 7k = 12m$

$\therefore$ smallest positive integer $k = 12$ $(12 \times 210° = 2520° = 7 \times 360°)$

(c) $N = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$

(d) $MA = \begin{pmatrix} -\sqrt{3}/2 & 1/2 & 0 \\ -1/2 & -\sqrt{3}/2 & 0 \\ 0 & 0 & 1 \end{pmatrix}\begin{pmatrix} -2 \\ 4 \\ 3 \end{pmatrix} = \begin{pmatrix} \sqrt{3}+2 \\ 1-2\sqrt{3} \\ 3 \end{pmatrix}$

$NMA = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}\begin{pmatrix} \sqrt{3}+2 \\ 1-2\sqrt{3} \\ 3 \end{pmatrix} = \begin{pmatrix} \sqrt{3}+2 \\ 2\sqrt{3}-1 \\ 3 \end{pmatrix}$

$\therefore$ B is $(2 + \sqrt{3}, 2\sqrt{3} - 1, 3)$ as required

(e) $\begin{pmatrix} -2 \\ 4 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} 2+\sqrt{3} \\ 2\sqrt{3}-1 \\ 3 \end{pmatrix} = -4 - 2\sqrt{3} + 8\sqrt{3} - 4 + 9 = 1 + 6\sqrt{3}$

$|OA| = \sqrt{(4 + 16 + 9)} = \sqrt{29}$, $|OB| = \sqrt{29}$ (rotation and reflection preserve length)

$\cos AOB = \dfrac{1 + 6\sqrt{3}}{\sqrt{29}\,\sqrt{29}} = \dfrac{1 + 6\sqrt{3}}{29} = 0.39283...$

$AOB = 66.868...° = 66.9°$ (3 s.f.) as required

(f) Area $= \dfrac{1}{2} \times \sqrt{29} \times \sqrt{29} \times \sin 66.9° = 14.5 \times \sin 66.9°$

$= 13.337... = 13.3$ (3 s.f.)

Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 2 continued

(Total for Question 2 is 10 marks)

3. (a) Use the standard results for summations to show that, for all positive integers n,

$$\sum_{r=1}^{n} r^2(r+1) = \frac{1}{12}n(n+1)(n+2)(an+b)$$

where a and b are integers to be determined.

(4)

(b) Hence show that, for all positive integers k,

$$\sum_{r=k+1}^{3k} r^2(r+1) = \frac{1}{3}k(3k+1)(Ak^2+Bk+C)$$

where A, B and C are integers to be determined.

(3)

(c) Hence, using algebra and making your method clear, determine the value of k for which

$$25\sum_{r=k+1}^{3k} r^2(r+1) = 192k^3(3k+1)$$

(3)

(a) $\sum_{r=1}^{n} r^2(r+1) = \sum_{r=1}^{n} r^3 + \sum_{r=1}^{n} r^2 = \frac{1}{4}n^2(n+1)^2 + \frac{1}{6}n(n+1)(2n+1)$

$= \frac{1}{12}n(n+1)[3n(n+1) + 2(2n+1)]$

$= \frac{1}{12}n(n+1)(3n^2+7n+2)$

$= \frac{1}{12}n(n+1)(n+2)(3n+1)$ $\quad \therefore a = 3,\ b = 1$

(b) $\sum_{r=k+1}^{3k} r^2(r+1) = \sum_{r=1}^{3k} r^2(r+1) - \sum_{r=1}^{k} r^2(r+1)$

$= \frac{1}{12}(3k)(3k+1)(3k+2)(9k+1) - \frac{1}{12}k(k+1)(k+2)(3k+1)$

$= \frac{1}{12}k(3k+1)[3(3k+2)(9k+1) - (k+1)(k+2)]$

$= \frac{1}{12}k(3k+1)[81k^2+63k+6-k^2-3k-2]$

$= \frac{1}{12}k(3k+1)(80k^2+60k+4)$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 3 continued

$= \frac{1}{3}k(3k + 1)(20k^2 + 15k + 1)$ $\quad \therefore A = 20, \ B = 15, \ C = 1$

(c) $\frac{25}{3}k(3k + 1)(20k^2 + 15k + 1) = 192k^3(3k + 1)$

$k > 0$ so $k(3k + 1) \neq 0$: divide by $k(3k + 1)$ and multiply by 3

$25(20k^2 + 15k + 1) = 576k^2$

$500k^2 + 375k + 25 = 576k^2$

$76k^2 - 375k - 25 = 0$

$(k - 5)(76k + 5) = 0$

$k = 5$ or $k = -\frac{5}{76}$ (rejected, k must be a positive integer)

$\therefore k = 5$

Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 3 continued

(Total for Question 3 is 10 marks)

4.

$$\mathbf{A} = \begin{pmatrix} -1 & -2 & -7 \\ 3 & k & 2 \\ 1 & 1 & 4 \end{pmatrix} \qquad \mathbf{B} = \begin{pmatrix} 4k-2 & 1 & 7k-4 \\ -10 & 3 & -19 \\ 3-k & -1 & 6-k \end{pmatrix}$$

where k is a constant.

(a) Determine the value of the constant c for which

$$\mathbf{AB} = (3k + c)\mathbf{I}$$

(2)

(b) Hence determine the value of k for which $\mathbf{A}^{-1}$ does not exist.

(2)

Given that $\mathbf{A}^{-1}$ does exist,

(c) write down $\mathbf{A}^{-1}$ in terms of k.

(1)

(d) Use the answer to part (c) to solve the simultaneous equations

$$\begin{aligned} -x - 2y - 7z &= 10 \\ 3x + ky + 2z &= 3 \\ x + y + 4z &= 1 \end{aligned}$$

giving the values of x, y and z in simplest form in terms of k.

(3)

(a) Leading diagonal of AB, first element:

$(-1)(4k - 2) + (-2)(-10) + (-7)(3 - k) = -4k + 2 + 20 - 21 + 7k = 3k + 1$

(other elements check: e.g. row 1 × col 2 = $-1 - 6 + 7 = 0$)

$AB = (3k + 1)I \Rightarrow 3k + c = 3k + 1 \Rightarrow c = 1$

(b) $\det(AB) = (3k + 1)^3 = \det A \times \det B$, and $\det A = 3k + 1$

A^{-1} does not exist when $\det A = 0$: $3k + 1 = 0 \Rightarrow k = -\frac{1}{3}$

(c) $A^{-1} = \frac{1}{3k+1}\begin{pmatrix} 4k-2 & 1 & 7k-4 \\ -10 & 3 & -19 \\ 3-k & -1 & 6-k \end{pmatrix}$

(d) $\begin{pmatrix} x \\ y \\ z \end{pmatrix} = A^{-1}\begin{pmatrix} 10 \\ 3 \\ 1 \end{pmatrix} = \frac{1}{3k+1}\begin{pmatrix} 4k-2 & 1 & 7k-4 \\ -10 & 3 & -19 \\ 3-k & -1 & 6-k \end{pmatrix}\begin{pmatrix} 10 \\ 3 \\ 1 \end{pmatrix}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

$$= \frac{1}{3k+1}\begin{pmatrix} 40k - 20 + 3 + 7k - 4 \\ -100 + 9 - 19 \\ 30 - 10k - 3 + 6 - k \end{pmatrix} = \frac{1}{3k+1}\begin{pmatrix} 47k - 21 \\ -110 \\ 33 - 11k \end{pmatrix}$$

$$x = \frac{47k - 21}{3k + 1}, \quad y = -\frac{110}{3k + 1}, \quad z = \frac{33 - 11k}{3k + 1}$$

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 4 continued

(Total for Question 4 is 8 marks)

5. Given that on an Argand diagram the locus of points defined by $|z + 5 - 12i| = 10$ is a circle,

(a) write down,

(i) the coordinates of the centre of this circle,

(ii) the radius of this circle.

(2)

(b) Show, by shading on an Argand diagram, the set of points defined by

$$|z + 5 - 12i| \leqslant 10$$

(1)

(c) For the set of points defined in part (b), determine the maximum value of $|z|$

(3)

The set of points A is defined by

$$A = \{z : 0 \leqslant \arg(z + 5 - 20i) \leqslant \pi\} \cap \{z : |z + 5 - 12i| \leqslant 10\}$$


(d) Determine the area of the region defined by A, giving your answer to 3 significant figures.

(4)

(a) (i) centre (−5, 12)

(ii) radius = 10

(b)



(c) Distance of centre from O = $\sqrt{(5^2 + 12^2)} = \sqrt{169} = 13$

max |z| is at the point of the circle furthest from O, on the line through O and the centre:

$|z|_{max} = 13 + 10 = 23$

(d) $0 \leqslant \arg(z + 5 - 20i) \leqslant \pi$ is the region on or above the line $y = 20$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 5 continued

(boundary is the horizontal line through $-5 + 20i$)

so A is the segment of the circle above $y = 20$. Centre is $20 - 12 = 8$ below the chord.

$\cos\left(\frac{\theta}{2}\right) = \frac{8}{10} \Rightarrow \frac{\theta}{2} = 0.6435... \Rightarrow \theta = 1.2870...$ rad (angle at centre)

half-chord $= \sqrt{(10^2 - 8^2)} = 6$

Area $= \frac{1}{2} \times 10^2 \times \theta - \frac{1}{2} \times 12 \times 8$

$= 50 \times 1.2870... - 48 = 64.350... - 48 = 16.350...$

Area $= 16.4$ (3 s.f.)

Question 5 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 5 continued

(Total for Question 5 is 10 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

6. The drainage system for a sports field consists of underground pipes.

This situation is modelled with respect to a fixed origin O.

According to the model,

- the surface of the sports field is a plane with equation $z = 0$
- the pipes are straight lines
- one of the pipes, P_1, passes through the points $A(3, 4, -2)$ and $B(-2, -8, -3)$
- a different pipe, P_2, has equation $\dfrac{x-1}{2} = \dfrac{y-3}{4} = \dfrac{z+1}{-2}$
- the units are metres

(a) Determine a vector equation of the line representing the pipe P_1

(2)

(b) Determine the coordinates of the point at which the pipe P_1 meets the surface of the playing field, according to the model.

(2)

Determine, according to the model,

(c) the acute angle between pipes P_1 and P_2, giving your answer in degrees to 3 significant figures,

(3)

(d) the shortest distance between pipes P_1 and P_2

(5)

(a) $\overrightarrow{AB} = \begin{pmatrix}-2\\-8\\-3\end{pmatrix} - \begin{pmatrix}3\\4\\-2\end{pmatrix} = \begin{pmatrix}-5\\-12\\-1\end{pmatrix} \Rightarrow \mathbf{r} = \begin{pmatrix}3\\4\\-2\end{pmatrix} + \lambda\begin{pmatrix}5\\12\\1\end{pmatrix}$

(b) Surface $z = 0$: $-2 + \lambda = 0 \Rightarrow \lambda = 2$

$x = 3 + 10 = 13$, $y = 4 + 24 = 28$ $\Rightarrow$ point $(13, 28, 0)$

(c) Directions: $\mathbf{d}_1 = \begin{pmatrix}5\\12\\1\end{pmatrix}$, $\mathbf{d}_2 = \begin{pmatrix}2\\4\\-2\end{pmatrix}$ $\mathbf{d}_1 \cdot \mathbf{d}_2 = 10 + 48 - 2 = 56$

$\cos\theta = \dfrac{56}{\sqrt{(5^2 + 12^2 + 1^2)}\sqrt{(2^2 + 4^2 + 2^2)}} = \dfrac{56}{\sqrt{170}\sqrt{24}} = 0.87671\ldots$

$\theta = 28.751\ldots° = 28.8°$ (3 s.f.)

(d) General points: P_1: $(3 + 5\lambda, 4 + 12\lambda, -2 + \lambda)$, P_2: $(1 + 2\mu, 3 + 4\mu, -1 - 2\mu)$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 6 continued

$P_1 - P_2 = \begin{pmatrix} 2 + 5\lambda - 2\mu \\ 1 + 12\lambda - 4\mu \\ -1 + \lambda + 2\mu \end{pmatrix}$ perpendicular to both pipes:

$\begin{pmatrix} 2 + 5\lambda - 2\mu \\ 1 + 12\lambda - 4\mu \\ -1 + \lambda + 2\mu \end{pmatrix} \cdot \begin{pmatrix} 5 \\ 12 \\ 1 \end{pmatrix} = 0 \quad \Rightarrow \quad 170\lambda - 56\mu = -21$

$\begin{pmatrix} 2 + 5\lambda - 2\mu \\ 1 + 12\lambda - 4\mu \\ -1 + \lambda + 2\mu \end{pmatrix} \cdot \begin{pmatrix} 2 \\ 4 \\ -2 \end{pmatrix} = 0 \quad \Rightarrow \quad 56\lambda - 24\mu = -10$

Solving: $\lambda = \frac{7}{118}, \quad \mu = \frac{131}{236}$

$P_1 - P_2 = \frac{70}{59}i - \frac{30}{59}j + \frac{10}{59}k$

$|P_1 - P_2| = \frac{\sqrt{(70^2 + 30^2 + 10^2)}}{59} = \frac{\sqrt{5900}}{59} = \frac{10\sqrt{59}}{59} = 1.3018...$

Shortest distance = 1.30 m (3 s.f.)

Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 6 continued

(Total for Question 6 is 12 marks)

7. (i) Prove by induction that, for all positive integers n,

$$\sum_{r=1}^{n} \frac{1}{r(r+1)} = \frac{n}{n+1}$$

(5)

(ii) Prove by induction that, for all positive integers n,

$$f(n) = 3^{2n+4} - 2^{2n}$$

is divisible by 5

(5)

(i) $n = 1$: LHS $= \frac{1}{1 \times 2} = \frac{1}{2}$, RHS $= \frac{1}{1+1} = \frac{1}{2}$ so true for $n = 1$

Assume true for $n = k$: $\sum_{r=1}^{k} \frac{1}{r(r+1)} = \frac{k}{k+1}$

Then $\sum_{r=1}^{k+1} \frac{1}{r(r+1)} = \frac{k}{k+1} + \frac{1}{(k+1)(k+2)} = \frac{k(k+2)+1}{(k+1)(k+2)}$

$= \frac{k^2+2k+1}{(k+1)(k+2)} = \frac{(k+1)^2}{(k+1)(k+2)} = \frac{k+1}{k+2} = \frac{k+1}{(k+1)+1}$

which is the result for $n = k + 1$.

If the result is true for $n = k$ then it has been shown to be true for $n = k + 1$,

and as it is true for $n = 1$, the statement is true for all positive integers n.

(ii) $n = 1$: $f(1) = 3^6 - 2^2 = 729 - 4 = 725 = 5 \times 145$, divisible by 5, so true for $n = 1$

Assume true for $n = k$: $f(k) = 3^{2k+4} - 2^{2k}$ is divisible by 5

$f(k+1) = 3^{2(k+1)+4} - 2^{2(k+1)} = 3^{2k+6} - 2^{2k+2} = 9 \times 3^{2k+4} - 4 \times 2^{2k}$

$= 9 \times 3^{2k+4} - 9 \times 2^{2k} + 5 \times 2^{2k}$

$= 9(3^{2k+4} - 2^{2k}) + 5 \times 2^{2k} = 9f(k) + 5 \times 2^{2k}$

$9f(k)$ is divisible by 5 (assumption) and 5×2^{2k} is divisible by 5, so $f(k+1)$ is divisible by 5.

If the result is true for $n = k$ then it has been shown to be true for $n = k + 1$,

and as it is true for $n = 1$, the statement is true for all positive integers n.

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 7 continued

Question 7 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 7 continued

(Total for Question 7 is 10 marks)

8.




Figure 1




Figure 2

Figure 1 shows the central vertical cross-section, *OABCDEO*, of the design for a solid glass ornament.

Figure 2 shows the finite region, *R*, which is bounded by the *y*-axis, the horizontal line *CB*, the vertical line *BA*, and the curve *AO*.

The ornament is formed by rotating the region *R* through 360° about the *y*-axis.

The curve *AO* is modelled by the equation

$$x = ky^2 + \sqrt{y} \qquad 0 \leqslant y \leqslant 4$$

where *k* is a constant.

The point *A* has coordinates (0.4, 4) and the point *B* has coordinates (0.4, 4.5)

The units are centimetres.

(a) Determine the value of *k* according to this model.

(2)

(b) Use algebraic integration to determine the exact volume of glass that would be required to make the ornament, according to the model.

(7)

(c) State a limitation of the model.

(1)

When the ornament was manufactured, 9 cm^3 of glass was required.

(d) Use this information and your answer to part (b) to evaluate the model, explaining your reasoning.

(1)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 8 continued

(a) $A(0.4, 4)$: $0.4 = k \times 4^2 + \sqrt{4} \Rightarrow 0.4 = 16k + 2 \Rightarrow 16k = -1.6 \Rightarrow k = -0.1$

(b) Neck (cylinder, $4 \leq y \leq 4.5$): $V_1 = \pi \times 0.4^2 \times 0.5 = 0.08\pi = \frac{2}{25}\pi$

Curve: $V_2 = \pi\int_0^4 x^2\,dy = \pi\int_0^4 (\sqrt{y} - 0.1y^2)^2\,dy = \pi\int_0^4 (y - 0.2y^{5/2} + 0.01y^4)\,dy$

$= \pi\left[\frac{y^2}{2} - \frac{2}{35}y^{7/2} + \frac{1}{500}y^5\right]_0^4$

$= \pi\left(8 - \frac{2}{35} \times 128 + \frac{1}{500} \times 1024\right) - 0 = \pi\left(8 - \frac{256}{35} + \frac{256}{125}\right) = \frac{2392}{875}\pi$

$V = \frac{2392}{875}\pi + \frac{2}{25}\pi = \frac{2392}{875}\pi + \frac{70}{875}\pi = \frac{2462}{875}\pi \text{ cm}^3$

(c) There may be flaws or air bubbles within the glass, so the solid is not exactly this shape.

(d) $\frac{2462}{875}\pi = 8.8395\ldots \approx 8.84 \text{ cm}^3$, which is only 0.16 cm^3 less than 9 cm^3 (about 1.8% too small), so the model gives a good estimate of the glass required.

Question 8 continued

(Total for Question 8 is 11 marks)

TOTAL FOR CORE PURE MATHEMATICS IS 80 MARKS

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA