Please check the examination details below before entering your candidate information

Candidate surname StemUp Model Solution | Other names

Pearson Edexcel
International Advanced Level

Centre Number | Candidate Number

Time 1 hours 30 minutes

Paper reference **WFM03/01**

Mathematics

International Advanced Subsidiary/Advanced Level
Further Pure Mathematics F3

You must have:
Mathematical Formulae and Statistical Tables (Yellow), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
  – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets
  – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.
- Good luck with your examination.

Turn over ▶

P60705A

©2021 Pearson Education Ltd.

1/1/1/1/1/

P 6 0 7 0 5 A 0 1 3 2

Pearson

Leave blank

1. (a) Using the definitions of hyperbolic functions in terms of exponentials, show that

$$1 - \tanh^2 x \equiv \operatorname{sech}^2 x$$

(3)

(b) Solve the equation

$$2\operatorname{sech}^2 x + 3\tanh x = 3$$

giving your answer as an exact logarithm.

(3)

(a) $LHS = 1 - \tanh^2 x$

$= 1 - \left(\frac{e^x - e^{-x}}{e^x + e^{-x}}\right)^2$

$= \frac{(e^x + e^{-x})^2 - (e^x - e^{-x})^2}{(e^x + e^{-x})^2}$

$= \frac{(e^{2x} + 2 + e^{-2x}) - (e^{2x} - 2 + e^{-2x})}{(e^x + e^{-x})^2}$

$= \frac{4}{(e^x + e^{-x})^2} = \left(\frac{2}{e^x + e^{-x}}\right)^2 = \operatorname{sech}^2 x$

$\therefore 1 - \tanh^2 x = \operatorname{sech}^2 x$ as required

(b) $2(1 - \tanh^2 x) + 3\tanh x = 3$

$2 - 2\tanh^2 x + 3\tanh x = 3$

$2\tanh^2 x - 3\tanh x + 1 = 0$

$(2\tanh x - 1)(\tanh x - 1) = 0$

$\tanh x = \frac{1}{2}$ ($\tanh x = 1$ has no solution since $-1 < \tanh x < 1$)

$x = \operatorname{artanh}\frac{1}{2} = \frac{1}{2}\ln\left(\frac{1 + 0.5}{1 - 0.5}\right) = \frac{1}{2}\ln 3$

$x = \ln\sqrt{3}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 1 continued

Q1

(Total 6 marks)

Leave blank

2. A curve has equation

$$y = \sqrt{9 - x^2} \qquad 0 \leqslant x \leqslant 3$$

(a) Using calculus, show that the length of the curve is $\frac{3\pi}{2}$

(4)

The curve is rotated through 2π radians about the x-axis.

(b) Using calculus, find the exact area of the surface generated.

(3)

(a) $\frac{dy}{dx} = \frac{1}{2}(9 - x^2)^{-1/2} \times (-2x) = \frac{-x}{\sqrt{(9 - x^2)}}$

$1 + \left(\frac{dy}{dx}\right)^2 = 1 + \frac{x^2}{9 - x^2} = \frac{9}{9 - x^2}$

$s = \int_0^3 \sqrt{(1 + (dy/dx)^2)}\, dx = \int_0^3 \frac{3}{\sqrt{(9 - x^2)}}\, dx$

$= \left[3\arcsin\frac{x}{3}\right]_0^3$

$= 3\arcsin(1) - 3\arcsin(0) = 3 \times \frac{\pi}{2} - 0$

$= \frac{3\pi}{2}$ as required

(b) $S = 2\pi\int_0^3 y\sqrt{(1 + (dy/dx)^2)}\, dx$

$= 2\pi\int_0^3 \sqrt{(9 - x^2)} \times \frac{3}{\sqrt{(9 - x^2)}}\, dx$

$= 2\pi\int_0^3 3\, dx = 6\pi\left[x\right]_0^3$

$= 6\pi \times 3 = 18\pi$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 2 continued

Leave blank

Question 2 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 2 continued

Q2

(Total 7 marks)

Leave blank

3. $\mathbf{M} = \begin{pmatrix} 3 & 1 & p \\ 1 & 1 & 2 \\ -1 & p & 2 \end{pmatrix}$ where p is a real constant

(a) Find the exact values of p for which $\mathbf{M}$ has no inverse.

(4)

Given that $\mathbf{M}$ does have an inverse,

(b) find $\mathbf{M}^{-1}$ in terms of p.

(5)

(a) $\det M = 3(1\times2 - 2p) - 1(1\times2 - 2\times(-1)) + p(1\times p - 1\times(-1))$

$= 3(2 - 2p) - 1(4) + p(p + 1)$

$= 6 - 6p - 4 + p^2 + p = p^2 - 5p + 2$

No inverse when $\det M = 0$: $p^2 - 5p + 2 = 0$

$p = \frac{5 \pm \sqrt{(25 - 8)}}{2} \Rightarrow p = \frac{5 + \sqrt{17}}{2}$ or $p = \frac{5 - \sqrt{17}}{2}$

(b) Matrix of minors:

$\begin{pmatrix} 2 - 2p & 4 & p + 1 \\ 2 - p^2 & 6 + p & 3p + 1 \\ 2 - p & 6 - p & 2 \end{pmatrix}$

Matrix of cofactors:

$\begin{pmatrix} 2 - 2p & -4 & p + 1 \\ p^2 - 2 & 6 + p & -3p - 1 \\ 2 - p & p - 6 & 2 \end{pmatrix}$

Transpose (adjugate) and divide by $\det M$:

$M^{-1} = \frac{1}{p^2 - 5p + 2} \begin{pmatrix} 2 - 2p & p^2 - 2 & 2 - p \\ -4 & 6 + p & p - 6 \\ p + 1 & -3p - 1 & 2 \end{pmatrix}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 3 continued

Leave blank

Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 3 continued

Q3

(Total 9 marks)

Leave blank

4. (i) $f(x) = x \arccos x \quad -1 \leqslant x \leqslant 1$

Find the exact value of $f'(0.5)$.

(3)

(ii) $g(x) = \arctan(e^{2x})$

Show that

$$g''(x) = k \operatorname{sech}(2x) \tanh(2x)$$

where k is a constant to be found.

(5)

(i) $f'(x) = \arccos x + x \times \left(\frac{-1}{\sqrt{(1 - x^2)}} \right) = \arccos x - \frac{x}{\sqrt{(1 - x^2)}}$

$f'(0.5) = \arccos 0.5 - \frac{0.5}{\sqrt{(1 - 0.5^2)}} = \frac{\pi}{3} - \frac{0.5}{\sqrt{0.75}}$

$= \frac{\pi}{3} - \frac{1}{\sqrt{3}} = \frac{\pi}{3} - \frac{\sqrt{3}}{3} = \frac{\pi - \sqrt{3}}{3}$

(ii) $g'(x) = \frac{1}{1 + (e^{2x})^2} \times 2e^{2x} = \frac{2e^{2x}}{1 + e^{4x}}$

$= \frac{2}{e^{-2x} + e^{2x}}$ (dividing by e^{2x}) $= \frac{1}{\cosh(2x)} = \operatorname{sech}(2x)$

$g''(x) = -\operatorname{sech}(2x)\tanh(2x) \times 2$

$g''(x) = -2\operatorname{sech}(2x)\tanh(2x)$, so $k = -2$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 4 continued

Leave blank

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave blank

Question 4 continued

Q4

(Total 8 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave blank

5. $$I_n = \int \sec^n x \, dx \qquad n \geqslant 0$$

(a) Prove that for $n \geqslant 2$

$$(n-1)I_n = \tan x \sec^{n-2} x + (n-2)I_{n-2}$$

(6)

(b) Hence, showing each step of your working, find the exact value of

$$\int_0^{\frac{\pi}{4}} \sec^6 x \, dx$$

(4)

(a) $I_n = \int \sec^n x \, dx = \int \sec^{n-2} x \sec^2 x \, dx$

Let $u = \sec^{n-2} x$, $\frac{dv}{dx} = \sec^2 x \Rightarrow \frac{du}{dx} = (n-2)\sec^{n-3} x \sec x \tan x$, $v = \tan x$

$I_n = \sec^{n-2} x \tan x - \int (n-2)\sec^{n-2} x \tan^2 x \, dx$

$= \sec^{n-2} x \tan x - (n-2)\int \sec^{n-2} x(\sec^2 x - 1) \, dx$

$= \sec^{n-2} x \tan x - (n-2)\int \sec^n x \, dx + (n-2)\int \sec^{n-2} x \, dx$

$I_n = \tan x \sec^{n-2} x - (n-2)I_n + (n-2)I_{n-2}$

$I_n + (n-2)I_n = \tan x \sec^{n-2} x + (n-2)I_{n-2}$

$(n-1)I_n = \tan x \sec^{n-2} x + (n-2)I_{n-2}$ as required

(b) Let $I_n = \int_0^{\pi/4} \sec^n x \, dx$. $\tan\frac{\pi}{4} = 1$, $\sec\frac{\pi}{4} = \sqrt{2}$

$I_2 = \int_0^{\pi/4} \sec^2 x \, dx = \left[\tan x\right]_0^{\pi/4} = 1 - 0 = 1$

$n = 4$: $3I_4 = \left[\tan x \sec^2 x\right]_0^{\pi/4} + 2I_2 = 1 \times (\sqrt{2})^2 - 0 + 2(1) = 4$

$I_4 = \frac{4}{3}$

$n = 6$: $5I_6 = \left[\tan x \sec^4 x\right]_0^{\pi/4} + 4I_4 = 1 \times (\sqrt{2})^4 - 0 + 4 \times \frac{4}{3}$

$5I_6 = 4 + \frac{16}{3} = \frac{28}{3} \Rightarrow I_6 = \frac{28}{15}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 5 continued

$$\int_0^{\pi/4} \sec^6 x \, dx = \frac{28}{15}$$

Leave blank

Question 5 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 5 continued

(Total 10 marks)

Q5

Leave blank

6. The line l_1 has equation

$$\mathbf{r} = \mathbf{i} + \mathbf{j} + \mathbf{k} + \lambda(\mathbf{i} + 3\mathbf{k})$$

and the line l_2 has equation

$$\mathbf{r} = 2\mathbf{i} + s\mathbf{j} + \mu(\mathbf{i} - 2\mathbf{j} + \mathbf{k})$$

where s is a constant and λ and μ are scalar parameters.

Given that l_1 and l_2 both lie in a common plane Π_1

(a) show that an equation for Π_1 is $3x + y - z = 3$

(4)

(b) find the value of s.

(1)

The plane Π_2 has equation $\mathbf{r}.(\mathbf{i} + \mathbf{j} - 2\mathbf{k}) = 3$

(c) Find an equation for the line of intersection of Π_1 and Π_2

(4)

(d) Find the acute angle between Π_1 and Π_2 giving your answer in degrees to 3 significant figures.

(4)

(a) $n = (i + 3k) \times (i - 2j + k)$

$$n = \begin{vmatrix} i & j & k \\ 1 & 0 & 3 \\ 1 & -2 & 1 \end{vmatrix} = (0 + 6)i - (1 - 3)j + (-2 - 0)k$$

$n = 6i + 2j - 2k$, or use $3i + j - k$

$(1, 1, 1)$ on l_1: $3(1) + 1(1) - 1(1) = 3$

$\therefore$ Π_1: $3x + y - z = 3$ as required

(b) $(2, s, 0)$ lies on Π_1: $3(2) + s - 0 = 3 \Rightarrow s = -3$

(c) Direction of line $= (3i + j - k) \times (i + j - 2k)$

$$= \begin{vmatrix} i & j & k \\ 3 & 1 & -1 \\ 1 & 1 & -2 \end{vmatrix} = (-2 + 1)i - (-6 + 1)j + (3 - 1)k$$

$= -i + 5j + 2k$, so direction $i - 5j - 2k$

Π_2: $x + y - 2z = 3$. Let $x = 0$:

$y - z = 3$ and $y - 2z = 3 \Rightarrow z = 0$, $y = 3$, point $(0, 3, 0)$

$r = 3j + \lambda(i - 5j - 2k)$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 6 continued

(d) $(3\mathbf{i} + \mathbf{j} - \mathbf{k}).(\mathbf{i} + \mathbf{j} - 2\mathbf{k}) = 3 + 1 + 2 = 6$

$$\cos\theta = \frac{6}{\sqrt{(9+1+1)}\sqrt{(1+1+4)}} = \frac{6}{\sqrt{11}\sqrt{6}} = \frac{\sqrt{6}}{\sqrt{11}}$$

$\theta = 42.392...^\circ = 42.4^\circ$ (3 s.f.)

Leave blank

Question 6 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 6 continued

(Total 13 marks)

Q6

Leave blank

7. Using calculus, find the exact values of

(i) $\int_1^2 \frac{1}{x^2 - 4x + 5}\,\mathrm{d}x$

(3)

(ii) $\int_{\sqrt{3}}^3 \frac{\sqrt{x^2 - 3}}{x^2}\,\mathrm{d}x$

(5)

(i) $x^2 - 4x + 5 = (x - 2)^2 + 1$

$$\int_1^2 \frac{1}{(x-2)^2 + 1}\,dx = \Big[\arctan(x - 2)\Big]_1^2$$

$$= \arctan 0 - \arctan(-1) = 0 - \left(-\frac{\pi}{4}\right) = \frac{\pi}{4}$$

(ii) $u = \sqrt{(x^2 - 3)}$, $\frac{du}{dx} = \frac{x}{\sqrt{(x^2 - 3)}}$; $\frac{dv}{dx} = \frac{1}{x^2}$, $v = -\frac{1}{x}$

$$\int \frac{\sqrt{(x^2 - 3)}}{x^2}\,dx = -\frac{\sqrt{(x^2 - 3)}}{x} + \int \frac{1}{x} \times \frac{x}{\sqrt{(x^2 - 3)}}\,dx$$

$$= -\frac{\sqrt{(x^2 - 3)}}{x} + \int \frac{1}{\sqrt{(x^2 - 3)}}\,dx = -\frac{\sqrt{(x^2 - 3)}}{x} + \operatorname{arcosh}\left(\frac{x}{\sqrt{3}}\right) + c$$

$$\int_{\sqrt{3}}^3 \frac{\sqrt{(x^2 - 3)}}{x^2}\,dx = \left[-\frac{\sqrt{(x^2 - 3)}}{x} + \operatorname{arcosh}\left(\frac{x}{\sqrt{3}}\right)\right]_{\sqrt{3}}^3$$

$$= \left(-\frac{\sqrt{6}}{3} + \operatorname{arcosh}\sqrt{3}\right) - \left(-0 + \operatorname{arcosh} 1\right)$$

$$= \operatorname{arcosh}\sqrt{3} - \frac{\sqrt{6}}{3} = \ln(\sqrt{3} + \sqrt{2}) - \frac{\sqrt{6}}{3}$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 7 continued

Leave blank

Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave blank

Question 7 continued

Q7

(Total 8 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave blank

8. The hyperbola H has equation

$$4x^2 - y^2 = 4$$

(a) Write down the equations of the asymptotes of H.

(1)

(b) Find the coordinates of the foci of H.

(2)

The point $P(\sec\theta, 2\tan\theta)$ lies on H.

(c) Using calculus, show that the equation of the tangent to H at the point P is

$$y\tan\theta = 2x\sec\theta - 2$$

(4)

The point $V(-1, 0)$ and the point $W(1, 0)$ both lie on H.

The point $Q(\sec\theta, -2\tan\theta)$ also lies on H.

Given that P, Q, V and W are distinct points on H and that the lines VP and WQ intersect at the point S,

(d) show that, as θ varies, S lies on an ellipse with equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

where a and b are integers to be found.

(7)

(a) $\frac{x^2}{1} - \frac{y^2}{4} = 1$, $a = 1$, $b = 2$. Asymptotes: $y = 2x$ and $y = -2x$

(b) $b^2 = a^2(e^2 - 1)$: $4 = 1(e^2 - 1) \Rightarrow e = \sqrt{5}$

Foci $(\pm ae, 0)$: $(\sqrt{5}, 0)$ and $(-\sqrt{5}, 0)$

(c) $8x - 2y\frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = \frac{4x}{y} = \frac{4\sec\theta}{2\tan\theta} = \frac{2\sec\theta}{\tan\theta}$

$y - 2\tan\theta = \frac{2\sec\theta}{\tan\theta}(x - \sec\theta)$

$y\tan\theta - 2\tan^2\theta = 2x\sec\theta - 2\sec^2\theta$

$y\tan\theta = 2x\sec\theta - 2(\sec^2\theta - \tan^2\theta)$

$\sec^2\theta - \tan^2\theta = 1 \Rightarrow y\tan\theta = 2x\sec\theta - 2$ as required

(d) VP: gradient $= \frac{2\tan\theta - 0}{\sec\theta + 1}$, $y = \frac{2\tan\theta}{\sec\theta + 1}(x + 1)$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 8 continued

WQ: gradient $= \dfrac{-2\tan\theta - 0}{\sec\theta - 1}$, $y = \dfrac{-2\tan\theta}{\sec\theta - 1}(x - 1)$

At S: $\dfrac{2\tan\theta}{\sec\theta + 1}(x + 1) = \dfrac{-2\tan\theta}{\sec\theta - 1}(x - 1)$

$(x + 1)(\sec\theta - 1) = -(x - 1)(\sec\theta + 1)$

$x\sec\theta - x + \sec\theta - 1 = -x\sec\theta - x + \sec\theta + 1$

$2x\sec\theta = 2 \quad \Rightarrow \quad x = \cos\theta$

$y = \dfrac{2\tan\theta}{\sec\theta + 1}(\cos\theta + 1) = \dfrac{2\tan\theta\cos\theta(\cos\theta + 1)}{1 + \cos\theta} = 2\sin\theta$

$x = \cos\theta$, $y = 2\sin\theta$ and $\cos^2\theta + \sin^2\theta = 1$

$\therefore \quad x^2 + \dfrac{y^2}{4} = 1$, i.e. $\dfrac{x^2}{1^2} + \dfrac{y^2}{2^2} = 1$, so $a = 1$, $b = 2$

Leave blank

Question 8 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 8 continued

Leave blank

Question 8 continued

Q8

(Total 14 marks)

TOTAL FOR PAPER: 75 MARKS

END

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA