Please check the examination details below before entering your candidate information

Candidate surname StemUp Model Solution | Other names

Pearson Edexcel
International Advanced Level

Centre Number | Candidate Number

Thursday 7 January 2021

Morning (Time: 1 hour 30 minutes) | Paper Reference **WMA14/01**

Mathematics

International Advanced Subsidiary/Advanced Level
Pure Mathematics P4

You must have:
Mathematical Formulae and Statistical Tables (Lilac), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
  – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 10 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
  – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶

P67754A

©2021 Pearson Education Ltd.

1/1/1/1/

P 6 7 7 5 4 A 0 1 2 8

Pearson

Leave blank

1. (a) Find the first 4 terms, in ascending powers of x, of the binomial expansion of

$$\left(\frac{1}{4} - 5x\right)^{\frac{1}{2}} \qquad |x| < \frac{1}{20}$$

giving each coefficient in its simplest form.

(5)

By substituting $x = \frac{1}{100}$ into the answer for (a),

(b) find an approximation for $\sqrt{5}$

Give your answer in the form $\frac{a}{b}$ where a and b are integers to be found.

(2)

(a) $\left(\frac{1}{4} - 5x\right)^{\frac{1}{2}} = \left(\frac{1}{4}\right)^{\frac{1}{2}}(1 - 20x)^{\frac{1}{2}} = \frac{1}{2}(1 - 20x)^{\frac{1}{2}}$

$(1 - 20x)^{\frac{1}{2}} = 1 + (\frac{1}{2})(-20x) + \frac{(\frac{1}{2})(-\frac{1}{2})}{2!}(-20x)^2$

$+ \frac{(\frac{1}{2})(-\frac{1}{2})(-3/2)}{3!}(-20x)^3 + \ldots$

$= 1 - 10x - 50x^2 - 500x^3 + \ldots$

$\left(\frac{1}{4} - 5x\right)^{\frac{1}{2}} = \frac{1}{2}(1 - 10x - 50x^2 - 500x^3 + \ldots)$

$= \frac{1}{2} - 5x - 25x^2 - 250x^3 + \ldots$

(b) $x = \frac{1}{100} : \left(\frac{1}{4} - \frac{5}{100}\right)^{\frac{1}{2}} = \left(\frac{1}{5}\right)^{\frac{1}{2}} = \frac{1}{\sqrt{5}} = \frac{\sqrt{5}}{5}$

$\frac{\sqrt{5}}{5} \approx \frac{1}{2} - 5\left(\frac{1}{100}\right) - 25\left(\frac{1}{100}\right)^2 - 250\left(\frac{1}{100}\right)^3$

$\frac{\sqrt{5}}{5} \approx 0.5 - 0.05 - 0.0025 - 0.00025 = 0.44725 = \frac{1789}{4000}$

$\sqrt{5} \approx 5 \times \frac{1789}{4000} = \frac{1789}{800}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 1 continued

Q1

(Total 7 marks)

Leave blank

2.




Figure 1

Figure 1 shows a sketch of parallelogram $ABCD$.

Given that $\overrightarrow{AB} = 6\mathbf{i} - 2\mathbf{j} + 3\mathbf{k}$ and $\overrightarrow{BC} = 2\mathbf{i} + 5\mathbf{j} + 8\mathbf{k}$

(a) find the size of angle ABC, giving your answer in degrees, to 2 decimal places.

(3)

(b) Find the area of parallelogram $ABCD$, giving your answer to one decimal place.

(2)

(a) $\overrightarrow{BA} = -6\mathbf{i} + 2\mathbf{j} - 3\mathbf{k}$, $\overrightarrow{BC} = 2\mathbf{i} + 5\mathbf{j} + 8\mathbf{k}$

$\overrightarrow{BA} \cdot \overrightarrow{BC} = (-6)(2) + (2)(5) + (-3)(8) = -12 + 10 - 24 = -26$

$|\overrightarrow{BA}| = \sqrt{(6^2 + 2^2 + 3^2)} = \sqrt{49} = 7$, $|\overrightarrow{BC}| = \sqrt{(2^2 + 5^2 + 8^2)} = \sqrt{93}$

$\cos ABC = \dfrac{\overrightarrow{BA} \cdot \overrightarrow{BC}}{|\overrightarrow{BA}||\overrightarrow{BC}|} = \dfrac{-26}{7\sqrt{93}} = -0.38515\ldots$

$ABC = 112.6532\ldots^\circ = 112.65^\circ$

(b) Area $= |\overrightarrow{BA}||\overrightarrow{BC}| \sin ABC$

$= 7\sqrt{93} \times \sin 112.6532\ldots^\circ$

$= 62.2976\ldots = 62.3$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 2 continued

Q2

(Total 5 marks)

Leave blank

3. Prove by contradiction that there is no greatest odd integer.

(2)

Assume that there is a greatest odd integer, n.
Then n + 2 is also an odd integer (odd + even = odd)
and n + 2 > n.
So n + 2 is an odd integer greater than the greatest odd
integer n. This is a contradiction.
Hence there is no greatest odd integer.

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 3 continued

Q3

(Total 2 marks)

Leave blank

4. The curve C is defined by the parametric equations

$$x = \frac{1}{t} + 2 \qquad y = \frac{1 - 2t}{3 + t} \qquad t > 0$$

(a) Show that the equation of C can be written in the form $y = g(x)$ where g is the function

$$g(x) = \frac{ax + b}{cx + d} \qquad x > k$$

where a, b, c, d and k are integers to be found.

(5)

(b) Hence, or otherwise, state the range of g.

(2)

(a) $t > 0 \Rightarrow \frac{1}{t} > 0 \Rightarrow x > 2$, so $k = 2$

$x = \frac{1}{t} + 2 \Rightarrow \frac{1}{t} = x - 2 \Rightarrow t = \frac{1}{x - 2}$

$y = \frac{1 - 2t}{3 + t} = \frac{1 - 2/(x - 2)}{3 + 1/(x - 2)}$ ($x(x - 2)$ top and bottom)

$y = \frac{(x - 2) - 2}{3(x - 2) + 1} = \frac{x - 4}{3x - 6 + 1}$

$g(x) = \frac{x - 4}{3x - 5}$, $x > 2$

$a = 1, b = -4, c = 3, d = -5, k = 2$

(b) $g(2) = \frac{2 - 4}{6 - 5} = -2$; as $x \to \infty$, $g(x) \to \frac{1}{3}$

$g'(x) = \frac{(3x - 5) - 3(x - 4)}{(3x - 5)^2} = \frac{7}{(3x - 5)^2} > 0$, g increasing

Range: $-2 < g(x) < \frac{1}{3}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 4 continued

Q4

(Total 7 marks)

Leave blank

5. **In this question you should show all stages of your working.**

Solutions relying on calculator technology are not acceptable.

Using the substitution $u = 3 + \sqrt{2x-1}$ find the exact value of

$$\int_{1}^{13} \frac{4}{3+\sqrt{2x-1}}\,\mathrm{d}x$$

giving your answer in the form $p + q\ln 2$, where p and q are integers to be found.

(8)

$u = 3 + \sqrt{(2x-1)} \Rightarrow (u-3)^2 = 2x - 1$

$x = \frac{(u-3)^2 + 1}{2} \Rightarrow \frac{dx}{du} = u - 3$

$x = 1 : \ u = 3 + \sqrt{1} = 4 ; \quad x = 13 : \ u = 3 + \sqrt{25} = 8$

$\int_1^{13} \frac{4}{3+\sqrt{(2x-1)}}\,dx = \int_4^8 \frac{4}{u} \times (u-3)\,du$

$= \int_4^8 \left(4 - \frac{12}{u}\right) du$

$= \left[4u - 12\ln u\right]_4^8$

$= (4 \times 8 - 12\ln 8) - (4 \times 4 - 12\ln 4)$

$= 32 - 12\ln 8 - 16 + 12\ln 4$

$= 16 - 36\ln 2 + 24\ln 2$

$= 16 - 12\ln 2 \qquad (p = 16, q = -12)$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 5 continued

Q5

(Total 8 marks)

Leave blank

6. A curve has equation

$$4y^2 + 3x = 6ye^{-2x}$$

(a) Find $\frac{dy}{dx}$ in terms of x and y.

(5)

The curve crosses the y-axis at the origin and at the point P.

(b) Find the equation of the normal to the curve at P, writing your answer in the form $y = mx + c$ where m and c are constants to be found.

(4)

(a) $8y\frac{dy}{dx} + 3 = -12ye^{-2x} + 6e^{-2x}\frac{dy}{dx}$

$\frac{dy}{dx}(8y - 6e^{-2x}) = -12ye^{-2x} - 3$

$\frac{dy}{dx} = \frac{12ye^{-2x} + 3}{6e^{-2x} - 8y}$

(b) $x = 0$: $4y^2 = 6y \Rightarrow 2y(2y - 3) = 0 \Rightarrow y = 0$ or $y = 3/2$

P is $\left(0, \frac{3}{2}\right)$

$\frac{dy}{dx} = \frac{12(3/2)(1) + 3}{6(1) - 8(3/2)} = \frac{21}{-6} = -\frac{7}{2}$

Gradient of normal $= -1 \div \left(-\frac{7}{2}\right) = \frac{2}{7}$

$y - \frac{3}{2} = \frac{2}{7}(x - 0)$

$y = \frac{2}{7}x + \frac{3}{2}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 6 continued

Q6

(Total 9 marks)

Leave blank

7.




Figure 2

(a) Find $\int e^{2x} \sin x \, dx$

(5)

Figure 2 shows a sketch of part of the curve with equation

$$y = e^{2x} \sin x \qquad x \geqslant 0$$

The finite region R is bounded by the curve and the x-axis and is shown shaded in Figure 2.

(b) Show that the exact area of R is $\dfrac{e^{2\pi} + 1}{5}$

(*Solutions relying on calculator technology are not acceptable.*)

(2)

(a) Let $I = \int e^{2x} \sin x \, dx$; $u = \sin x$, $dv/dx = e^{2x}$

$I = \frac{1}{2} e^{2x} \sin x - \int \frac{1}{2} e^{2x} \cos x \, dx$

$\int \frac{1}{2} e^{2x} \cos x \, dx = \frac{1}{4} e^{2x} \cos x + \int \frac{1}{4} e^{2x} \sin x \, dx$

$I = \frac{1}{2} e^{2x} \sin x - \frac{1}{4} e^{2x} \cos x - \frac{1}{4} I$

$\frac{5}{4} I = \frac{1}{2} e^{2x} \sin x - \frac{1}{4} e^{2x} \cos x$

$\int e^{2x} \sin x \, dx = \frac{2}{5} e^{2x} \sin x - \frac{1}{5} e^{2x} \cos x + c$

(b) Curve meets x-axis when $\sin x = 0$: $x = 0$ and $x = \pi$

Area $= \left[\frac{2}{5} e^{2x} \sin x - \frac{1}{5} e^{2x} \cos x \right]_0^{\pi}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 7 continued

$$= \left(\frac{2}{5}e^{2\pi}\sin\pi - \frac{1}{5}e^{2\pi}\cos\pi\right) - \left(\frac{2}{5}e^{0}\sin 0 - \frac{1}{5}e^{0}\cos 0\right)$$

$$= \left(0 + \frac{1}{5}e^{2\pi}\right) - \left(0 - \frac{1}{5}\right) = \frac{1}{5}e^{2\pi} + \frac{1}{5} = \frac{e^{2\pi}+1}{5} \quad \text{as required}$$

Leave blank

Question 7 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 7 continued

(Total 7 marks)

Q7

Leave blank

8. With respect to a fixed origin O, the lines l_1 and l_2 are given by the equations

$$l_1: \quad \mathbf{r} = \begin{pmatrix} -1 \\ 5 \\ 4 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ -1 \\ 5 \end{pmatrix} \qquad l_2: \quad \mathbf{r} = \begin{pmatrix} 2 \\ -2 \\ -5 \end{pmatrix} + \mu \begin{pmatrix} 4 \\ -3 \\ b \end{pmatrix}$$

where λ and μ are scalar parameters and b is a constant.

Prove that for all values of $b \neq 7$, the lines l_1 and l_2 are skew.

(6)

If the lines meet:

$i: \quad -1 + 2\lambda = 2 + 4\mu \qquad (1)$

$j: \quad 5 - \lambda = -2 - 3\mu \qquad (2)$

$k: \quad 4 + 5\lambda = -5 + b\mu \qquad (3)$

$(1) + 2\times(2): \quad 9 = -2 - 2\mu \Rightarrow \mu = -11/2$

$(2): \quad \lambda = 7 + 3\mu = 7 - 33/2 \Rightarrow \lambda = -19/2$

Check (1): $-1 + 2(-19/2) = -20$, $2 + 4(-11/2) = -20$ ✓

$(3): \quad 4 + 5\left(-\frac{19}{2}\right) = -5 + b\left(-\frac{11}{2}\right)$

$-\frac{87}{2} = -5 - \frac{11b}{2} \Rightarrow -87 = -10 - 11b \Rightarrow 11b = 77 \Rightarrow b = 7$

So the lines intersect only when $b = 7$; when $b \neq 7$ equation (3) is not satisfied, so the lines do not intersect.

Directions $2i - j + 5k$ and $4i - 3j + bk$: $4/2 = 2$ but $-3/-1 = 3$, so they are not multiples and the lines are not parallel for any b.

$\therefore$ for $b \neq 7$ the lines neither meet nor are parallel, so l_1 and l_2 are skew.

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 8 continued

Q8

(Total 6 marks)

Leave blank

9.




Figure 3

Figure 3 shows a sketch of part of the curve with parametric equations

$$x = \tan\theta \qquad y = 2\sin 2\theta \qquad \theta \geqslant 0$$

The finite region, shown shaded in Figure 3, is bounded by the curve, the x-axis and the line with equation $x = \sqrt{3}$

The region is rotated through 2π radians about the x-axis to form a solid of revolution.

(a) Show that the exact volume of this solid of revolution is given by

$$\int_0^k p(1 - \cos 2\theta)\,\mathrm{d}\theta$$

where p and k are constants to be found.

(7)

(b) Hence find, by algebraic integration, the exact volume of this solid of revolution.

(3)

(a) $x = 0 \Rightarrow \theta = 0$; $x = \sqrt{3} \Rightarrow \tan\theta = \sqrt{3} \Rightarrow \theta = \frac{\pi}{3}$ so $k = \frac{\pi}{3}$

$x = \tan\theta \Rightarrow \frac{dx}{d\theta} = \sec^2\theta$

$V = \pi \int y^2\,dx = \pi \int_0^{\pi/3} (2\sin 2\theta)^2 \sec^2\theta\,d\theta$

$= \pi \int_0^{\pi/3} 4(2\sin\theta\cos\theta)^2 \times \frac{1}{\cos^2\theta}\,d\theta$

$= \pi \int_0^{\pi/3} 16\sin^2\theta\,d\theta$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 9 continued

$= 16\pi \int_0^{\pi/3} \frac{1 - \cos 2\theta}{2}\, d\theta \quad \left(\sin^2\theta = \frac{1 - \cos 2\theta}{2}\right)$

$V = \int_0^{\pi/3} 8\pi(1 - \cos 2\theta)\, d\theta \qquad p = 8\pi, \quad k = \frac{\pi}{3}$

(b) $V = 8\pi \left[\theta - \frac{1}{2}\sin 2\theta\right]_0^{\pi/3}$

$= 8\pi\left(\frac{\pi}{3} - \frac{1}{2}\sin\frac{2\pi}{3}\right) - 8\pi(0 - 0)$

$= 8\pi\left(\frac{\pi}{3} - \frac{\sqrt{3}}{4}\right)$

$V = \frac{8}{3}\pi^2 - 2\sqrt{3}\,\pi$

Leave blank

Question 9 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 9 continued

(Total 10 marks)

Q9

Leave blank

10. (a) Write $\dfrac{1}{(H-5)(H+3)}$ in partial fraction form.

(3)

The depth of water in a storage tank is being monitored.

The depth of water in the tank, H metres, is modelled by the differential equation

$$\frac{\mathrm{d}H}{\mathrm{d}t} = -\frac{(H-5)(H+3)}{40}$$

where t is the time, in days, from when monitoring began.

Given that the initial depth of water in the tank was 13 m,

(b) solve the differential equation to show that

$$H = \frac{10 + 3\mathrm{e}^{-0.2t}}{2 - \mathrm{e}^{-0.2t}}$$

(7)

(c) Hence find the time taken for the depth of water in the tank to fall to 8 m.

(*Solutions relying entirely on calculator technology are not acceptable.*)

(3)

According to the model, the depth of water in the tank will eventually fall to k metres.

(d) State the value of the constant k.

(1)

(a) $\dfrac{1}{(H-5)(H+3)} = \dfrac{A}{H-5} + \dfrac{B}{H+3} \Rightarrow 1 = A(H+3) + B(H-5)$

$H = 5: \ 1 = 8A \Rightarrow A = \frac{1}{8}; \quad H = -3: \ 1 = -8B \Rightarrow B = -\frac{1}{8}$

$\dfrac{1}{(H-5)(H+3)} = \dfrac{1}{8(H-5)} - \dfrac{1}{8(H+3)}$

(b) $\displaystyle\int \frac{1}{(H-5)(H+3)}\,dH = \int -\frac{1}{40}\,dt$

$\displaystyle\int \left(\frac{1}{8(H-5)} - \frac{1}{8(H+3)}\right) dH = -\frac{1}{40}\int 1\,dt$

$\frac{1}{8}\ln|H-5| - \frac{1}{8}\ln|H+3| = -\frac{1}{40}t + c$

$t = 0, H = 13: \ \frac{1}{8}\ln 8 - \frac{1}{8}\ln 16 = c \Rightarrow c = \frac{1}{8}\ln\left(\frac{1}{2}\right)$

$\times 8: \ \ln|H-5| - \ln|H+3| = -\frac{1}{5}t + \ln\left(\frac{1}{2}\right)$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 10 continued

$\ln\left(\frac{H-5}{H+3}\right) = -0.2t + \ln\frac{1}{2} \Rightarrow \frac{H-5}{H+3} = \frac{1}{2}e^{-0.2t}$

$2(H-5) = (H+3)e^{-0.2t}$

$2H - 10 = He^{-0.2t} + 3e^{-0.2t}$

$2H - He^{-0.2t} = 10 + 3e^{-0.2t}$

$H(2 - e^{-0.2t}) = 10 + 3e^{-0.2t}$

$H = \frac{10 + 3e^{-0.2t}}{2 - e^{-0.2t}}$ as required

(c) $\frac{10 + 3e^{-0.2t}}{2 - e^{-0.2t}} = 8 \Rightarrow 10 + 3e^{-0.2t} = 16 - 8e^{-0.2t}$

$11e^{-0.2t} = 6 \Rightarrow e^{-0.2t} = \frac{6}{11}$

$-0.2t = \ln\left(\frac{6}{11}\right) \Rightarrow t = -5\ln\left(\frac{6}{11}\right) = 5\ln\left(\frac{11}{6}\right)$

$t = 3.0306\ldots = 3.03$ days

(d) As $t \to \infty$, $e^{-0.2t} \to 0$, so $H \to 10/2 = 5$

$k = 5$

Leave blank

Question 10 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 10 continued

Leave blank

Question 10 continued

Q10

(Total 14 marks)

TOTAL FOR PAPER IS 75 MARKS

END

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA