Please check the examination details below before entering your candidate information

Candidate surname StemUp Model Solution | Other names

Centre Number | Candidate Number

Pearson Edexcel International Advanced Level

Time 1 hour 30 minutes

Paper reference **WFM01/01**

Mathematics

International Advanced Subsidiary/ Advanced Level
Further Pure Mathematics F1

You must have:
Mathematical Formulae and Statistics Tables (Yellow), calculator

Total Marks

Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
  – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 9 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
  – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶

P72464A

©2022 Pearson Education Ltd.
J:1/1/1/1/

P 7 2 4 6 4 A 0 1 3 2

Pearson

1. Given that

$$\mathbf{A} = \begin{pmatrix} 2 & -1 & 3 \\ -2 & 3 & 0 \end{pmatrix} \quad \text{and} \quad \mathbf{B} = \begin{pmatrix} 1 & k \\ 0 & -3 \\ 2k & 2 \end{pmatrix}$$

where k is a non-zero constant,

(a) determine the matrix **AB**

(2)

(b) determine the value of k for which $\det(\mathbf{AB}) = 0$

(3)

(a) $\begin{pmatrix} 2 & -1 & 3 \\ -2 & 3 & 0 \end{pmatrix} \begin{pmatrix} 1 & k \\ 0 & -3 \\ 2k & 2 \end{pmatrix} = \begin{pmatrix} 2+0+6k & 2k+3+6 \\ -2+0+0 & -2k-9+0 \end{pmatrix}$

$AB = \begin{pmatrix} 2+6k & 2k+9 \\ -2 & -2k-9 \end{pmatrix}$

(b) $\det AB = (2+6k)(-2k-9) - (-2)(2k+9)$

$= -4k - 18 - 12k^2 - 54k + 4k + 18$

$= -12k^2 - 54k$

$\det AB = 0 \Rightarrow -6k(2k+9) = 0$

$k \neq 0 \Rightarrow k = -\frac{9}{2}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 1 continued

(Total for Question 1 is 5 marks)

2. **In this question you must show all stages of your working.**

Solutions relying entirely on calculator technology are not acceptable.

Use the standard results for $\sum_{r=1}^{n} r$ and $\sum_{r=1}^{n} r^2$ to show that for all positive integers n

$$\sum_{r=1}^{n} (7r-5)^2 = \frac{n}{6}(7n+1)(An+B)$$

where A and B are integers to be determined.

(6)

$(7r-5)^2 = 49r^2 - 70r + 25$

$\sum_{r=1}^{n} (7r-5)^2 = 49\sum_{r=1}^{n} r^2 - 70\sum_{r=1}^{n} r + \sum_{r=1}^{n} 25$

$= 49 \times \frac{n}{6}(n+1)(2n+1) - 70 \times \frac{n}{2}(n+1) + 25n$

$= \frac{n}{6}\left[49(2n^2+3n+1) - 210(n+1) + 150\right]$

$= \frac{n}{6}(98n^2 + 147n + 49 - 210n - 210 + 150)$

$= \frac{n}{6}(98n^2 - 63n - 11)$

$= \frac{n}{6}(7n+1)(14n-11)$

so $A = 14, \quad B = -11$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 2 continued

(Total for Question 2 is 6 marks)

3. **In this question you must show all stages of your working.**

Solutions relying entirely on calculator technology are not acceptable.

$$f(z) = 4z^3 + pz^2 - 24z + 108$$

where p is a constant.

Given that -3 is a root of the equation $f(z) = 0$

(a) determine the value of p

(2)

(b) using algebra, solve $f(z) = 0$ completely, giving the roots in simplest form,

(4)

(c) determine the modulus of the complex roots of $f(z) = 0$

(2)

(d) show the roots of $f(z) = 0$ on a single Argand diagram.

(2)

(a) $f(-3) = 0 \Rightarrow 4(-3)^3 + p(-3)^2 - 24(-3) + 108 = 0$

$-108 + 9p + 72 + 108 = 0 \Rightarrow 9p = -72$

$p = -8$

(b) $4z^3 - 8z^2 - 24z + 108 = (z + 3)(4z^2 - 20z + 36)$

$4z^2 - 20z + 36 = 0 \Rightarrow z^2 - 5z + 9 = 0$

$z = \frac{5 \pm \sqrt{(25 - 36)}}{2} = \frac{5 \pm \sqrt{(-11)}}{2} = \frac{5 \pm i\sqrt{11}}{2}$

Roots: $z = -3$, $z = \frac{5 + i\sqrt{11}}{2}$, $z = \frac{5 - i\sqrt{11}}{2}$

(c) $|z|^2 = \left(\frac{5}{2}\right)^2 + \left(\frac{\sqrt{11}}{2}\right)^2 = \frac{25}{4} + \frac{11}{4} = 9 \Rightarrow |z| = \sqrt{9}$

Modulus = 3

(d) Argand diagram:


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 3 continued



Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 3 continued

(Total for Question 3 is 10 marks)

4.

$$f(x) = 1 - \frac{1}{8x^4} + \frac{2}{7\sqrt{x^7}} \qquad x > 0$$

The equation $f(x) = 0$ has a single root, α, that lies in the interval $[0.15, 0.25]$

(a) (i) Determine $f'(x)$

(ii) Explain why 0.25 cannot be used as an initial approximation for α in the Newton-Raphson process.

(iii) Taking 0.15 as a first approximation to α apply the Newton-Raphson process once to $f(x)$ to obtain a second approximation to α
Give your answer to 3 decimal places.

(5)

(b) Use linear interpolation once on the interval $[0.15, 0.25]$ to find another approximation to α
Give your answer to 3 decimal places.

(3)

(a)(i) $f(x) = 1 - \frac{1}{8}x^{-4} + \frac{2}{7}x^{-7/2}$

$f'(x) = \frac{4}{8}x^{-5} - \frac{7}{2} \times \frac{2}{7}x^{-9/2} = \frac{1}{2x^5} - \frac{1}{x^{9/2}}$

(ii) $f'(0.25) = \frac{1}{2(0.25)^5} - \frac{1}{(0.25)^{9/2}} = 512 - 512 = 0$

The Newton-Raphson process would require division by $f'(0.25) = 0$, so it cannot be used.

(iii) $f(0.15) = -27.3325...$, $f'(0.15) = 1484.137...$

$\alpha_2 = 0.15 - \frac{f(0.15)}{f'(0.15)} = 0.15 - \frac{-27.3325...}{1484.137...}$

$= 0.168416...$

$\alpha_2 = 0.168$ (3 d.p.)

(b) $f(0.15) = -27.3325...$, $f(0.25) = 5.5714...$

$\frac{\alpha - 0.15}{0 - f(0.15)} = \frac{0.25 - \alpha}{f(0.25) - 0} \Rightarrow \frac{\alpha - 0.15}{27.3325...} = \frac{0.25 - \alpha}{5.5714...}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

$$\alpha = 0.15 + \frac{0.1 \times 27.3325...}{5.5714... + 27.3325...} = 0.233067...$$

$\alpha = 0.233$ (3 d.p.)

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

(Total for Question 4 is 8 marks)

5. The quadratic equation

$$4x^2 + 3x + k = 0$$

where k is an integer, has roots α and β

(a) Write down, in terms of k where appropriate, the value of $\alpha + \beta$ and the value of $\alpha\beta$

(2)

(b) Determine, in simplest form in terms of k, the value of $\dfrac{\alpha}{\beta^2} + \dfrac{\beta}{\alpha^2}$

(4)

(c) Determine a quadratic equation which has roots

$$\frac{\alpha}{\beta^2} \text{ and } \frac{\beta}{\alpha^2}$$

giving your answer in the form $px^2 + qx + r = 0$ where p, q and r are integer values in terms of k

(3)

(a) $\alpha + \beta = -\frac{3}{4}$, $\quad \alpha\beta = \frac{k}{4}$

(b) $\frac{\alpha}{\beta^2} + \frac{\beta}{\alpha^2} = \frac{\alpha^3 + \beta^3}{\alpha^2\beta^2} = \frac{(\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)}{(\alpha\beta)^2}$

$= \frac{(-3/4)^3 - 3(k/4)(-3/4)}{(k/4)^2} = \frac{-27/64 + 9k/16}{k^2/16}$

$= \frac{36k - 27}{64} \times \frac{16}{k^2} = \frac{36k - 27}{4k^2}$

(c) Sum of roots $= \frac{36k - 27}{4k^2}$

Product of roots $= \frac{\alpha}{\beta^2} \times \frac{\beta}{\alpha^2} = \frac{1}{\alpha\beta} = \frac{4}{k}$

$x^2 - \left(\frac{36k - 27}{4k^2}\right)x + \frac{4}{k} = 0$

$4k^2x^2 - (36k - 27)x + 16k = 0$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 5 continued

Question 5 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 5 continued

(Total for Question 5 is 9 marks)

6. **In this question you must show all stages of your working.**

Solutions relying entirely on calculator technology are not acceptable.

The rectangular hyperbola H has equation $xy = 20$

The point $P\left(2t\sqrt{a}, \frac{2\sqrt{a}}{t}\right)$, $t \neq 0$, where a is a constant, is a general point on H

(a) State the value of a

(1)

(b) Show that the normal to H at the point P has equation

$$ty - t^3x - 2\sqrt{5}\left(1 - t^4\right) = 0$$

(4)

The points A and B lie on H

The point A has parameter $t = c$ and the point B has parameter $t = -\frac{1}{2c}$, where c is a constant.

The normal to H at A meets H again at B

(c) Determine the possible values of c

(4)

(a) $xy = 2t\sqrt{a} \times \frac{2\sqrt{a}}{t} = 4a = 20 \Rightarrow a = 5$

(b) $y = \frac{20}{x} \Rightarrow \frac{dy}{dx} = -\frac{20}{x^2}$; at P, $x = 2t\sqrt{5}$: $\frac{dy}{dx} = -\frac{20}{20t^2} = -\frac{1}{t^2}$

Gradient of normal $= \frac{-1}{-1/t^2} = t^2$

Normal: $y - \frac{2\sqrt{5}}{t} = t^2(x - 2t\sqrt{5})$

$ty - 2\sqrt{5} = t^3x - 2t^4\sqrt{5}$

$ty - t^3x - 2\sqrt{5} + 2\sqrt{5}t^4 = 0$

$ty - t^3x - 2\sqrt{5}(1 - t^4) = 0$ as required

(c) Normal at A ($t = c$): $cy - c^3x - 2\sqrt{5}(1 - c^4) = 0$

B: $t = -\frac{1}{2c} \Rightarrow x = 2t\sqrt{5} = -\frac{\sqrt{5}}{c}$, $y = \frac{2\sqrt{5}}{t} = -4c\sqrt{5}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 6 continued

$c(-4c\sqrt{5}) - c^3\left(-\frac{\sqrt{5}}{c}\right) - 2\sqrt{5}(1 - c^4) = 0$

$-4c^2\sqrt{5} + c^2\sqrt{5} - 2\sqrt{5} + 2\sqrt{5}c^4 = 0$

$2c^4 - 3c^2 - 2 = 0$

$(2c^2 + 1)(c^2 - 2) = 0 \Rightarrow c^2 = 2 \text{ or } c^2 = -\frac{1}{2}$

$c^2 > 0 \Rightarrow c^2 = 2$

$c = \sqrt{2} \text{ or } c = -\sqrt{2}$

Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 6 continued

(Total for Question 6 is 9 marks)

7. (i)

$$\mathbf{P} = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$$

The matrix **P** represents a geometrical transformation U

(a) Describe U fully as a single geometrical transformation.

(2)

The transformation V, represented by the 2×2 matrix **Q**, is a rotation through 240° anticlockwise about the origin followed by an enlargement about (0, 0) with scale factor 6

(b) Determine the matrix **Q**, giving each entry in exact numerical form.

(2)

Given that U followed by V is the transformation T, which is represented by the matrix **R**

(c) determine the matrix **R**

(2)

(ii) The transformation W is represented by the matrix

$$\begin{pmatrix} -2 & 2\sqrt{3} \\ 2\sqrt{3} & 2 \end{pmatrix}$$

Show that there is a real number λ for which W maps the point $(\lambda, 1)$ onto the point $(4\lambda, 4)$, giving the exact value of λ

(5)

(i)(a) Reflection in the line $y = -x$

(b) $Q = 6\begin{pmatrix} \cos 240° & -\sin 240° \\ \sin 240° & \cos 240° \end{pmatrix} = 6\begin{pmatrix} -1/2 & \sqrt{3}/2 \\ -\sqrt{3}/2 & -1/2 \end{pmatrix}$

$Q = \begin{pmatrix} -3 & 3\sqrt{3} \\ -3\sqrt{3} & -3 \end{pmatrix}$

(c) $R = QP = \begin{pmatrix} -3 & 3\sqrt{3} \\ -3\sqrt{3} & -3 \end{pmatrix}\begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 7 continued

$$R = \begin{pmatrix} -3\sqrt{3} & 3 \\ 3 & 3\sqrt{3} \end{pmatrix}$$

(ii) $\begin{pmatrix} -2 & 2\sqrt{3} \\ 2\sqrt{3} & 2 \end{pmatrix}\begin{pmatrix} \lambda \\ 1 \end{pmatrix} = \begin{pmatrix} 4\lambda \\ 4 \end{pmatrix} \Rightarrow \begin{pmatrix} -2\lambda + 2\sqrt{3} \\ 2\sqrt{3}\lambda + 2 \end{pmatrix} = \begin{pmatrix} 4\lambda \\ 4 \end{pmatrix}$

$-2\lambda + 2\sqrt{3} = 4\lambda \Rightarrow 6\lambda = 2\sqrt{3} \Rightarrow \lambda = \frac{\sqrt{3}}{3}$

$2\sqrt{3}\lambda + 2 = 4 \Rightarrow 2\sqrt{3}\lambda = 2 \Rightarrow \lambda = \frac{1}{\sqrt{3}} = \frac{\sqrt{3}}{3}$

Both equations give the same value, so such a real λ exists: $\lambda = \frac{\sqrt{3}}{3}$

Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 7 continued

(Total for Question 7 is 11 marks)

8. A parabola C has equation $y^2 = 4ax$ where a is a positive constant.

The point S is the focus of C

The line l_1 with equation $y = k$ where k is a positive constant, intersects C at the point P

(a) Show that

$$PS = \frac{k^2 + 4a^2}{4a}$$

(3)

The line l_2 passes through P and intersects the directrix of C on the x-axis.

The line l_2 intersects the y-axis at the point A

(b) Show that the y coordinate of A is $\dfrac{4a^2k}{k^2 + 4a^2}$

(3)

The line l_1 intersects the directrix of C at the point B

Given that the areas of triangles BPA and OSP, where O is the origin, satisfy the ratio

$$\text{area } BPA : \text{area } OSP = 4k^2 : 1$$

(c) determine the exact value of a

(5)

(a) $y = k$: $k^2 = 4ax \Rightarrow x = \frac{k^2}{4a}$, so $P\left(\frac{k^2}{4a}, k\right)$

S is $(a, 0)$ and the directrix is $x = -a$

PS = distance from P to the directrix (focus-directrix property)

$PS = \frac{k^2}{4a} + a = \frac{k^2 + 4a^2}{4a}$ as required

(b) l_2 passes through P and $(-a, 0)$

Gradient of $l_2 = \frac{k - 0}{k^2/4a + a} = \frac{k}{(k^2 + 4a^2)/4a} = \frac{4ak}{k^2 + 4a^2}$

l_2: $y = \frac{4ak}{k^2 + 4a^2}(x + a)$

At A, $x = 0$: $y = \frac{4ak}{k^2 + 4a^2} \times a = \frac{4a^2k}{k^2 + 4a^2}$ as required

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 8 continued

(c) B is $(-a, k)$. Area $OSP = \frac{1}{2} \times a \times k = \frac{ak}{2}$

$BP = \frac{k^2}{4a} + a = \frac{k^2 + 4a^2}{4a}$, height of A below $l_1 = k - \frac{4a^2k}{k^2 + 4a^2} = \frac{k^3}{k^2 + 4a^2}$

Area $BPA = \frac{1}{2} \times \frac{k^2 + 4a^2}{4a} \times \frac{k^3}{k^2 + 4a^2} = \frac{k^3}{8a}$

$\frac{\text{Area } BPA}{\text{Area } OSP} = \frac{k^3/8a}{ak/2} = \frac{k^2}{4a^2} = 4k^2$

$\Rightarrow 16a^2 = 1 \Rightarrow a^2 = \frac{1}{16}$

$a > 0 \Rightarrow a = \frac{1}{4}$

Question 8 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 8 continued

(Total for Question 8 is 11 marks)

9. Prove by induction that for all positive integers n

$$\sum_{r=1}^{n} \log(2r-1) = \log\left(\frac{(2n)!}{2^n n!}\right)$$

(6)

$n = 1$: LHS $= \sum_{r=1}^{1} \log(2r-1) = \log 1$, RHS $= \log\left(\frac{2!}{2^1 \times 1!}\right) = \log 1$

LHS = RHS, so the result is true for $n = 1$

Assume true for $n = k$: $\sum_{r=1}^{k} \log(2r-1) = \log\left(\frac{(2k)!}{2^k k!}\right)$

Then $\sum_{r=1}^{k+1} \log(2r-1) = \log\left(\frac{(2k)!}{2^k k!}\right) + \log(2(k+1)-1)$

$= \log\left(\frac{(2k)!}{2^k k!}\right) + \log(2k+1) = \log\left(\frac{(2k)! \times (2k+1)}{2^k k!}\right)$

$= \log\left(\frac{(2k+1)!}{2^k k!}\right) = \log\left(\frac{(2k+1)!}{2^k k!}\right) \times \frac{2k+2}{2k+2}$

$= \log\left(\frac{(2k+2)!}{2^k \times 2(k+1) \times k!}\right) = \log\left(\frac{(2k+2)!}{2^{k+1}(k+1)!}\right)$

which is the result with $n = k + 1$.

So if the result is true for $n = k$, it is true for $n = k + 1$.

As it is true for $n = 1$, by mathematical induction the result is true for all positive integers n.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 9 continued

Question 9 continued

(Total for Question 9 is 6 marks)

TOTAL FOR PAPER IS 75 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA