Please check the examination details below before entering your candidate information

Candidate surname StemUp Model Solution | Other names

Centre Number | Candidate Number

Pearson Edexcel International Advanced Level

Time 1 hour 30 minutes

Paper reference **WFM01/01**

Mathematics

International Advanced Subsidiary/Advanced Level
Further Pure Mathematics F1

You must have:
Mathematical Formulae and Statistical Tables (Yellow), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 9 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶

P73334A

©2022 Pearson Education Ltd.

Q:1/1/1/

P 7 3 3 3 4 A 0 1 3 6

Pearson

1. $z_1 = 3 + 3\text{i}$ $z_2 = p + q\text{i}$ $p, q \in \mathbb{R}$

Given that $|z_1 z_2| = 15\sqrt{2}$

(a) determine $|z_2|$

(2)

Given also that $p = -4$

(b) determine the possible values of q

(2)

(c) Show z_1 and the possible positions for z_2 on the same Argand diagram.

(2)

(a) $|z_1| = \sqrt{(3^2 + 3^2)} = \sqrt{18} = 3\sqrt{2}$

$|z_1 z_2| = |z_1||z_2|$

$3\sqrt{2}\,|z_2| = 15\sqrt{2}$


$|z_2| = \frac{15\sqrt{2}}{3\sqrt{2}} = 5$

(b) $|z_2| = 5 \Rightarrow p^2 + q^2 = 25$

$(-4)^2 + q^2 = 25 \Rightarrow q^2 = 9$

$q = 3$ or $q = -3$

(c) Argand diagram:



DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 1 continued

Question 1 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 1 continued

(Total for Question 1 is 6 marks)

2. $$f(x) = 10 - 2x - \frac{1}{2\sqrt{x}} - \frac{1}{x^3} \qquad x > 0$$

(a) Show that the equation $f(x) = 0$ has a root α in the interval [0.4, 0.5]

(2)

(b) Determine $f'(x)$.

(3)

(c) Using $x_0 = 0.5$ as a first approximation to α, apply the Newton-Raphson procedure once to $f(x)$ to find a second approximation to α, giving your answer to 3 decimal places.

(2)

The equation $f(x) = 0$ has another root β in the interval [4.8, 4.9]

(d) Use linear interpolation once on the interval [4.8, 4.9] to find an approximation to β, giving your answer to 3 decimal places.

(2)

(a) $f(0.4) = -7.2155... < 0$

$f(0.5) = 0.2928... > 0$

There is a sign change (negative to positive) and $f(x)$ is continuous on [0.4, 0.5], therefore a root α lies in the interval [0.4, 0.5]

(b) $f(x) = 10 - 2x - \frac{1}{2}x^{-1/2} - x^{-3}$

$f'(x) = -2 + \frac{1}{4}x^{-3/2} + 3x^{-4}$

(c) $x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$

$f(0.5) = 0.29289321...$, $f'(0.5) = 46.70710678...$

$x_1 = 0.5 - \frac{0.29289321...}{46.70710678...} = 0.493729...$

$x_1 = 0.494$ (3 d.p.)

(d) $f(4.8) = 0.16274...$, $f(4.9) = -0.034376...$

$\frac{\beta - 4.8}{0.16274...} = \frac{4.9 - \beta}{0.034376...}$

$\beta = \frac{4.8 \times 0.034376... + 4.9 \times 0.16274...}{0.034376... + 0.16274...}$

$\beta = 4.88256...$

$\beta = 4.883$ (3 d.p.)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 2 continued

Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 2 continued

(Total for Question 2 is 9 marks)

3. $\mathbf{M} = \begin{pmatrix} k & k \\ 3 & 5 \end{pmatrix}$ where k is a non-zero constant

(a) Determine $\mathbf{M}^{-1}$, giving your answer in simplest form in terms of k.

(2)

Hence, given that $\mathbf{N}^{-1} = \begin{pmatrix} k & k \\ 4 & -1 \end{pmatrix}$

(b) determine $(\mathbf{MN})^{-1}$, giving your answer in simplest form in terms of k.

(2)

(a) $\det M = 5k - 3k = 2k$

$$M^{-1} = \frac{1}{2k}\begin{pmatrix} 5 & -k \\ -3 & k \end{pmatrix}$$

(b) $$(MN)^{-1} = N^{-1}M^{-1} = \frac{1}{2k}\begin{pmatrix} k & k \\ 4 & -1 \end{pmatrix}\begin{pmatrix} 5 & -k \\ -3 & k \end{pmatrix}$$

$$= \frac{1}{2k}\begin{pmatrix} 5k - 3k & -k^2 + k^2 \\ 20 + 3 & -4k - k \end{pmatrix}$$

$$= \frac{1}{2k}\begin{pmatrix} 2k & 0 \\ 23 & -5k \end{pmatrix}$$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 3 continued

(Total for Question 3 is 4 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

4. $f(z) = 2z^4 - 19z^3 + Az^2 + Bz - 156$

where A and B are constants.

The complex number $5 - i$ is a root of the equation $f(z) = 0$

(a) Write down another complex root of this equation.

(1)

(b) Solve the equation $f(z) = 0$ completely.

(5)

(c) Determine the value of A and the value of B.

(2)

(a) $5 + i$

(b) $(z - (5 + i))(z - (5 - i)) = (z - 5)^2 - i^2$

$= z^2 - 10z + 25 + 1 = z^2 - 10z + 26$

$f(z) = (z^2 - 10z + 26)(2z^2 + az + b)$

constant: $26b = -156 \Rightarrow b = -6$

z^3: $-20 + a = -19 \Rightarrow a = 1$

$f(z) = (z^2 - 10z + 26)(2z^2 + z - 6)$

$2z^2 + z - 6 = 0 \Rightarrow (2z - 3)(z + 2) = 0$

$z = 5 + i, \quad z = 5 - i, \quad z = \frac{3}{2}, \quad z = -2$

(c) $(z^2 - 10z + 26)(2z^2 + z - 6)$

$= 2z^4 + z^3 - 6z^2 - 20z^3 - 10z^2 + 60z + 52z^2 + 26z - 156$

$= 2z^4 - 19z^3 + 36z^2 + 86z - 156$

$A = 36, \quad B = 86$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 4 continued

(Total for Question 4 is 8 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

5. The quadratic equation

$$2x^2 - 3x + 5 = 0$$

has roots α and β

Without solving the equation,

(a) write down the value of $(\alpha + \beta)$ and the value of $\alpha\beta$

(1)

(b) determine the value of

(i) $\alpha^2 + \beta^2$

(ii) $\alpha^3 + \beta^3$

(4)

(c) find a quadratic equation which has roots

$$(\alpha^3 - \beta) \text{ and } (\beta^3 - \alpha)$$

giving your answer in the form $px^2 + qx + r = 0$ where p, q and r are integers to be determined.

(5)

(a) $\alpha + \beta = \frac{3}{2}$, $\quad \alpha\beta = \frac{5}{2}$

(b)(i) $\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta = \left(\frac{3}{2}\right)^2 - 2\left(\frac{5}{2}\right)$

$= \frac{9}{4} - 5 = -\frac{11}{4}$

(ii) $\alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta) = \left(\frac{3}{2}\right)^3 - 3\left(\frac{5}{2}\right)\left(\frac{3}{2}\right)$

$= \frac{27}{8} - \frac{45}{4} = -\frac{63}{8}$

(c) Sum $= \alpha^3 + \beta^3 - (\alpha + \beta) = -\frac{63}{8} - \frac{3}{2} = -\frac{75}{8}$

Product $= (\alpha^3 - \beta)(\beta^3 - \alpha) = (\alpha\beta)^3 - \alpha^4 - \beta^4 + \alpha\beta$

$\alpha^4 + \beta^4 = (\alpha^2 + \beta^2)^2 - 2(\alpha\beta)^2 = \left(-\frac{11}{4}\right)^2 - 2\left(\frac{5}{2}\right)^2$

$= \frac{121}{16} - \frac{200}{16} = -\frac{79}{16}$

Product $= \left(\frac{5}{2}\right)^3 + \frac{79}{16} + \frac{5}{2} = \frac{250}{16} + \frac{79}{16} + \frac{40}{16} = \frac{369}{16}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 5 continued

$x^2 - (\text{sum})x + \text{product} = 0 \Rightarrow x^2 + \frac{75}{8}x + \frac{369}{16} = 0$

$16x^2 + 150x + 369 = 0$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 5 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 5 continued

(Total for Question 5 is 10 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

6. The parabola C has equation $y^2 = 36x$

The point $P(9t^2, 18t)$, where $t \neq 0$, lies on C

(a) Use calculus to show that the normal to C at P has equation

$$y + tx = 9t^3 + 18t$$

(4)

(b) Hence find the equations of the two normals to C which pass through the point (54, 0), giving your answers in the form $y = px + q$ where p and q are constants to be determined.

(4)

Given that

- the normals found in part (b) intersect the directrix of C at the points A and B
- the point F is the focus of C

(c) determine the area of triangle AFB

(3)

(a) $y^2 = 36x \Rightarrow 2y\frac{dy}{dx} = 36 \Rightarrow \frac{dy}{dx} = \frac{18}{y}$

At P: $\frac{dy}{dx} = \frac{18}{18t} = \frac{1}{t}$ so $m_T = \frac{1}{t}$, $m_N = -t$

Normal: $y - 18t = -t(x - 9t^2)$

$y - 18t = -tx + 9t^3$

$\Rightarrow y + tx = 9t^3 + 18t$ as required

(b) $x = 54, y = 0$: $54t = 9t^3 + 18t$

$9t^3 - 36t = 0 \Rightarrow 9t(t^2 - 4) = 0$

$t \neq 0 \Rightarrow t = 2$ or $t = -2$

$t = 2$: $y + 2x = 72 + 36 = 108 \Rightarrow y = -2x + 108$

$t = -2$: $y - 2x = -72 - 36 = -108 \Rightarrow y = 2x - 108$

(c) $4a = 36 \Rightarrow a = 9$: focus $F(9, 0)$, directrix $x = -9$

$x = -9$: $y = -2(-9) + 108 = 126 \Rightarrow A(-9, 126)$

$x = -9$: $y = 2(-9) - 108 = -126 \Rightarrow B(-9, -126)$

$AB = 126 - (-126) = 252$, height $= 9 - (-9) = 18$

Area of triangle $AFB = \frac{1}{2} \times 252 \times 18 = 2268$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 6 continued

Question 6 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 6 continued

(Total for Question 6 is 11 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

7. $$\mathbf{A} = \begin{pmatrix} -\frac{\sqrt{3}}{2} & -\frac{1}{2} \\ \frac{1}{2} & -\frac{\sqrt{3}}{2} \end{pmatrix}$$

(a) Determine the matrix $\mathbf{A}^2$

(1)

(b) Describe fully the single geometrical transformation represented by the matrix $\mathbf{A}^2$

(2)

(c) Hence determine the smallest positive integer value of n for which $\mathbf{A}^n = \mathbf{I}$

(1)

The matrix **B** represents a stretch scale factor 4 parallel to the x-axis.

(d) Write down the matrix **B**

(1)

The transformation represented by matrix **A** followed by the transformation represented by matrix **B** is represented by the matrix **C**

(e) Determine the matrix **C**

(2)

The parallelogram P is transformed onto the parallelogram P' by the matrix **C**

(f) Given that the area of parallelogram P' is 20 square units, determine the area of parallelogram P

(2)

(a) $A^2 = \begin{pmatrix} -\sqrt{3}/2 & -1/2 \\ 1/2 & -\sqrt{3}/2 \end{pmatrix}\begin{pmatrix} -\sqrt{3}/2 & -1/2 \\ 1/2 & -\sqrt{3}/2 \end{pmatrix}$

$= \begin{pmatrix} 3/4 - 1/4 & \sqrt{3}/4 + \sqrt{3}/4 \\ -\sqrt{3}/4 - \sqrt{3}/4 & -1/4 + 3/4 \end{pmatrix} = \begin{pmatrix} 1/2 & \sqrt{3}/2 \\ -\sqrt{3}/2 & 1/2 \end{pmatrix}$

(b) $\cos\theta = \frac{1}{2}$, $\sin\theta = -\frac{\sqrt{3}}{2} \Rightarrow \theta = -60°$

A^2 represents a rotation of 60° clockwise about the origin

(i.e. 300° anticlockwise about (0, 0))

(c) A^2 is a rotation of −60°, so $(A^2)^6$ = rotation of −360° = I

so $A^{12} = I$ (A is a rotation of 150° and 12 × 150° = 1800° = 5 × 360°)

$n = 12$

(d) $B = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}$

(e) $C = BA = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}\begin{pmatrix} -\sqrt{3}/2 & -1/2 \\ 1/2 & -\sqrt{3}/2 \end{pmatrix}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 7 continued

$$C = \begin{pmatrix} -2\sqrt{3} & -2 \\ 1/2 & -\sqrt{3}/2 \end{pmatrix}$$

(f) $\det C = (-2\sqrt{3})\left(-\frac{\sqrt{3}}{2}\right) - (-2)\left(\frac{1}{2}\right) = 3 + 1 = 4$

Area of $P = \frac{20}{\det C} = \frac{20}{4} = 5$ square units

Question 7 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 7 continued

(Total for Question 7 is 9 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

8. (a) Use the standard results for $\sum_{r=1}^{n} r^2$ and $\sum_{r=1}^{n} r$ to show that for all positive integers n

$$\sum_{r=0}^{n} (r+1)(r+2) = \frac{1}{3}(n+1)(n+2)(n+3)$$

(5)

(b) Hence determine the value of

$$10 \times 11 + 11 \times 12 + 12 \times 13 + \ldots + 100 \times 101$$

(3)

(a) $\sum_{r=0}^{n} (r+1)(r+2) = \sum_{r=0}^{n} (r^2 + 3r + 2) = 2 + \sum_{r=1}^{n} r^2 + 3\sum_{r=1}^{n} r + \sum_{r=1}^{n} 2$

$= 2 + \frac{1}{6} n(n+1)(2n+1) + \frac{3}{2} n(n+1) + 2n$

$= \frac{1}{6} n(n+1)(2n+1) + \frac{3}{2} n(n+1) + 2(n+1)$

$= \frac{1}{6} (n+1)[n(2n+1) + 9n + 12]$

$= \frac{1}{6} (n+1)(2n^2 + 10n + 12)$

$= \frac{1}{3} (n+1)(n^2 + 5n + 6)$

$= \frac{1}{3} (n+1)(n+2)(n+3)$ as required

(b) 10×11 is the term with $r = 9$, 100×101 is the term with $r = 99$

Sum $= \sum_{r=0}^{99} (r+1)(r+2) - \sum_{r=0}^{8} (r+1)(r+2)$

$= \frac{1}{3} (100)(101)(102) - \frac{1}{3} (9)(10)(11)$

$= 343400 - 330$

$= 343070$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 8 continued

Question 8 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 8 continued

(Total for Question 8 is 8 marks)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

9. (i) A sequence of numbers is defined by

$$u_1 = 3$$

$$u_{n+1} = 2u_n - 2^{n+1} \qquad n \geqslant 1$$

Prove by induction that, for $n \in \mathbb{N}$

$$u_n = 5 \times 2^{n-1} - n \times 2^n$$

(5)

(ii) Prove by induction that, for $n \in \mathbb{N}$

$$f(n) = 5^{n+2} - 4n - 9$$

is divisible by 16

(5)

(i) $n = 1$: $u_1 = 5 \times 2^0 - 1 \times 2^1 = 5 - 2 = 3$, so true for $n = 1$

Assume true for $n = k$: $u_k = 5 \times 2^{k-1} - k \times 2^k$

$u_{k+1} = 2u_k - 2^{k+1} = 2(5 \times 2^{k-1} - k \times 2^k) - 2^{k+1}$

$= 5 \times 2^k - k \times 2^{k+1} - 2^{k+1}$

$= 5 \times 2^{(k+1)-1} - (k + 1) \times 2^{k+1}$

which is the result with $n = k + 1$.

If the result is true for $n = k$ then it is true for $n = k + 1$. As the result has been shown to be true for $n = 1$, then the result is true for all $n \in \mathbb{N}$ by mathematical induction.

(ii) $n = 1$: $f(1) = 5^3 - 4 - 9 = 112 = 16 \times 7$, so divisible by 16

Assume true for $n = k$: $f(k) = 5^{k+2} - 4k - 9$ is divisible by 16

$f(k + 1) = 5^{k+3} - 4(k + 1) - 9 = 5 \times 5^{k+2} - 4k - 13$

$= 5(5^{k+2} - 4k - 9) + 20k + 45 - 4k - 13$

$= 5(5^{k+2} - 4k - 9) + 16k + 32$

$= 5f(k) + 16(k + 2)$

$5f(k)$ and $16(k + 2)$ are both divisible by 16, so $f(k + 1)$ is divisible by 16.

If the result is true for $n = k$ then it is true for $n = k + 1$. As the result has been shown to be true for $n = 1$, then the result is true for all $n \in \mathbb{N}$ by mathematical induction.

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 9 continued

Question 9 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 9 continued

Question 9 continued

(Total for Question 9 is 10 marks)

TOTAL FOR PAPER: 75 MARKS

END

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA