Please check the examination details below before entering your candidate information

Candidate surname StemUp Model Solution | Other names

Pearson Edexcel
International
Advanced Level

Centre Number | Candidate Number

Tuesday 8 January 2019

Morning (Time: 1 hour 30 minutes) | Paper Reference **WMA11/01**

Mathematics

Advanced Subsidiary
Pure Mathematics P1

You must have:
Mathematical Formulae and Statistical Tables (Lilac), calculator

Total Marks

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
  – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 12 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
  – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶

P60791A

©2019 Pearson Education Ltd.

1/1/1/1/1/C2/

P 6 0 7 9 1 A 0 1 2 8

Pearson

Leave blank

Answer ALL questions. Write your answers in the spaces provided.

1. Find

$$\int\left(\frac{2}{3}x^3 - \frac{1}{2x^3} + 5\right)\mathrm{d}x$$

simplifying your answer.

(4)

$\int\left(\frac{2}{3}x^3 - \frac{1}{2}x^{-3} + 5\right)dx$

$= \frac{2}{3} \times \frac{x^4}{4} - \frac{1}{2} \times \frac{x^{-2}}{-2} + 5x + c$

$= \frac{1}{6}x^4 + \frac{1}{4}x^{-2} + 5x + c$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 1 continued

Q1

(Total 4 marks)

Leave blank

2. Given

$$\frac{3^x}{3^{4y}} = 27\sqrt{3}$$

find y as a simplified function of x.

(3)

$27\sqrt{3} = 3^3 \times 3^{1/2} = 3^{7/2}$

$\frac{3^x}{3^{4y}} = 3^{x-4y} \Rightarrow 3^{x-4y} = 3^{7/2}$

$x - 4y = \frac{7}{2} \Rightarrow 4y = x - \frac{7}{2}$

$y = \frac{1}{4}x - \frac{7}{8}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 2 continued

Q2

(Total 3 marks)

Leave blank

3. The line l_1 has equation $3x + 5y - 7 = 0$

(a) Find the gradient of l_1

(2)

The line l_2 is perpendicular to l_1 and passes through the point $(6, -2)$.

(b) Find the equation of l_2 in the form $y = mx + c$, where m and c are constants.

(3)

(a) $5y = -3x + 7 \Rightarrow y = -\frac{3}{5}x + \frac{7}{5}$

Gradient of $l_1 = -\frac{3}{5}$

(b) Gradient of $l_2 = -1 \div \left(-\frac{3}{5}\right) = \frac{5}{3}$

$y - (-2) = \frac{5}{3}(x - 6)$

$y + 2 = \frac{5}{3}x - 10$

$y = \frac{5}{3}x - 12$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 3 continued

Q3

(Total 5 marks)

Leave blank

4.




Figure 1

Figure 1 shows a line l_1 with equation $2y = x$ and a curve C with equation $y = 2x - \frac{1}{8}x^2$

The region R, shown unshaded in Figure 1, is bounded by the line l_1, the curve C and a line l_2

Given that l_2 is parallel to the y-axis and passes through the intercept of C with the positive x-axis, identify the inequalities that define R.

(3)

C meets x-axis: $2x - \frac{1}{8}x^2 = 0 \Rightarrow x(2 - \frac{1}{8}x) = 0$

$\Rightarrow$ $x = 0$ or $x = 16$, so l_2 is $x = 16$

l_2 is dashed (strict), l_1 and C are solid:

$x < 16$, $2y \leq x$, $y \geq 2x - \frac{1}{8}x^2$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 4 continued

Q4

(Total 3 marks)

Leave blank

5.




Figure 2

Figure 2 shows a plot of part of the curve with equation $y = \cos 2x$ with x being measured in radians.

The point P, shown on Figure 2, is a minimum point on the curve.

(a) State the coordinates of P.

(2)

A copy of Figure 2, called Diagram 1, is shown at the top of the next page.

(b) Sketch, on Diagram 1, the curve with equation $y = \sin x$

(2)

(c) Hence, or otherwise, deduce the number of solutions of the equation

(i) $\cos 2x = \sin x$ that lie in the region $0 \leqslant x \leqslant 20\pi$

(ii) $\cos 2x = \sin x$ that lie in the region $0 \leqslant x \leqslant 21\pi$

(2)

(a) $y = \cos 2x$ has its first minimum when $2x = \pi$

P $\left(\frac{\pi}{2}, -1\right)$

(c) In $0 \leq x \leq 2\pi$ the curves meet 3 times

(i) $20\pi = 10 \times 2\pi \Rightarrow 10 \times 3 = 30$ solutions

(ii) in 20π to 21π (like 0 to π) 2 more $\Rightarrow$ 32 solutions

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 5 continued




Diagram 1

(b) $y = \sin x$ drawn on Diagram 1 (period 2π, max 1, min -1)

(c) check: $\cos 2x = \sin x \Rightarrow 1 - 2\sin^2 x = \sin x$

$2\sin^2 x + \sin x - 1 = 0 \Rightarrow (2\sin x - 1)(\sin x + 1) = 0$

$\sin x = \frac{1}{2}$: $x = \frac{\pi}{6}, \frac{5\pi}{6}$; $\sin x = -1$: $x = \frac{3\pi}{2}$

Q5

(Total 6 marks)

Leave blank

6. (*Solutions based entirely on graphical or numerical methods are not acceptable.*)

Given

$$f(x) = 2x^{\frac{5}{2}} - 40x + 8 \qquad x > 0$$

(a) solve the equation $f'(x) = 0$

(4)

(b) solve the equation $f''(x) = 5$

(3)

(a) $f'(x) = 2 \times \frac{5}{2} x^{3/2} - 40 = 5x^{3/2} - 40$

$5x^{3/2} - 40 = 0 \Rightarrow x^{3/2} = 8$

$x = 8^{2/3} = 4$

(b) $f''(x) = \frac{3}{2} \times 5x^{1/2} = \frac{15}{2} x^{1/2}$

$\frac{15}{2} x^{1/2} = 5 \Rightarrow x^{1/2} = \frac{10}{15} = \frac{2}{3}$

$x = (\frac{2}{3})^2 = \frac{4}{9}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank


Question 6 continued

Q6

(Total 7 marks)

Leave blank

7.



Not to scale

Figure 3

Figure 3 shows the design for a structure used to support a roof.

The structure consists of four wooden beams, AB, BD, BC and AD.

Given $AB = 6.5\,\text{m}$, $BC = BD = 4.7\,\text{m}$ and angle $BAC = 35°$

(a) find, to one decimal place, the size of angle ACB,

(3)

(b) find, to the nearest metre, the total length of wood required to make this structure.

(3)

(a) $\frac{\sin \angle ACB}{6.5} = \frac{\sin 35°}{4.7}$

$\sin \angle ACB = \frac{6.5 \sin 35°}{4.7} = 0.79324...$

$\angle ACB = 52.4897...°$ or $180° - 52.4897...° = 127.5102...°$

From the figure $\angle ACB$ is obtuse: $\angle ACB = 127.5°$

(b) $\angle ABC = 180° - 35° - 127.51° = 17.49°$

$AC = \frac{6.5 \sin 17.49°}{\sin 127.51°} = 2.4626...\ m$

Triangle BCD is isosceles: $\angle BCD = \angle BDC = 52.49°$, $\angle CBD = 75.02°$

$CD = \frac{4.7 \sin 75.02°}{\sin 52.49°} = 5.7236...\ m$

$AD = AC + CD = 2.4626... + 5.7236... = 8.1863...\ m$

$Total = AB + BC + BD + AD = 6.5 + 4.7 + 4.7 + 8.1863...$

$= 24.086...\ m \approx 24\ m$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 7 continued

Q7

(Total 6 marks)

Leave blank

8.




Figure 4

The curve C with equation $y = \mathrm{f}(x)$ is shown in Figure 4.

The curve C

- has a single turning point, a maximum at (4, 9)
- crosses the coordinate axes at only two places, (−3, 0) and (0, 6)
- has a single asymptote with equation $y = 4$

as shown in Figure 4.

(a) State the equation of the asymptote to the curve with equation $y = \mathrm{f}(-x)$.

(1)

(b) State the coordinates of the turning point on the curve with equation $y = \mathrm{f}\left(\frac{1}{4}x\right)$.

(1)

Given that the line with equation $y = k$, where k is a constant, intersects C at exactly one point,

(c) state the possible values for k.

(2)

The curve C is transformed to a new curve that passes through the origin.

(d) (i) Given that the new curve has equation $y = \mathrm{f}(x) - a$, state the value of the constant a.

(ii) Write down an equation for another single transformation of C that also passes through the origin.

(2)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 8 continued

(a) $y = 4$

(b) $(16, 9)$

(c) $k \leqslant 4, \quad k = 9$

(d) (i) $f(0) = 6$, so $y = f(x) - 6$ passes through 0: $a = 6$

(ii) $y = f(x - 3)$ (moves $(-3, 0)$ to the origin)

Q8

(Total 6 marks)

Leave blank

9. The equation

$$\frac{3}{x} + 5 = -2x + c$$

where c is a constant, has no real roots.

Find the range of possible values of c.

(7)

$\frac{3}{x} + 5 = -2x + c \quad (\times x)$

$3 + 5x = -2x^2 + cx$

$2x^2 + (5 - c)x + 3 = 0$

No real roots: $b^2 - 4ac < 0$

$(5 - c)^2 - 4 \times 2 \times 3 < 0 \Rightarrow (5 - c)^2 - 24 < 0$

Critical values: $(5 - c)^2 = 24 \Rightarrow 5 - c = \pm\sqrt{24}$

$c = 5 \pm \sqrt{24} = 5 \pm 2\sqrt{6}$

Quadratic in c is negative between its roots:

$5 - 2\sqrt{6} < c < 5 + 2\sqrt{6}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 9 continued

Q9

(Total 7 marks)

Leave blank

10. A sector AOB, of a circle centre O, has radius r cm and angle θ radians.

Given that the area of the sector is 6 cm^2 and that the perimeter of the sector is 10 cm,

(a) show that

$$3\theta^2 - 13\theta + 12 = 0$$

(4)

(b) Hence find possible values of r and θ.

(3)

(a) Area: $\frac{1}{2} r^2\theta = 6$ Perimeter: $2r + r\theta = 10$

$r(2 + \theta) = 10 \Rightarrow r = \frac{10}{2 + \theta}$

$\frac{1}{2}\left(\frac{10}{2 + \theta}\right)^2 \theta = 6 \Rightarrow \frac{50\theta}{(2 + \theta)^2} = 6$

$50\theta = 6(4 + 4\theta + \theta^2) = 24 + 24\theta + 6\theta^2$

$6\theta^2 - 26\theta + 24 = 0$

$3\theta^2 - 13\theta + 12 = 0$ as required

(b) $(3\theta - 4)(\theta - 3) = 0$

$\theta = \frac{4}{3}$ or $\theta = 3$

$\theta = \frac{4}{3}$: $r = \frac{10}{2 + 4/3} = \frac{10}{10/3} = 3$

$\theta = 3$: $r = \frac{10}{2 + 3} = 2$

$r = 3, \theta = \frac{4}{3}$ or $r = 2, \theta = 3$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 10 continued

Q10

(Total 7 marks)

Leave blank

11. (a) On Diagram 1 sketch the graphs of

(i) $y = x(3 - x)$

(ii) $y = x(x - 2)(5 - x)$

showing clearly the coordinates of the points where the curves cross the coordinate axes.

(4)

(b) Show that the x coordinates of the points of intersection of

$$y = x(3 - x) \text{ and } y = x(x - 2)(5 - x)$$

are given by the solutions to the equation $x(x^2 - 8x + 13) = 0$

(3)

The point P lies on both curves. Given that P lies in the first quadrant,

(c) find, using algebra and showing your working, the exact coordinates of P.

(5)

(b) $x(3 - x) = x(x - 2)(5 - x)$

$3x - x^2 = x(-x^2 + 7x - 10) = -x^3 + 7x^2 - 10x$

$x^3 - 8x^2 + 13x = 0$

$x(x^2 - 8x + 13) = 0$ as required

(c) $x^2 - 8x + 13 = 0 \Rightarrow x = \frac{8 \pm \sqrt{(64 - 52)}}{2} = \frac{8 \pm \sqrt{12}}{2}$

$x = 4 \pm \sqrt{3}$

First quadrant needs $y = x(3 - x) > 0$, so $0 < x < 3$:

$4 + \sqrt{3} \approx 5.73$ rejected, $x = 4 - \sqrt{3} \approx 2.27$

$y = (4 - \sqrt{3})(3 - (4 - \sqrt{3})) = (4 - \sqrt{3})(-1 + \sqrt{3})$

$y = -4 + 4\sqrt{3} + \sqrt{3} - 3 = 5\sqrt{3} - 7$

$P\ (4 - \sqrt{3},\ 5\sqrt{3} - 7)$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 11 continued




Diagram 1

(a) (i) $y = x(3 - x)$: ∩ shape, crosses x-axis at (0, 0) and (3, 0)

(ii) $y = x(x - 2)(5 - x)$: negative cubic, crosses at (0, 0), (2, 0), (5, 0)

Leave blank

Question 11 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 11 continued

Q11

(Total 12 marks)

Leave blank

12. The curve with equation $y = f(x), x > 0,$ passes through the point $P(4, -2)$.

Given that

$$\frac{dy}{dx} = 3x\sqrt{x} - 10x^{-\frac{1}{2}}$$

(a) find the equation of the tangent to the curve at P, writing your answer in the form $y = mx + c$, where m and c are integers to be found.

(4)

(b) Find $f(x)$.

(5)

(a) $x = 4$: $\frac{dy}{dx} = 3 \times 4 \times \sqrt{4} - 10 \times 4^{-1/2} = 24 - \frac{10}{2} = 19$

Tangent: $y - (-2) = 19(x - 4)$

$y + 2 = 19x - 76$

$y = 19x - 78$

(b) $\frac{dy}{dx} = 3x^{3/2} - 10x^{-1/2}$

$f(x) = \frac{3x^{5/2}}{5/2} - \frac{10x^{1/2}}{1/2} + c = \frac{6}{5}x^{5/2} - 20x^{1/2} + c$

$P(4, -2)$: $-2 = \frac{6}{5} \times 32 - 20 \times 2 + c$

$-2 = 38.4 - 40 + c \Rightarrow c = -0.4$

$f(x) = \frac{6}{5}x^{5/2} - 20x^{1/2} - \frac{2}{5}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 12 continued

Leave blank

Question 12 continued

Q12

(Total 9 marks)

TOTAL FOR PAPER IS 75 MARKS

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA