Please check the examination details below before entering your candidate information

Candidate surname	Other names
StemUp Model Solution	

Centre Number | Candidate Number

Pearson Edexcel International Advanced Level

Time 1 hour 30 minutes

Paper reference **WMA12/01**

Mathematics

International Advanced Subsidiary/Advanced Level
Pure Mathematics P2

You must have:
Mathematical Formulae and Statistical Tables (Yellow), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
  – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 10 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
  – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶

P71377A

©2022 Pearson Education Ltd.

Q:1/1/1/1/

P 7 1 3 7 7 A 0 1 3 2

Pearson

Leave blank

1. Find the first four terms, in ascending powers of x, of the binomial expansion of

$$\left(2+\frac{3}{8}x\right)^{10}$$

Give each coefficient as an integer.

(4)

$(2 + (3/8)x)^{10}$

$= 2^{10} + {}^{10}C_1 \times 2^9 \times ((3/8)x) + {}^{10}C_2 \times 2^8 \times ((3/8)x)^2$

$\quad + {}^{10}C_3 \times 2^7 \times ((3/8)x)^3 + \ldots$

$= 1024 + 10 \times 512 \times (3/8)x + 45 \times 256 \times (9/64)x^2$

$\quad + 120 \times 128 \times (27/512)x^3 + \ldots$

$= 1024 + 1920x + 1620x^2 + 810x^3 + \ldots$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 1 continued

Q1

(Total 4 marks)

Leave blank

2.




Figure 1

Figure 1 shows the graph of

$$y = 1 - \log_{10}(\sin x) \qquad 0 < x < \pi$$

where x is in radians.

The table below shows some values of x and y for this graph, with values of y given to 3 decimal places.

x	0.5	1	1.5	2	2.5	3
y	1.319	1.075	1.001	1.041	1.223	1.850

(a) Complete the table above, giving values of y to 3 decimal places.

(2)

(b) Use the trapezium rule with all the y values in the completed table to find, to 2 decimal places, an estimate for

$$\int_{0.5}^{3} \left(1 - \log_{10}(\sin x)\right) \mathrm{d}x$$

(3)

(c) Use your answer to part (b) to find an estimate for

$$\int_{0.5}^{3} \left(3 + \log_{10}(\sin x)\right) \mathrm{d}x$$

(3)

(a) $x = 1$: $y = 1 - \log_{10}(\sin 1) = 1.07496... = 1.075$

$x = 2$: $y = 1 - \log_{10}(\sin 2) = 1.04129... = 1.041$

(b) $h = 0.5$

$\int \approx (0.5/2) \times [1.319 + 1.850 + 2(1.075 + 1.001 + 1.041 + 1.223)]$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 2 continued

$= 0.25 \times [3.169 + 8.680] = 0.25 \times 11.849$

$= 2.96225...$

$= 2.96$ (2 d.p.)

(c) $3 + \log_{10}(\sin x) = 4 - (1 - \log_{10}(\sin x))$

$\int_{0.5}^{3} (3 + \log_{10}(\sin x))\, dx = \int_{0.5}^{3} 4\, dx - \int_{0.5}^{3} (1 - \log_{10}(\sin x))\, dx$

$= [4x]_{0.5}^{3} - 2.96$

$= 12 - 2 - 2.96$

$= 7.04$

Leave blank

Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 2 continued

Q2

(Total 8 marks)

Leave blank

3. (i) Show that the following statement is **false**:

"$(n + 1)^3 - n^3$ is prime for all $n \in \mathbb{N}$"

(2)

(ii) Given that the points $A(1, 0)$, $B(3, -10)$ and $C(7, -6)$ lie on a circle, prove that AB is a diameter of this circle.

(5)

(i) $n = 1$: $2^3 - 1^3 = 7$ (prime), $n = 2$: $3^3 - 2^3 = 19$ (prime), ...

$n = 5$: $6^3 - 5^3 = 216 - 125 = 91 = 7 \times 13$

91 is not prime, so the statement is false (counterexample $n = 5$).

(ii) Gradient of AC: $m_{AC} = (-6 - 0)/(7 - 1) = -6/6 = -1$

Gradient of BC: $m_{BC} = (-6 - (-10))/(7 - 3) = 4/4 = 1$

$m_{AC} \times m_{BC} = -1 \times 1 = -1$

$\therefore$ AC is perpendicular to BC, so angle ACB = 90°

The angle in a semicircle is a right angle. Since A, B and C lie on the circle and angle ACB = 90°, AB must be a diameter of the circle.

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 3 continued

Leave blank

Leave blank

Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 3 continued

Q3

(Total 7 marks)

Leave blank

4. **In this question you must show all stages of your working.**

Give your answers in fully simplified surd form.

Given that a and b are positive constants, solve the simultaneous equations

$$a - b = 8$$
$$\log_4 a + \log_4 b = 3$$

(6)

$\log_4 a + \log_4 b = 3 \Rightarrow \log_4(ab) = 3$

$\Rightarrow ab = 4^3 = 64$

$a - b = 8 \Rightarrow a = b + 8$

$\Rightarrow (b + 8)b = 64$

$\Rightarrow b^2 + 8b - 64 = 0$

$b = (-8 \pm \sqrt{(8^2 - 4 \times 1 \times (-64))})/(2 \times 1) = (-8 \pm \sqrt{320})/2$

$= (-8 \pm 8\sqrt{5})/2 = -4 \pm 4\sqrt{5}$

$b > 0 \Rightarrow b = -4 + 4\sqrt{5}$

$a = b + 8 = 4 + 4\sqrt{5}$

$a = 4 + 4\sqrt{5}, \quad b = -4 + 4\sqrt{5}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 4 continued

Q4

(Total 6 marks)

Leave blank

5. **In this question you must show all stages of your working.**

Solutions relying entirely on calculator technology are not acceptable.

Solve, for $-180° < \theta \leqslant 180°$, the equation

$$3\tan(\theta + 43°) = 2\cos(\theta + 43°)$$

(6)

$3\tan(\theta + 43°) = 2\cos(\theta + 43°)$

$\Rightarrow 3\sin(\theta + 43°)/\cos(\theta + 43°) = 2\cos(\theta + 43°)$

$\Rightarrow 3\sin(\theta + 43°) = 2\cos^2(\theta + 43°)$

$\Rightarrow 3\sin(\theta + 43°) = 2(1 - \sin^2(\theta + 43°))$

$\Rightarrow 2\sin^2(\theta + 43°) + 3\sin(\theta + 43°) - 2 = 0$

$\Rightarrow (2\sin(\theta + 43°) - 1)(\sin(\theta + 43°) + 2) = 0$

$\Rightarrow \sin(\theta + 43°) = 1/2$ $(\sin(\theta + 43°) = -2$ has no solutions$)$

$-137° < \theta + 43° \leq 223°$

$\theta + 43° = 30°$ or $\theta + 43° = 180° - 30° = 150°$

$\theta = 30° - 43° = -13°$, $\theta = 150° - 43° = 107°$

$\theta = -13°, 107°$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 5 continued

Q5

(Total 6 marks)

Leave blank

6. In a geometric sequence $u_1, u_2, u_3, \ldots$

- the common ratio is r
- $u_2 + u_3 = 6$
- $u_4 = 8$

(a) Show that r satisfies

$$3r^2 - 4r - 4 = 0$$

(3)

Given that the geometric sequence has a sum to infinity,

(b) find u_1

(3)

(c) find S_∞

(2)

(a) $u_n = ar^{n-1}$

$u_2 + u_3 = 6 \Rightarrow ar + ar^2 = 6$

$u_4 = 8 \Rightarrow ar^3 = 8$

Dividing: $(ar + ar^2)/(ar^3) = 6/8$

$\Rightarrow (1 + r)/r^2 = 3/4$

$\Rightarrow 4 + 4r = 3r^2$

$\Rightarrow 3r^2 - 4r - 4 = 0$ as required

(b) $(3r + 2)(r - 2) = 0 \Rightarrow r = -2/3$ or $r = 2$

Sum to infinity exists so $|r| < 1 \Rightarrow r = -2/3$

$ar^3 = 8 \Rightarrow a = 8 \div (-2/3)^3 = 8 \div (-8/27) = -27$

$u_1 = -27$

(c) $S_\infty = a/(1 - r) = -27/(1 - (-2/3)) = -27 \div (5/3)$

$S_\infty = -81/5 \ (= -16.2)$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 6 continued

Leave blank

Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 6 continued

Q6

(Total 8 marks)

Leave blank

7. $f(x) = Ax^3 + 6x^2 - 4x + B$

where A and B are constants.

Given that

- $(x + 2)$ is a factor of $f(x)$
- $\int_3^5 f(x)\,dx = 176$

find the value of A and the value of B.

(7)

$(x + 2)$ is a factor $\Rightarrow f(-2) = 0$

$A(-2)^3 + 6(-2)^2 - 4(-2) + B = 0$

$\Rightarrow -8A + 24 + 8 + B = 0 \Rightarrow 8A - B = 32 \quad (1)$

$\int f(x)\,dx = (A/4)x^4 + 2x^3 - 2x^2 + Bx$

$\int_3^5 f(x)\,dx = [(A/4)x^4 + 2x^3 - 2x^2 + Bx]_3^5 = 176$

$\Rightarrow (625A/4 + 250 - 50 + 5B) - (81A/4 + 54 - 18 + 3B) = 176$

$\Rightarrow 136A + 164 + 2B = 176$

$\Rightarrow 136A + 2B = 12 \quad (2)$

$(1) \times 2$: $16A - 2B = 64$

Adding: $152A = 76 \Rightarrow A = 1/2$

$B = 8A - 32 = 4 - 32 = -28$

$A = 1/2, \quad B = -28$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 7 continued

Q7

(Total 7 marks)

Leave blank

8. **In this question you must show all stages of your working.**

Solutions relying entirely on calculator technology are not acceptable.

A curve has equation

$$y = 256x^4 - 304x - 35 + \frac{27}{x^2} \qquad x \neq 0$$

(a) Find $\frac{dy}{dx}$

(3)

(b) Hence find the coordinates of the stationary points of the curve.

(5)

$y = 256x^4 - 304x - 35 + 27x^{-2}$

(a) $dy/dx = 4 \times 256x^3 - 304 - 2 \times 27x^{-3}$

$dy/dx = 1024x^3 - 304 - 54/x^3$

(b) At stationary points $dy/dx = 0$:

$1024x^3 - 304 - 54/x^3 = 0 \quad (\times x^3)$

$\Rightarrow 1024x^6 - 304x^3 - 54 = 0$

$\Rightarrow 512(x^3)^2 - 152x^3 - 27 = 0$

$\Rightarrow (8x^3 + 1)(64x^3 - 27) = 0$

$\Rightarrow x^3 = -1/8$ or $x^3 = 27/64$

$\Rightarrow x = -1/2$ or $x = 3/4$

$x = -1/2$: $y = 256(1/16) - 304(-1/2) - 35 + 27 \div (1/4)$

$= 16 + 152 - 35 + 108 = 241$

$x = 3/4$: $y = 256(81/256) - 304(3/4) - 35 + 27 \div (9/16)$

$= 81 - 228 - 35 + 48 = -134$

Stationary points: $(-1/2, 241)$ and $(3/4, -134)$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 8 continued

Q8

(Total 8 marks)

Leave blank

9. A scientist is using carbon-14 dating to determine the age of some wooden items.

The equation for carbon-14 dating an item is given by

$$N = k\lambda^t$$

where

- N grams is the amount of carbon-14 **currently** present in the item
- k grams was the **initial** amount of carbon-14 present in the item
- t is the number of years since the item was made
- λ is a constant, with $0 < \lambda < 1$

(a) Sketch the graph of N against t for $k = 1$

(2)

Given that it takes 5700 years for the amount of carbon-14 to reduce to half its initial value,

(b) show that the value of the constant λ is 0.999878 to 6 decimal places.

(2)

Given that Item A

- is known to have had 15 grams of carbon-14 present initially
- is thought to be 3250 years old

(c) calculate, to 3 significant figures, how much carbon-14 the equation predicts is currently in Item A.


(2)

Item B is known to have initially had 25 grams of carbon-14 present, but only 18 grams now remain.

(d) Use algebra to calculate the age of Item B to the nearest 100 years.

(3)

(a)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 9 continued

(b) After 5700 years, $N = k/2$:

$(1/2)k = k\lambda^{5700} \Rightarrow \lambda^{5700} = 1/2$

$\Rightarrow \lambda = (1/2)^{1/5700}$

$= 0.99987840... = 0.999878$ (6 d.p.) as required

(c) $k = 15$, $t = 3250$:

$N = 15 \times 0.999878^{3250}$

$= 10.0898...$

$N = 10.1$ grams (3 s.f.)

(d) $18 = 25 \times 0.999878^t$

$\Rightarrow 0.999878^t = 18/25 = 0.72$

$\Rightarrow t \log 0.999878 = \log 0.72$

$\Rightarrow t = \log 0.72 \div \log 0.999878$

$= 2692.49...$

Age of Item B = 2700 years (to the nearest 100 years)

Leave blank

Question 9 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 9 continued

Q9

(Total 9 marks)

Leave blank

10. The circle C has centre $X(3, 5)$ and radius r

The line l has equation $y = 2x + k$, where k is a constant.

(a) Show that l and C intersect when

$$5x^2 + (4k - 26)x + k^2 - 10k + 34 - r^2 = 0$$

(3)

Given that l is a tangent to C,

(b) show that $5r^2 = (k + p)^2$, where p is a constant to be found.

(3)




Figure 2

The line l

- cuts the y-axis at the point A
- touches the circle C at the point B

as shown in Figure 2.

Given that $AB = 2r$

(c) find the value of k

(6)

(a) $C: (x - 3)^2 + (y - 5)^2 = r^2$, $l: y = 2x + k$

$(x - 3)^2 + (2x + k - 5)^2 = r^2$

$x^2 - 6x + 9 + 4x^2 + 4(k - 5)x + (k - 5)^2 = r^2$

$5x^2 + (-6 + 4k - 20)x + 9 + k^2 - 10k + 25 - r^2 = 0$

$\Rightarrow 5x^2 + (4k - 26)x + k^2 - 10k + 34 - r^2 = 0$ as required

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 10 continued

(b) l is a tangent $\Rightarrow$ repeated root $\Rightarrow b^2 - 4ac = 0$

$(4k - 26)^2 - 4 \times 5 \times (k^2 - 10k + 34 - r^2) = 0$

$16k^2 - 208k + 676 - 20k^2 + 200k - 680 + 20r^2 = 0$

$\Rightarrow 20r^2 = 4k^2 + 8k + 4$

$\Rightarrow 5r^2 = k^2 + 2k + 1$

$\Rightarrow 5r^2 = (k + 1)^2$, so $p = 1$

(c) $A = (0, k)$. XB is perpendicular to l (radius $\perp$ tangent),

so triangle AXB is right-angled at B: $AB^2 + XB^2 = XA^2$

$XA^2 = (3 - 0)^2 + (5 - k)^2 = 9 + (5 - k)^2$

$AB = 2r \Rightarrow AB^2 + r^2 = 4r^2 + r^2 = 5r^2$

$\Rightarrow 5r^2 = 9 + (5 - k)^2$

Using (b): $(k + 1)^2 = 9 + (5 - k)^2$

$k^2 + 2k + 1 = 9 + 25 - 10k + k^2$

$\Rightarrow 12k = 33$

$k = 11/4$

Leave blank

Question 10 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 10 continued

Leave blank

Question 10 continued

Q10

(Total 12 marks)

TOTAL FOR PAPER IS 75 MARKS

END

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA