Surname	Centre Number	Candidate Number
First name(s)		2

GCE A LEVEL

A400U10-1

S25-A400U10-1

THURSDAY, 5 JUNE 2025 – AFTERNOON

BIOLOGY – A level component 1
Energy for Life

2 hours

For Examiner's use only		
Question	**Maximum Mark**	**Mark Awarded**
1.	**12**	
2.	**14**	
3.	**17**	
4.	**21**	
5.	**19**	
6.	**8**	
7.	**9**	
Total	**100**	

A400U101
01

ADDITIONAL MATERIALS

A calculator and a ruler.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen. Do not use gel pen or correction fluid.

You may use a pencil for graphs and diagrams only.

Write your name, centre number and candidate number in the spaces at the top of this page.

Answer **all** questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the additional page(s) at the back of the booklet, taking care to number the question(s) correctly.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

The assessment of the quality of extended response (QER) will take place in question **7**.

The quality of written communication will affect the awarding of marks.

JUN25A400U10101

© WJEC CBAC Ltd.

BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE

© WJEC CBAC Ltd.

Examiner only

Answer **all** questions.

1. The oxidation of respiratory substrates is catalysed by enzymes. In a series of reactions hydrogen ions are produced which are transferred to the hydrogen carriers involved in respiration. These carriers then deliver hydrogen to the site of ATP synthesis.

(a) (i) Name **two** hydrogen carriers that are involved in aerobic respiration in mitochondria. [1]

....

(ii) Describe the role of hydrogen ions in ATP synthesis. [3]

.....

.....

.....

.....

.....

.....

.....

(iii) Explain why the two hydrogen carriers involved in aerobic respiration result in the synthesis of different numbers of ATP molecules. [2]

.....

.....

.....

.....

.....

.....


A400U101
03

© WJEC CBAC Ltd.

Examiner only

(b) Resazurin dye changes colour when it is reduced and will change colour in the presence of respiring cells. If resazurin is present it will be reduced by hydrogen and changes colour from dark blue to pink to pale pink as shown in **Image 1.1**.

Image 1.1



Students used resazurin to investigate the action of dehydrogenase in yeast. They set up a thermostatically controlled water bath at 35 °C. Two test tubes were prepared as follows and left in the water bath for five minutes:

Test tube **A**: 3 cm^3 of boiled 2.5% yeast suspension + 3 cm^3 of 5% glucose solution
Test tube **B**: 3 cm^3 of 2.5% yeast suspension + 3 cm^3 of 5% glucose solution

After five minutes, 3 cm^3 of 0.01% resazurin was added to test tubes **A** and **B**. The resazurin solution was at the same temperature as the yeast and glucose mixture.

(i) Explain why the resazurin was brought to the same temperature as the yeast and glucose mixtures before mixing. [1]

.....

.....

.....

(ii) Explain the purpose of test tube **A**. [2]

.....


.....

.....

.....

The colour of the contents of each test tube was recorded every three minutes for a period of 30 minutes. The colour of each tube was matched to the colour chart shown in **Image 1.2**. The numbers indicate the extent of reduction of resazurin, with 10 indicating no reduction.

Image 1.2



© WJEC CBAC Ltd.

Examiner only

Table 1.3 shows one student's results.

Table 1.3

Time / minutes	Extent of reduction of resazurin / au	
	test tube **A**	test tube **B**
0	10	8
3	10	8
6	10	7
9	10	6
12	10	5
15	10	4
18	10	3
21	10	2
24	10	1
27	10	1
30	10	1

The results of other students varied considerably from those shown in the table.

A400U101 05

(iii) I. Use all the information given to identify and explain how **one** limitation in this investigation could account for the variation in the results of other students.

[2]

.....

.....

.....

.....

II. Suggest and explain **one** improvement that would provide results that are more accurate. [1]

.....

.....

.....

.....

12

© WJEC CBAC Ltd.

Examiner only

2. (a) *Streptococcus mutans* is one of at least 25 species of streptococcus bacteria found in the human mouth. A chain of *S. mutans* cells following Gram staining is shown in **Image 2.1**.

Image 2.1



(i) Classification of the different streptococcus bacteria is tentative as all these species appear similar after Gram staining.
Explain the meaning of the term **tentative** when referring to classification. [1]

.....

.....

(ii) Explain how **one** biochemical method of analysis could be used to distinguish between the different species of *Streptococcus*. [2]

.....

.....

.....

.....


© WJEC CBAC Ltd.

Examiner only

(b) A microscope was used to measure the diameter of one of the bacteria shown in **Image 2.1**. The microscope was first calibrated using an eyepiece graticule and a stage micrometer. Divisions marked on the stage micrometer have a constant value of 0.01 mm but the value of one eyepiece unit (epu) varies depending on the magnification of the objective lens.

Image 2.2 shows how a stage micrometer and eyepiece graticule would appear when viewed down a microscope at high magnification.

Image 2.2



(i) Name scale **X** in **Image 2.2**. [1]

.....

(ii) At this magnification it was calculated that 1 epu = 1.05 μm.
The diameter of one of the bacteria in **Image 2.1** was measured at 2 epu.
Calculate the actual diameter of this bacterium. [1]

Diameter of bacterium = ... μm

A400U101
07

© WJEC CBAC Ltd.


Examiner only

(c) In humans, *S. mutans* is most prevalent on the surface of teeth and is one of the main organisms responsible for tooth decay.
S. mutans can convert sucrose into two main products:

- lactic acid
- dextran – a sticky polysaccharide

S. mutans can also metabolise other sugars, including glucose, fructose and lactose to produce lactic acid.
Image 2.3 shows a cavity in a human premolar caused by tooth decay.

Image 2.3



(i) With reference to the habitat of this species of bacteria and its metabolism of sugars, conclude whether *S. mutans* is an aerobe, an obligate anaerobe or a facultative anaerobe. Explain your answer. [3]

.....

.....

.....

.....

.....

.....

.....

© WJEC CBAC Ltd.

Examiner only

(ii) Use the information provided to explain why **not** brushing your teeth can increase the risk of cavity formation. [2]

.....

.....

.....

.....

(d) Dextran solutions are used to treat cases of very low blood pressure caused by excessive blood loss. When dextran solutions are injected into the blood, they significantly reduce the plasma solute potential.
Use your knowledge of tissue fluid formation and osmosis to conclude how dextran solutions help to restore blood pressure. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

A400U101
09

14

© WJEC CBAC Ltd.

Examiner only

3. For healthy and productive growth, plants require a range of minerals.
Calcium, sulphur, nitrogen, phosphorus and magnesium are termed macronutrients, as plants require relatively high levels of them.
Some elements are only required in low levels; these are called micronutrients.

(a) (i) State which **two** of the macronutrients listed above are required to produce the amino acids involved in cross-linking polypeptide chains. [2]

....

(ii) State **one** role of each of the following elements **in plants**. [2]

calcium .....

magnesium .....

(b) An experiment was carried out to investigate the effects of some micronutrient deficiencies on duckweed (*Lemna minor*). This is a simple, aquatic plant that floats on the surface of ponds. It grows rapidly and individual plants can be counted easily. **Image 3.1** shows some *L. minor* plants.

Image 3.1



Five different nutrient media were tested. One contained complete nutrient medium containing all the minerals required for healthy growth. The others contained the same nutrient medium but with a source of one micronutrient omitted. The method used to set up the culture vessels is described below.

- wrap the sides of a 250 cm^3 beaker with aluminium foil to prevent entry of light
- place 150 cm^3 of nutrient medium in the beaker
- place two *L. minor* plants on the surface of the nutrient medium
- cover the top of the beaker with clear plastic film
- place the beaker in a room kept at a constant 20 °C under continuous light
- after 14 days
    1. record the colour of the leaves on a scale of 0–5 where 0 is complete chlorosis and 5 is normal dark green
    2. record the size of the leaves as T = tiny, S = small, M = medium and L = large
    3. harvest the plants and dry to constant mass
- repeat the whole experiment to gain five sets of data and calculate mean results for each nutrient medium tested

© WJEC CBAC Ltd.

Examiner only

(i) The beaker was wrapped in aluminium foil to reduce the growth of algae in the nutrient medium.
Explain how this increases the validity of the investigation. [2]

.....

.....

.....

.....

(ii) Explain **one** reason why the top of the beaker was covered with clear plastic film. [2]

.....

.....

.....

.....

(iii) Suggest the colour of leaves given a leaf colour score of 0. [1]

.....

© WJEC CBAC Ltd.

Turn over.


Examiner only

The results of the experiment are shown in **Table 3.2**.

Table 3.2

Nutrient medium – micronutrient omitted	Mean results		
	dry mass / mg	leaf colour score	leaf size
none	13.2	5	L
copper	10.6	4	M
iron	2.5	1	S
manganese	3.1	2	T
zinc	12.1	4	M

(iv) Use **Table 3.2** to plot a **bar chart** of the mean dry mass of *L. minor* harvested for each nutrient medium on the grid below. [3]



© WJEC CBAC Ltd.

Examiner only

(v) Iron is essential for the transfer of electrons from photosystem I to the final electron acceptor.
Manganese is involved in reactions that result in the photolysis of water.

Use the information above and your knowledge of the light-dependent and light-independent stages of photosynthesis to explain why a deficiency of iron and manganese results in a major decrease in growth of *L. minor* plants. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


17

© WJEC CBAC Ltd.

Examiner only

4. (a) Planetary boundaries represent the central concept in an Earth system framework proposed by scientists. **Image 4.1** shows the status of the nine planetary boundaries in 2022.

Image 4.1



(i) State the expected consequence of crossing the safe threshold for any planetary boundary. [1]

.....

.....

© WJEC CBAC Ltd.

Examiner only

(ii) The threshold for climate change has been crossed. This has been linked to human activities, including deforestation.
With reference to the role of **photoautotrophs** and **saprotrophs** in the carbon cycle, explain how deforestation could account for the status of the climate change planetary boundary shown in **Image 4.1**. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

A400U101
15

Question continued overleaf

© WJEC CBAC Ltd.


Examiner only

(b) The chemical pollution boundary includes a wide variety of chemicals released into the environment by humans. To meet the demands of an increasing human population, intensive farming is used to increase food production.

'Aphide' is a pesticide which is sprayed onto a wide range of crops to control aphids. Aphids feed on the contents of phloem sieve tubes and can spread fungal disease between plants.

Image 4.2 shows the feeding relationships in a community of interdependent species that includes aphids.

Image 4.2



(i) State what the arrows represent in **Image 4.2**. [1]

.....

(ii) Identify the following in **Image 4.2**:

I. an organism that is both a herbivore and a detritivore [1]

.....

II. an omnivore [1]

.....

(iii) Identify the original source of the energy transferred between organisms. [1]

.....

© WJEC CBAC Ltd.

Examiner only

(c) 'Aphide' acts by inhibiting the hydrolysis of acetylcholine in synapses. At low concentrations 'Aphide' affects only aphids. In high concentrations it can affect aphid predators and other organisms, including humans.

(i) Use the information given to explain why it is recommended that all fruit and vegetables should be washed before being eaten. [2]

.....

.....

.....

.....

(ii) In rats, it was found that the maximum 'safe' daily intake of 'Aphide' was $0.035\,mg\,kg^{-1}$. This is the highest dose that could be ingested daily without causing adverse effects to health.
Calculate the total mass of 'Aphide' that an adult human weighing 95 kg could ingest per day with no impact to their health.
Give your answer in grams and in standard form. [3]

Total mass of 'Aphide' = ... $g\,day^{-1}$

(iii) Suggest **one** disadvantage **to humans** of using rats to determine the maximum 'safe' daily intake of 'Aphide' for a human. [1]

.....

.....

© WJEC CBAC Ltd.

Examiner only

(d) The presence and concentration of 'Aphide' can be monitored using a biosensor. This includes immobilised molecules of the enzyme laccase.

(i) State **two** advantages of using **immobilised enzymes** in a biosensor. [2]

I. .....

.....

II. .....

.....

(ii) Laccase is a soluble, globular protein. A ribbon diagram of its structure is shown in **Image 4.3**.

Image 4.3



Identify the secondary protein structure labelled **X** in **Image 4.3** and name the type of bond involved in maintaining this structure. [2]

X = ...

Bond = ...

© WJEC CBAC Ltd.

Examiner only

(iii) There are over 350 000 **synthetic** chemicals in production globally. At present only a very small fraction of these can be identified using biosensors.
Use your knowledge of enzymes to suggest why it is proving difficult to develop biosensors to identify these chemicals. [2]

.....

.....

.....

.....

21

© WJEC CBAC Ltd.

Examiner only

5. In eukaryotic cells most of the ATP is synthesised in the mitochondria. Only a small number of ATP molecules are synthesised in the cytoplasm.

(a) (i) Name the stage of respiration that takes place in the cytoplasm. [1]


.....

(ii) State the type of reaction that produces ATP in the cytoplasm. [1]

.....

Image 5.1 shows the stages of respiration that take place in the mitochondria.

Image 5.1



(iii) Name **two** types of enzyme involved in the conversion of pyruvate to acetyl CoA in the stage labelled **X** in **Image 5.1**. [1]

....

© WJEC CBAC Ltd.

Examiner only

(iv) Triacetin is a triglyceride that can be used as an alternative respiratory substrate to glucose. From each molecule of triacetin, four molecules of acetyl CoA are formed.
Use **Image 5.1** and your knowledge of aerobic respiration to calculate the number of molecules of ATP that could be synthesised as a result of the breakdown of one molecule of triacetin. [2]

Number of molecules of ATP = ...


Question continued overleaf

© WJEC CBAC Ltd.

Examiner only

(b) **Image 5.2** shows the formulae of citrate and isocitrate, two of the molecules formed during Krebs cycle.

Image 5.2



citrate **isocitrate**

(i) Explain why the molecules shown in **Image 5.2** are structural isomers. [1]

.....

.....

(ii) Both citrate and isocitrate contain three of the same functional group.
This functional group is also found in amino acids.
State the name and formula of this functional group. [1]

Name .....


Formula .....

© WJEC CBAC Ltd.

Examiner only

(c) Isocitrate is converted to α-ketoglutarate by the enzyme isocitrate dehydrogenase (IDH). NAD is also converted to NADH in this reaction as shown in **Image 5.3**.

Image 5.3



NAD NADH

(i) Use information from **Image 5.3** to state whether NAD is being oxidised or reduced in the reaction labelled **X**. Explain your answer. [2]

.....


.....

.....

.....

(ii) **Image 5.4** shows the conversion of isocitrate to α-ketoglutarate.

Image 5.4



Use information from **Images 5.3** and **5.4** to explain how NADH can act as a competitive inhibitor of IDH and regulate the production of α-ketoglutarate. [3]

.....

.....

.....

.....

.....

.....

© WJEC CBAC Ltd.

Examiner only

(d) An investigation was carried out to determine how the concentration of NADH affects the rate at which isocitrate is converted to α-ketoglutarate.
The same concentration of IDH was used throughout. All solutions were at pH 7.4 and 25 °C.

(i) Explain how **and** why the pH was maintained at pH 7.4. [2]

.....

.....

.....

The inhibitory effect of NADH on the activity of IDH is given as percent inhibition (I%).
Table 5.5 shows the rates of reaction of IDH and I% for a range of NADH concentrations.

Table 5.5

Concentration of NADH / μmol dm^{-3}	Rate of reaction / μmol minute^{-1} mg^{-1} IDH	I% (percentage inhibition) / %
0	79	0
25	64	19
50	51	.....
100	39	51
150	25	68
200	16	80
250	10	87
300	7	91

© WJEC CBAC Ltd.

Examiner only

(ii) Use the following equation and data from **Table 5.5** to calculate the percentage inhibition (I%) for a concentration of 50 μmol NADH. [2]

$$I\% = \frac{(V_0 - V)}{V_0} \times 100$$

where V_0 = the rate of reaction in the absence of NADH,
and V = the rate of reaction in the presence of NADH

I% = ...

(iii) The rate at which NADH transports hydrogen to the electron transport chain increases as the concentration of ATP in the matrix decreases. ATP cannot be stored in cells. However, demand for ATP varies at different times of the day and the rate of ATP synthesis varies to meet demand.

Suggest how the inhibitory effect of NADH on IDH is involved in regulating the rate at which ATP is synthesised when **an organism's activity levels are low**. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

19

© WJEC CBAC Ltd.

Examiner only

6. In any ecosystem, population numbers vary over time due to many factors. These factors can be biotic or abiotic and their effects can be density dependent or density independent. Increasing human pressures on the environment have increased the risk of extinction for many species.

(a) (i) State the term used to describe a species that is at risk of extinction. [1]

.....

(ii) **Table 6.1** lists three factors that could affect the size of a population.

Table 6.1

Factor	Abiotic	Biotic	Density-dependent	Density-independent
Atmospheric temperature				
Disease				
Drought				

Place a tick (✓) in **two** boxes in each row in **Table 6.1** to correctly identify if these factors are **abiotic** or **biotic** and whether their effects are **density-dependent** or **density-independent**. [2]

(b) The population size of many species is now monitored to inform political decisions that could support conservation. The data collected is used to predict the time taken for a species to be no longer at risk of extinction.

The numbers of African elephants (*Loxodonta africana*) decreased rapidly in the 20th century, largely due to human activities. Conservation efforts are helping to restore population numbers in some areas of Africa.

(i) Suggest **two** human activities that could have contributed to the decline in the number of African elephants. [1]

.....

.....

© WJEC CBAC Ltd.

Examiner only

(ii) Under optimal conditions, African elephants have a generation time of 22 years. This is the time taken for a population to double in number.
Generation time (G) is calculated using the formula:

$$\text{Generation time (G)} = \frac{t}{3.3 \times \log_{10}\left(\frac{N_{final}}{N_{start}}\right)}$$

where:

t = time
N_{final} = the final number of individuals in the population
N_{start} = the initial number of individuals in the population

Use the formula above to estimate the time it would take for a starting population of 10 elephants to reach a population of 1000.
Give your answer to two significant figures. [3]

Time taken to reach final population of 1000 = ... years

(iii) One conservation method used has been to set up enclosed areas where elephants are protected.
If the area is not enclosed, it could lead to a decrease in the time taken to reach the target population of 1000 elephants. Suggest why. [1]

.....

.....

8

© WJEC CBAC Ltd.

7. Succession is the change in structure and species composition of a community over time. Primary succession from bare rock to a mature community can take hundreds of years. Secondary succession, for example, following the felling of trees, takes far less time.

Forests are habitats that can support a high level of biodiversity. In some countries, forests are given nature conservation status and are unmanaged. There is limited harvesting of wood and no clearance of dead wood or undergrowth. This helps to preserve habitats.

In managed forests, secondary succession is encouraged by the regular felling of trees. The removal of dead and decaying wood and undergrowth in these forests reduces the spread of disease.

Evidence suggests that by using sustainable forest management practices, the demand for timber and the need for conservation can be met.

Table 7 compares data from an unmanaged nature conservation forest and a managed timber production forest. Both forests contained similar tree species.

Table 7

	Type of forest	
	unmanaged nature conservation forest	sustainably managed timber production forest
increase in volume of wood / m^3 hectare^{-1} year^{-1}	2	15
maximum volume of wood / m^3 hectare^{-1}	624	757
time for trees to reach felling height / years	175	50
useful wood harvest / million m^3 year^{-1}	10–15	56–86
carbon storage / tonnes CO_2 hectare^{-1} year^{-1}	0.4	3.2–3.5
increase in species biodiversity over time / %	0	0.005

© WJEC CBAC Ltd.

Examiner only

Use correct terminology to describe the stages in primary succession that could result in the initial formation of a forest from bare rock.
Explain why biodiversity increases as the forest community matures.
With reference to the information opposite, explain how managed forest succession can meet the need for timber production and be of benefit to the environment. [9 QER]

© WJEC CBAC Ltd.

Examiner only

© WJEC CBAC Ltd.

Examiner only

END OF PAPER

9

© WJEC CBAC Ltd.

Question number	Additional page, if required. Write the question number(s) in the left-hand margin.

Examiner only

© WJEC CBAC Ltd.