

GCE

Chemistry A

H432/03: Unified chemistry

Advanced GCE

Mark Scheme for November 2020

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.


© OCR 2020

Annotations

Annotation	Meaning
	Correct response
	Incorrect response
	Omission mark
	Benefit of doubt given
	Contradiction
	Rounding error
	Error in number of significant figures
	Error carried forward
	Level 1
	Level 2
	Level 3
	Benefit of doubt not given
	Noted but no credit given
	Ignore

Abbreviations, annotations and conventions used in the detailed Mark Scheme (to include abbreviations and subject-specific conventions).





Annotation	Meaning
DO NOT ALLOW	Answers which are not worthy of credit
IGNORE	Statements which are irrelevant
ALLOW	Answers that can be accepted
()	Words which are not essential to gain credit
—	Underlined words must be present in answer to score a mark
ECF	Error carried forward
AW	Alternative wording
ORA	Or reverse argument



Question	Answer	Marks	AO element	Guidance
1 (a) (i)	 Water flow AND condenser Water in at bottom and out at top AND condenser ✓ Flask and technique Pear-shaped/round-bottom flask AND reflux ✓	2	1.2 × 2	DO NOT ALLOW conical flask, volumetric flask, beaker in place of round bottom/pear shaped flask

Question			Answer	Marks	AO element	Guidance
		(ii)	Diagram showing knowledge of filtration under reduced pressure Diagram showing Buchner flask <i>must have <u>ONE</u> side arm</i> AND Buchner/Hirsh funnel on top of flask ✓ <i>Labels not required</i> ----- Further details: • Funnel sealed or stoppered to flask AND • Apparatus capable of filtering under reduced pressure AND • Label for setup from side arm to indicate reduced pressure AND • Label for Buchner flask OR Buchner/Hirsh funnel ✓ <i>ALLOW slips in spelling of 'Buchner'</i>	2		Labels NOT required for diagram ALLOW diagram of a conical flask with a filtering setup above AND Side arm either in conical flask OR between flask and filter paper of funnel IGNORE absence of seals ----- MUST imply some type of seal between filter setup and flask. ALLOW <u>small</u> gaps Examples of suitable labels (may have arrow from side arm or tube attached) • to pump • to vacuum • air out • suction • reduced pressure • etc. For Buchner flask and Buchner funnel DO NOT ALLOW just 'flask OR 'funnel' <i>Flask and funnel used in normal filtration</i>

Question			Answer	Marks	AO element	Guidance
	(b)	(i)	Comparison of branching and points of contact e.g. $\text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2$ has longer chain / straight chain / no branches AND e.g. $\text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2$ has more points of contact / more surface interaction (between molecules) ✓ Relative strength of force e.g. $\text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2$ has stronger/more induced dipole(–dipole) interactions OR London forces ✓ ----- Hydrogen bonds $\text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2$ OR $(\text{CH}_3)_2\text{CHNH}_2$ have hydrogen/H bonds OR $(\text{CH}_3)_3\text{N}$ has no hydrogen/H bonds ✓ Relative strength of force Hydrogen bonds are stronger than London forces /permanent dipole interactions ✓ ----- Comparison of energy required to break force e.g. More energy to break/overcome London forces/intermolecular forces in $\text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2$ OR More energy is needed to break H bonds (than London forces) ✓	5 → 4 max		ANNOTATE WITH TICKS AND CROSSES, etc. ----- ALLOW ORA throughout ALLOW 'The straighter the chain, the more points of contact' 1.2 IGNORE comparison using 'primary', 'secondary' and 'tertiary'. <i>Comparison of branching is required.</i> 2.1 For London forces, • ALLOW induced dipole(–dipole) interactions • IGNORE IDID OR van der Waals' forces/VDW DO NOT ALLOW $\text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2$ has more electrons <i>(number of electrons are the same)</i> 1.2 2.1 DO NOT ALLOW 'more energy to break covalent bonds' ALLOW little energy is required to break London forces (compared with H bonds)

Question			Answer	Marks	AO element	Guidance
	(b)	(ii)	FIRST CHECK MOLECULAR FORMULA and STRUCTURE IF molecular formula = C₅H₁₃N AND correct structure AND evidence of ideal gas equation → 6 marks Correct up to 87 AND C₅H₁₃N → 5 marks Correct up to 87 → 4 marks Rearranging ideal gas equation $n = \frac{pV}{RT} \checkmark$ Unit conversion AND substitution into $n = \frac{pV}{RT}$: • $R = 8.314$ OR 8.31 • $V = 72(.0) \times 10^{-6}$ • T in K: 373 K e.g. $\frac{1.00 \times 10^5 \times 72.0 \times 10^{-6}}{8.314 \times 373} \checkmark$ Calculation of n $n = 2.32 \times 10^{-3} \text{ (mol)} \checkmark$ Calculation of M $M = \frac{0.202}{2.32 \times 10^{-3}} = 87 \checkmark$ Molecular formula C₅H₁₃N $\checkmark$ Molecular formula required	6	2.2×4 3.2	IF $n = \frac{pV}{RT}$ is omitted, ALLOW when values are substituted into rearranged ideal gas equation. Calculator: $n = 2.321740325 \times 10^{-3}$ from 8.314 From 8.31, $n = 2.322857889 \times 10^{-3}$ ALLOW elements in any order ALLOW molecular formula = C₃H₉N₃ ALLOW other molecular formulae of an amine that has $M = 87$, e.g. C₄H₉NO

Question	Answer	Marks	AO element	Guidance
	Structure of amine A from $C_5H_{13}N$ ✓  OR 		3.2	ALLOW any combination of skeletal OR structural OR displayed formula as long as unambiguous ALLOW structures below from molecular formula = $C_3H_9N_3$  ALLOW ECF but only if structure has calculated M_r AND has 3 peaks in ^{13}C NMR spectrum.
Use of 24000	3 marks max possible for use of 72.0 cm^3 OR 0.720 dm^3 by ECF Calculation e.g. $n = \frac{72.0}{24000} = 3.00 \times 10^{-3}$ No mark (calculation much simpler) $M = \frac{0.202}{3.00 \times 10^{-3}} = 67.3$ OR 67 ✓ ECF Molecular formula = C_4H_5N ✓ ECF Structure  ✓ ECF			

Question			Answer	Marks	AO element	Guidance
	(c)		 Organic product and water marked independently. <i>1st mark</i> correct organic product OR water IGNORE balancing numbers <i>2nd mark</i> BOTH products AND correctly balanced.	2	3.2	ALLOW any combination of skeletal OR structural OR displayed formula as long as unambiguous ALLOW  NOTE: For ECF, any structure must have correct number of bonds to C, H, O and N DO NOT ALLOW structure of dimer <i>Question states molecular formula = C₃H₃NO</i>
			Total	16		

Question	Answer	Marks	AO element	Guidance
2*	Refer to marking instructions on page 5 of mark scheme for guidance on marking this question. Level 3 (5-6 marks) Comprehensive explanation of the terms, ligand and coordination number and ligand substitution AND 3D diagrams of suitable examples of 6 AND 4 coordinate complex ions with different shapes AND Ligand substitution illustrated with a balanced equation <i>There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated.</i> Level 2 (3-4 marks) Explanation of the terms, ligand and coordination number and ligand substitution with some errors or omissions AND: Diagrams of suitable examples of 6 AND 4 coordinate complex ions with different shapes OR A 3D wedged diagram of a suitable example of 6 OR 4 coordination OR A diagram of a suitable example of 6 OR 4 coordination AND ligand substitution illustrated with an equation OR Ligand substitution illustrated with a balanced equation <i>There is a line of reasoning presented with some structure. The information presented is relevant and supported by some evidence</i>	6	1.1×4 2.1×2	Indicative scientific points may include: Terms Ligand: Donates a lone pair to metal ion Forms dative covalent (coordinate) bond with metal ion Coordination number: Number of coordinate bonds to metal ion. Could be implicit in annotated diagrams NOTE: For monodentate ligands, 'number of ligands' is the same as the number of coordination number Ligand substitution: One ligand replacing another <u>Suitable examples of complex ions with different shapes</u> Coordination no 6 Octahedral e.g. $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$, $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$ Coordination no 4 Tetrahedral e.g. CuCl_4^{2-}, CoCl_4^{2-} OR Square planar Pt complexes, e.g. $\text{Pt}(\text{NH}_3)_2\text{Cl}_2$ <u>Diagrams and equations</u> Diagrams of complex ions (may be 3D) Equation for ligand substitution e.g. $[\text{Cu}(\text{H}_2\text{O})_6]^{2+} + 4\text{Cl}^- \rightarrow \text{CuCl}_4^{2-} + 6\text{H}_2\text{O}$ $[\text{Cu}(\text{H}_2\text{O})_6]^{2+} + 4\text{NH}_3 \rightarrow [\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+} + 4\text{H}_2\text{O}$ NOTE: A clear and logically structured response would link shapes with some of: coordination number, names of shapes, connectivity, involvement of lone pairs, bond angles, etc. (not inclusive) ALLOW minor slips NOTE: Levels and the mark within a level is a 'best-fit', not perfection

Question			Answer	Marks	AO element	Guidance
			Level 1 (1-2 marks) Explanation of some terms: ligand, coordination number and ligand substitution with some errors or omissions. AND A suitable example of a complex ion OR Ligand substitution illustrated with an equation with some errors <i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant.</i> 0 marks No response or no response worthy of credit.			
			Total	6		

Question			Answer	Marks	AO element	Guidance
3	(a)	(i)	FIRST, CHECK THE ANSWER ON ANSWER LINE IF $\Delta_c H = -1860$ OR -1850 (kJ mol^{-1}) with evidence of working, award 3 marks IF $\Delta_c H = -1862$, award 2 marks (not 3 SF) ----- Energy released in J OR kJ $= 100 \times 4.18 \times 24.5 = \pm 10241$ (J) OR ± 10.241 (kJ) ✓ 3 SF minimum required Calculates $n(\text{C}_3\text{H}_8)$ $= \frac{0.242}{44(.0)} = 0.0055(0)$ (mol) ✓ Calculates $\Delta_c H$ with – sign AND 3 SF (appropriate) $\Delta_c H = \frac{10241}{0.0055 \times 1000} = -1862$ No mark $= -1860$ OR -1.86×10^3 (kJ mol^{-1}) ✓ – sign AND 3 SF required	3		FULL ANNOTATIONS MUST BE USED ----- ALLOW ECF throughout DO NOT ALLOW $c = 4.2 \rightarrow 10290$ Next 2 marks available by ECF $\rightarrow -1870$ ALLOW 10240/10200 J OR 10.24/10.2 kJ IGNORE units ALLOW ECF from initial 3 SF rounding to 10.2 kJ: $\pm \frac{10200}{0.0055 \times 1000} \rightarrow \pm 1854.545455$ ✓ $\rightarrow 1850$ ✓ ----- Common errors $\Delta H = -54.6$ OR -54.7 2 marks by ECF from $mc\Delta T$ m wrong as 0.242 and ΔT wrong as 297.5 K) $\rightarrow mc\Delta T$ wrong as 300.9391 (J) $\Delta H = -4.51$ 2 marks by ECF from $mc\Delta T$ m wrong as 0.242 and ΔT correct as 24.5) $\rightarrow mc\Delta T$ wrong as 24.78322 (J) $\Delta H = -22600$ 2 marks by ECF from $mc\Delta T$ m correct as 100 and ΔT wrong as 297.5) $\rightarrow mc\Delta T$ wrong as 124355 (J)
	(a)	(ii)	Any two from: 1 MARK ONLY ✓ - Heat loss/released to surroundings - Incomplete combustion/reaction with oxygen or air OR not everything burns - Evaporation of water	1	1.2	IGNORE incomplete 'reaction' Needs link to combustion/burning/reaction with air/O₂ IGNORE evaporation of C₃H₈


Question	Answer	Marks	AO element	Guidance
(b)*	Refer to marking instructions on page 5 of mark scheme for guidance on marking this question. Level 3 (5-6 marks) Calculates $\Delta_r H$ for reaction 3.1 correctly with correct sign AND Calculates a value for $\Delta_c H^\circ$ of propane using $\Delta_r H$ AND $\pm 4 \times \Delta_{\text{vap}} H$ <i>There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated.</i> Level 2 (3-4 marks) Calculates $\Delta_r H$ for reaction 3.1 correctly with correct sign OR Calculates bonds broken OR bonds made correctly to obtain a value of $\Delta_r H$ for reaction 3.1 AND attempts to link $\Delta_r H$ with $\Delta_{\text{vap}} H$ OR calculates $4 \times \Delta_{\text{vap}} H$ <i>There is a line of reasoning presented with some structure. The information presented is relevant and supported by some evidence.</i> Level 1 (1-2 marks) Uses bond enthalpies for bonds broken and bonds made but may contain errors or omissions AND obtains a value for $\Delta_r H$. OR Calculates bonds broken OR bonds made correctly. <i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant.</i> 0 marks No response or no response worthy of credit.	6	2.4×2 3.1×2 3.2×2	Indicative scientific points may include: Bond enthalpy calculation of $\Delta_r H$ Bonds broken $= (2 \times 347) + (8 \times 413) + (5 \times 498)$ $= (694) + (3304) + (2490)$ $= \pm 6488 \text{ kJ mol}^{-1}$ Bonds made $= (6 \times 805) + (8 \times 464)$ $= (4830) + (3712) = \pm 8542 \text{ kJ mol}^{-1}$ $\Delta_r H = 6488 - 8542 = -2054 \text{ kJ mol}^{-1}$ NOTE: 3 C–C → 6835 for bond broken: $\Delta H = -1707$ 2 C–C omitted from bonds broken gives: $\Delta H = -2748$ ----- Determination of $\Delta_c H(\text{C}_3\text{H}_8)$ $\Delta_c H^\circ$ of propane using $\Delta_r H$ AND $\pm 4 \times \Delta_{\text{vap}} H$ Correct $\Delta_c H(\text{C}_3\text{H}_8) = \Delta_r H - 4 \times \Delta_{\text{vap}} H$ $= -2054 - (4 \times 40.65)$ $= -2054 - 162.6$ $= -2216.6 / -2217 \text{ kJ mol}^{-1}$ Incorrect $\Delta_c H(\text{C}_3\text{H}_8) = -2054 + (4 \times 40.65)$ $= -2054 + 162.6$ $= -1891.4 / -1891 \text{ kJ mol}^{-1}$ NOTE: A clear and logically structured response would include a correct energy cycle for $\Delta_c H(\text{C}_3\text{H}_8)$ using $\Delta_r H$ AND $4 \times \Delta_{\text{vap}} H$ in energy cycle or expression: ALLOW trailing zeroes OR minor slips
	Total	10		

Question			Answer	Marks	AO element	Guidance
4	(a)	(i)	Overall equation AND state symbols: $M(s) + 2HCl(aq) \rightarrow MCl_2(aq) + H_2(g)$ ✓ STATE SYMBOLS required in overall equation ONLY Half equations: Oxidation $M \rightarrow M^{2+} + 2e^-$ ✓ Reduction $2H^+ + 2e^- \rightarrow H_2$ OR $H^+ + e^- \rightarrow \frac{1}{2}H_2$ ✓	3	2.6×3	All 3 marks are independent. IGNORE charges/oxidation numbers shown around overall equation. <i>Treat as rough working</i> ALLOW overall equation shown with some or all ions that are present e.g. (with state symbols) $M + 2H^+ \rightarrow M^{2+} + H_2$ $M + 2HCl \rightarrow M^{2+} + 2Cl^- + H_2$ $M + 2H^+ + 2Cl^- \rightarrow M^{2+} + 2Cl^- + H_2$ In half equations, IGNORE state symbols even is wrong BUT half equations MUST only have species that change. For charges on half equations, ALLOW M^{+2} for M^{2+} OR H^{+1} for H^+ ALLOW $M - 2e^- \rightarrow M^{2+}$ If BOTH half equations are correct but shown with oxidation and reduction the wrong way around, award 1 mark from the 2 marks for half equations
	(a)	(ii)	Bubbles/effervescence/fizzing stops ✓ M/metal/solid has disappeared/dissolved ✓	2	3.3×2	Responses must imply that all fizzing has stopped and that all the solid has dissolved i.e. 'metal disappears' is not quite enough. 'All the metal disappears' is enough IGNORE constant mass IGNORE no increase in temperature
	(a)	(iii)	$H^+ + OH^- \rightarrow H_2O$ ✓	1	2.5	ALLOW multiples e.g. $2H^+ + 2OH^- \rightarrow 2H_2O$ IGNORE state symbols, even if wrong



Question			Answer	Marks	AO element	Guidance
	(a)	(iv)	Mean titre 1 mark $= \frac{(27.30 + 27.20)}{2} = 27.25 \text{ (cm}^3\text{)} \checkmark$ Analysis of results 5 marks $n(\text{NaOH}) = 27.25 \times \frac{0.320}{1000} = 8.72 \times 10^{-3} \text{ (mol)} \checkmark$ $n(\text{HCl}) \text{ in } 25.0 \text{ cm}^3 = n(\text{NaOH})$ $n(\text{HCl}) \text{ in } 250 \text{ cm}^3 = 8.72 \times 10^{-3} \times 10 = 8.72 \times 10^{-2} \text{ (mol)} \checkmark$ $n(\text{HCl}) \text{ that reacted with M} = 0.210 - 8.72 \times 10^{-2} = 0.1228 \text{ (mol)} \checkmark$ $n(\text{M}) \text{ that reacted} = \frac{0.1228}{2} = 0.0614 \text{ (mol)} \checkmark$ $A_r \text{ of M} = \frac{6.90}{0.0614} = 112.4 \text{ AND M} = \text{cadmium/Cd} \checkmark$	6	2.8×5 3.2	FULL ANNOTATIONS MUST BE USED Common error: Incorrect mean from all 3 titres = 27.35 cm³ Use ECF throughout Intermediate values for working to at least 3 SF. TAKE CARE: Value written down may be truncated calculator value. Depending on rounding, either can be credited. ALLOW 0.123 (mol) i.e. 3SF ALLOW 0.0615 (mol) IF 0.1228 rounded to 0.123 ALLOW 112.2 from 0.0615 AND Cd ALLOW A_r to nearest whole number ALLOW ECF for metal closest to calculated A_r DO NOT ALLOW Ga OR Sc (Form 3+ ions only)
			COMMON ERRORS: Mean of 27.35 (use of all 3 titres) → 8.752 × 10⁻³ → 8.752 × 10⁻² → 0.12248 → 0.06124 → 112.7 AND Cd: 5 marks No ÷2 to obtain n(M) → 56.2 AND Fe (from 27.25) 5 marks → 56.3 AND Fe (from 27.35) 4 marks No subtraction from 0.210 → 8.72 × 10⁻²/2 → 4.36 × 10⁻² → $\frac{6.90}{4.36 \times 10^{-2}}$ → 158.2 to 158.3 AND Tb 5 marks			No ×10 to obtain n(HCl) in 250 cm³ 5 marks 0.210 – 8.72 × 10⁻³ = 0.20128 OR 0.201 n(M) = 0.20128/2 = 0.10064 A_r = 6.90/0.10064 = 68.56 → Zn No ×10 and no ÷ 2 4 marks 0.210 – 8.72 × 10⁻³ = 0.20128 A_r = 6.9/0.20128 = 34.28 → Ca Omitting initial titration calculation Zero marks 0.210/2 = 0.105 → 6.9/0.105 = 65.71 → Zn

Question			Answer	Marks	AO element	Guidance
	(b)	(i)	$n(\text{CO}_2) = \frac{2.75}{44} = 0.0625 \text{ (mol)} \checkmark$	1	2.8	
	(b)	(ii)	$n(\text{X}_2\text{CO}_3) = 0.0625 \text{ (mol)}$ OR 0.0625 used in molar mass expression below $\checkmark$ $\text{Molar mass of X}_2\text{CO}_3 = \frac{14.57}{0.0625} = 233.12 \text{ (g mol}^{-1}\text{)} \checkmark$ Metal X = Rubidium/Rb $\checkmark$	3	1.2 2.8 3.2	ALLOW ECF from 4b(i) ALLOW to nearest whole number DO NOT ALLOW strontium/Sr <i>wrong carbonate formula</i> ALLOW ECF for X from calculated molar mass ONLY IF X is a Group 1 metal OR Ag Working: Mass of X in X_2CO_3 = 233.14 – 60 = 173.12 OR 173 A_r of X = $\frac{173.12}{2}$ OR 86.56 OR 85.6 OR 87
	(c)	(i)	Reweigh to constant mass $\checkmark$	1	3.4	ALLOW response implying leaving for longer and monitoring by reweighing to constant mass, e.g. Leave flask until the mass does not change IGNORE ‘leave for longer’ OR wait till fizzing stops <i>Needs link to constant mass</i> ALLOW Collect gas until gas volume is constant
	(c)	(ii)	Mass (CO_2) OR $n(\text{CO}_2)$ loss would be smaller OR Mass X_2CO_3 OR $n(\text{X}_2\text{CO}_3)$ reacted (seems to be) less $\checkmark$ Molar mass would be greater $\checkmark$	2	3.1 3.2	
			Total	19		

Question		Answer					Marks	AO element	Guidance		
5	(a)		T / K	500	600	700	800	2	1.2×2	Mark by row ALLOW 2 SF or more for $1/T$ but ignore trailing zeroes ALLOW whole numbers (~~±1~~) for $\ln K_p$ ALLOW 1 small slip in each row. e.g. 1.66 for 1.67; 71.7 for 71.8 <i>Check with calculator values below table</i> BUT DO NOT ALLOW whole number errors, e.g. 85 for 86 ☒	
			K_p	5.86×10^{45}	1.83×10^{37}	1.46×10^{31}	1.14×10^{26}				
			$\frac{1}{T} / K^{-1}$	2.00×10^{-3}	1.67×10^{-3}	1.43×10^{-3}	1.25×10^{-3}				✓
			$\ln K_p$	105	86	72	60				✓
			<i>Calculator values</i> $1/T / 10^{-3}$ 2.00 1.66 recurring 1.428571429 1.25 $\ln K_p$ 105.3844788 85.79996441 71.75857432 59.99824068								
	(b)		Equilibrium (position) shifts to the left AND (forward) reaction is exothermic ✓					1	2.2	ALLOW 'favours reverse reaction' <i>Implies shift to left</i> ALLOW 'shifts in endothermic direction' BUT only if (forward) reaction stated as exothermic	

Question		Answer	Marks	AO element	Guidance
(c)		Plotting of graph All points correctly plotted AND best-fit straight line ✓ Gradient Correct gradient of best-fit straight line within the range $\pm 57000 \rightarrow \pm 63000$ ✓ ΔH calculation (subsumes mark for gradient) $\Delta H = (-) \text{gradient} \times 8.31(4)$ OR calculated value ✓ <i>e.g. from ± 60000, $\Delta H = (+) 498840$ (J) OR ± 498.840 (kJ)</i> ΔH in kJ mol^{-1} ΔH correct in kJ mol^{-1} AND 3SF AND – sign ✓ <i>e.g. from ± 498840, $\Delta H = -499$ (kJ mol^{-1})</i>	4	3.1  3.1 3.2 3.2	ALLOW 4 points on graph Tolerance 1 small square ALLOW ΔH in range: $-480 \rightarrow -530$ (kJ mol^{-1}) This mark subsumes gradient mark
(d)		Extrapolate line to (y) intercept OR Measure/Use (y) intercept ✓ Intercept = $\frac{\Delta S}{R}$ OR $\Delta S = R \times (\text{y}) \text{ intercept}$ ✓ <i>This statement automatically subsumes 1st mark</i> NOTE: If ‘x’ intercept, DO NOT ALLOW 1st mark but 2nd mark available for $\times R$ as BOD	2	3.1×2	ALLOW substitute values of $\ln K_p$, $1/T$ and gradient into Equation 5.1 ✓ From provided values and gradient = 60000: $\frac{\Delta S}{R} = \ln K_p - \text{gradient} \times 1/T$ OR $135 - 60000 \times 2.50 \times 10^{-3} = -15$ ✓
Total			9		

Question			Answer	Marks	AO element	Guidance						
6	(a)		<table><tr><td>Bond angle</td><td>Name of shape</td></tr><tr><td>120(°)</td><td>Trigonal planar</td></tr><tr><td>104–105(°)</td><td>Non-linear</td></tr></table> Mark by row OR by column to give higher mark i.e. 2 bond angles correct ✓ 2 shapes correct ✓ OR i.e. bond angle AND shape correct in 1st row ✓ bond angle AND shape correct in 2nd row ✓	Bond angle	Name of shape	120(°)	Trigonal planar	104–105(°)	Non-linear	2	1.2×2	For non-linear, ALLOW bent, v-shaped, angular IGNORE planar, ‘not straight’
		Bond angle	Name of shape									
120(°)	Trigonal planar											
104–105(°)	Non-linear											
	(b)		CH₃SO₂OH + H₂O ⇌ CH₃SO₂O[−] + H₃O⁺ ✓ A1 B2 B1 A2 ✓ For an equilibrium shown using CH₃COOH instead of H₂O, mark acid–base pairs by ECF, i.e. CH₃SO₂OH + CH₃COOH ⇌ CH₃SO₂O[−] + CH₃COOH₂⁺ ☒ A1 B2 B1 A2 ECF ✓ CH₃SO₂OH dissociates more (than CH₃COOH) OR CH₃SO₂OH is a stronger acid ✓ ORA in terms of CH₃COOH being a weaker acid Student is correct AND (sulfonic acid has) lower p<i>K</i>_a/higher <i>K</i>_a OR greater [H⁺] ORA ✓	4	2.1×2	ALLOW → for ⇌ ALLOW acid–base pairs labelled other way round. i.e. CH₃SO₂OH + H₂O ⇌ CH₃SO₂O[−] + H₃O⁺ A2 B1 B2 A1 ALLOW small slip If ONE charge is missing from equilibrium. ALLOW ECF for acid–base pairs mark IGNORE ‘more acidic’ <i>Response needs strength/dissociation</i> ALLOW maths explanation for final 2 marks, e.g. <i>K</i>_a(CH₃COOH) = 10^{−(4.76)}} = 1.74 × 10^{−5} [H⁺] = √(1.74 × 10^{−5} × 1) = 4.17 × 10^{−3} pH = −log 4.17 × 10^{−3} = 2.38 ✓						
					3.1							
					3.2	<i>K</i>_a(CH₃SO₂OH) = 10^{−(−1.90)}} = 79.4 [H⁺] = √(79.4 × 1) = 8.91 pH = −log 8.91 = −0.95 ✓ BOTH pH calcs subsumes ‘Student is correct’						

Question	Answer	Marks	AO element	Guidance
(c)	 6 curly arrows correct ✓✓✓✓ 5 curly arrows correct ✓✓✓ 4 curly arrows correct ✓✓ 3 curly arrows correct ✓	4	3.1×4	IGNORE any added charges OR dipoles. <i>Marks solely for curly arrows</i> IGNORE any curly arrows on bottom structures (not in boxes): 
	Total	10		

OCR (Oxford Cambridge and RSA Examinations)
The Triangle Building
Shaftesbury Road
Cambridge
CB2 8EA

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored