OCR
Oxford Cambridge and RSA

Monday 3 June 2019 – Morning

A Level Further Mathematics B (MEI)

Y420/01 Core Pure

Time allowed: 2 hours 40 minutes

7768493368

You must have:
- Printed Answer Booklet
- Formulae Further Mathematics B (MEI)

You may use:
- a scientific or graphical calculator

INSTRUCTIONS
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Answer **all** the questions.
- **Write your answer to each question in the space provided in the Printed Answer Booklet.** If additional space is required, you should use the lined page(s) at the end of the Printed Answer Booklet. The question number(s) must be clearly shown.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.

INFORMATION
- The total number of marks for this paper is **144**.
- The marks for each question are shown in brackets **[ ]**.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is used. You should communicate your method with correct reasoning.
- The Printed Answer Booklet consists of **24** pages. The Question Paper consists of **8** pages.

© OCR 2019 [Y/508/5592]
DC (LK/TP) 173302/4

OCR is an exempt Charity

stemup.co.uk

Section A (34 marks)

Answer **all** the questions.

1 Find $\sum_{r=1}^{n}(2r^2-1)$, expressing your answer in fully factorised form. **[4]**

2 The plane $x+2y+cz=4$ is perpendicular to the plane $2x-cy+6z=9$, where c is a constant. Find the value of c. **[3]**

3 Matrices **A** and **B** are defined by $\mathbf{A}=\begin{pmatrix}3 & 1\\ 2 & 1\end{pmatrix}$ and $\mathbf{B}=\begin{pmatrix}k & 1\\ 2 & 0\end{pmatrix}$, where k is a constant.

(a) Verify the result $(\mathbf{AB})^{-1}=\mathbf{B}^{-1}\mathbf{A}^{-1}$ in this case. **[5]**

(b) Investigate whether **A** and **B** are commutative under matrix multiplication. **[2]**

4 **In this question you must show detailed reasoning.**

Fig. 4 shows the region bounded by the curve $y=\sec\frac{1}{2}x$, the x-axis, the y-axis and the line $x=\frac{1}{2}\pi$.




Fig. 4

This region is rotated through 2π radians about the x-axis.
Find, in exact form, the volume of the solid of revolution generated. **[3]**

5 Using the Maclaurin series for $\cos 2x$, show that, for small values of x,

$\sin^2 x \approx ax^2+bx^4+cx^6$,

where the values of a, b and c are to be given in exact form. **[5]**

© OCR 2019

stemup.co.uk

6 **In this question you must show detailed reasoning.**

Find $\int_2^\infty \frac{1}{4+x^2}\,\mathrm{d}x$. **[4]**

7 A curve has cartesian equation $(x^2+y^2)^2 = 2c^2xy$, where c is a positive constant.

(a) Show that the polar equation of the curve is $r^2 = c^2\sin 2\theta$. **[2]**

(b) Sketch the curves $r = c\sqrt{\sin 2\theta}$ and $r = -c\sqrt{\sin 2\theta}$ for $0 \leqslant \theta \leqslant \frac{1}{2}\pi$. **[3]**

(c) Find the area of the region enclosed by one of the loops in part **(b)**. **[3]**

© OCR 2019

stemup.co.uk

Section B (110 marks)

Answer **all** the questions.

8 **In this question you must show detailed reasoning.**

The roots of the equation $x^3 - x^2 + kx - 2 = 0$ are α, $\frac{1}{\alpha}$ and β.

(a) Evaluate, in exact form, the roots of the equation. **[6]**

(b) Find k. **[2]**

9 Prove by induction that $5^n + 2 \times 11^n$ is divisible by 3 for all positive integers n. **[7]**

10 **In this question you must show detailed reasoning.**

(a) You are given that $-1 + \mathrm{i}$ is a root of the equation $z^3 = a + b\mathrm{i}$, where a and b are real numbers. Find a and b. **[3]**

(b) Find all the roots of the equation in part **(a)**, giving your answers in the form $r\mathrm{e}^{\mathrm{i}\theta}$, where r and θ are exact. **[4]**

(c) Chris says "the complex roots of a polynomial equation come in complex conjugate pairs". Explain why this does **not** apply to the polynomial equation in part **(a)**. **[1]**

11 **(a)** Specify fully the transformations represented by the following matrices.

- $\mathbf{M}_1 = \begin{pmatrix} \frac{3}{5} & -\frac{4}{5} \\ \frac{4}{5} & \frac{3}{5} \end{pmatrix}$
- $\mathbf{M}_2 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ **[4]**

(b) Find the equation of the mirror line of the reflection R represented by the matrix $\mathbf{M}_3 = \mathbf{M}_1\mathbf{M}_2$. **[5]**

(c) It is claimed that the reflection represented by the matrix $\mathbf{M}_4 = \mathbf{M}_2\mathbf{M}_1$ has the same mirror line as R. Explain whether or not this claim is correct. **[3]**

12 Three intersecting lines L_1, L_2 and L_3 have equations

L_1: $\frac{x}{2} = \frac{y}{3} = \frac{z}{1}$, L_2: $\frac{x}{1} = \frac{y}{2} = \frac{z}{-4}$ and L_3: $\frac{x-1}{1} = \frac{y-2}{1} = \frac{z+4}{5}$.

Find the area of the triangle enclosed by these lines. **[9]**

© OCR 2019

stemup.co.uk

13 **(a)** Using the logarithmic form of $\operatorname{arcosh} x$, prove that the derivative of $\operatorname{arcosh} x$ is $\dfrac{1}{\sqrt{x^2-1}}$. **[5]**

(b) Hence find $\displaystyle\int_1^2 \operatorname{arcosh} x \, \mathrm{d}x$, giving your answer in exact logarithmic form. **[5]**

(c) Ali tries to evaluate $\displaystyle\int_0^1 \operatorname{arcosh} x \, \mathrm{d}x$ using his calculator, and gets an 'error'. Explain why. **[1]**

14 Three planes have equations

$$\begin{aligned} -x + ay &= 2 \\ 2x + 3y + z &= -3 \\ x + by + z &= c \end{aligned}$$

where a, b and c are constants.

(a) In the case where the planes **do not** intersect at a unique point,

(i) find b in terms of a, **[4]**

(ii) find the value of c for which the planes form a sheaf. **[3]**

(b) In the case where $b = a$ and $c = 1$, find the coordinates of the point of intersection of the planes in terms of a. **[6]**

15 **In this question you must show detailed reasoning.**

Show that $\displaystyle\int_{\frac{3}{4}}^{\frac{3}{2}} \frac{1}{\sqrt{4x^2-4x+2}} \, \mathrm{d}x = \frac{1}{2}\ln\left(\frac{3+\sqrt{5}}{2}\right)$. **[8]**

16 **(a)** Show that $(2-\mathrm{e}^{\mathrm{i}\theta})(2-\mathrm{e}^{-\mathrm{i}\theta}) = 5-4\cos\theta$. **[3]**

Series C and S are defined by

$$C = \frac{1}{2}\cos\theta + \frac{1}{4}\cos 2\theta + \frac{1}{8}\cos 3\theta + \ldots + \frac{1}{2^n}\cos n\theta,$$

$$S = \frac{1}{2}\sin\theta + \frac{1}{4}\sin 2\theta + \frac{1}{8}\sin 3\theta + \ldots + \frac{1}{2^n}\sin n\theta.$$

(b) Show that $C = \dfrac{2^n(2\cos\theta - 1) - 2\cos(n+1)\theta + \cos n\theta}{2^n(5-4\cos\theta)}$. **[9]**

© OCR 2019

stemup.co.uk

17 A cyclist accelerates from rest for 5 seconds then brakes for 5 seconds, coming to rest at the end of the 10 seconds. The total mass of the cycle and rider is m kg, and at time t seconds, for $0 \leqslant t \leqslant 10$, the cyclist's velocity is $v\,\text{m}\,\text{s}^{-1}$.

A resistance to motion, modelled by a force of magnitude $0.1mv$ N, acts on the cyclist during the whole 10 seconds.

(a) Explain why modelling the resistance to motion in this way is likely to be more realistic than assuming this force is constant. **[1]**

During the braking phase of the motion, for $5 \leqslant t \leqslant 10$, the brakes apply an additional constant resistance force of magnitude $2m$ N and the cyclist does not provide any driving force.

(b) Show that, for $5 \leqslant t \leqslant 10$, $\frac{\mathrm{d}v}{\mathrm{d}t} + 0.1v = -2$. **[1]**

(c) **(i)** Solve the differential equation in part **(b)**. **[5]**

(ii) Hence find the velocity of the cyclist when $t = 5$. **[1]**

During the acceleration phase ($0 \leqslant t \leqslant 5$), the cyclist applies a driving force of magnitude directly proportional to t.

(d) Show that, for $0 \leqslant t \leqslant 5$, $\frac{\mathrm{d}v}{\mathrm{d}t} + 0.1v = \lambda t$, where λ is a positive constant. **[1]**

(e) **(i)** Show by integration that, for $0 \leqslant t \leqslant 5$, $v = 10\lambda(t - 10 + 10\mathrm{e}^{-0.1t})$. **[5]**

(ii) Hence find λ. **[2]**

(f) Find the total distance, to the nearest metre, travelled by the cyclist during the motion. **[6]**

END OF QUESTION PAPER

© OCR 2019

stemup.co.uk

BLANK PAGE

© OCR 2019

stemup.co.uk

OCR
Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© OCR 2019

stemup.co.uk

Surname	StemUp Model Solution
Answer booklet	Write the question number in the left-hand margin at the start of each answer.

Question number

1

$$\sum_{r=1}^{n}(2r^2 - 1) = 2 \times \frac{1}{6}n(n+1)(2n+1) - n$$

$$= \frac{1}{3}n(n+1)(2n+1) - n$$

$$= \frac{1}{3}n[(2n^2 + 3n + 1) - 3]$$

$$= \frac{1}{3}n(2n^2 + 3n - 2)$$

$$= \frac{1}{3}n(2n-1)(n+2)$$

2

Normals: $n_1 = (1, 2, c)$, $n_2 = (2, -c, 6)$

Perpendicular planes $\Rightarrow n_1 \cdot n_2 = 0$

$1(2) + 2(-c) + c(6) = 0$

$2 - 2c + 6c = 0 \Rightarrow 4c = -2$

$$c = -\frac{1}{2}$$

3 (a)

$$AB = \begin{pmatrix} 3 & 1 \\ 2 & 1 \end{pmatrix}\begin{pmatrix} k & 1 \\ 2 & 0 \end{pmatrix} = \begin{pmatrix} 3k+2 & 3 \\ 2k+2 & 2 \end{pmatrix}$$

$\det(AB) = 2(3k+2) - 3(2k+2) = 6k + 4 - 6k - 6 = -2$

$$(AB)^{-1} = -\frac{1}{2}\begin{pmatrix} 2 & -3 \\ -2k-2 & 3k+2 \end{pmatrix}$$

$$\det A = 3 - 2 = 1, \quad A^{-1} = \begin{pmatrix} 1 & -1 \\ -2 & 3 \end{pmatrix}$$

Question number

$\det B = 0 - 2 = -2, \quad B^{-1} = -\frac{1}{2}\begin{pmatrix} 0 & -1 \\ -2 & k \end{pmatrix}$

$B^{-1}A^{-1} = -\frac{1}{2}\begin{pmatrix} 0 & -1 \\ -2 & k \end{pmatrix}\begin{pmatrix} 1 & -1 \\ -2 & 3 \end{pmatrix}$

$= -\frac{1}{2}\begin{pmatrix} 2 & -3 \\ -2-2k & 2+3k \end{pmatrix}$

This is the same as $(AB)^{-1}$, so $(AB)^{-1} = B^{-1}A^{-1}$ as required.

3 (b) $BA = \begin{pmatrix} k & 1 \\ 2 & 0 \end{pmatrix}\begin{pmatrix} 3 & 1 \\ 2 & 1 \end{pmatrix} = \begin{pmatrix} 3k+2 & k+1 \\ 6 & 2 \end{pmatrix}$

$AB = BA$ requires $3 = k + 1$ and $2k + 2 = 6$, both give $k = 2$

So A and B commute only when $k = 2$; for any other k, $AB \neq BA$.

4 $V = \pi\int_0^{\frac{\pi}{2}} \sec^2\left(\frac{1}{2}x\right)\,dx$

$= \pi[2\tan\left(\frac{1}{2}x\right)]$ (from 0 to ½π)

$= \pi\left(2\tan\frac{\pi}{4} - 2\tan 0\right)$

$= \pi(2 \times 1 - 0) = 2\pi$

5 $\cos 2x = 1 - \frac{(2x)^2}{2!} + \frac{(2x)^4}{4!} - \frac{(2x)^6}{6!} + \ldots$

$= 1 - 2x^2 + \frac{2}{3}x^4 - \frac{4}{45}x^6 + \ldots$

$\cos 2x = 1 - 2\sin^2 x \Rightarrow \sin^2 x = \frac{1}{2}(1 - \cos 2x)$

Question number

$\sin^2 x \approx \frac{1}{2}(2x^2 - \frac{2}{3}x^4 + \frac{4}{45}x^6)$

$= x^2 - \frac{1}{3}x^4 + \frac{2}{45}x^6$

$a = 1, \quad b = -\frac{1}{3}, \quad c = \frac{2}{45}$

6 $\int_2^\infty \frac{1}{4 + x^2}\, dx = \lim_{t\to\infty} \int_2^t \frac{1}{4 + x^2}\, dx$

$= \lim_{t\to\infty} [\frac{1}{2} \arctan\frac{x}{2}]$ (from 2 to t)

$= \lim_{t\to\infty} (\frac{1}{2} \arctan\frac{t}{2} - \frac{1}{2} \arctan 1)$

As $t \to \infty$, $\arctan\frac{t}{2} \to \frac{\pi}{2}$

$= \frac{1}{2} \times \frac{\pi}{2} - \frac{1}{2} \times \frac{\pi}{4}$

$= \frac{\pi}{4} - \frac{\pi}{8} = \frac{\pi}{8}$

7 (a) $x = r\cos\theta, \quad y = r\sin\theta, \quad x^2 + y^2 = r^2$


$(r^2)^2 = 2c^2(r\cos\theta)(r\sin\theta)$

$r^4 = c^2r^2(2\sin\theta\cos\theta) = c^2r^2\sin 2\theta$

$r^2 = c^2\sin 2\theta$ as required

Question number

7 (b)



Solid loop in the first quadrant: $r = c\sqrt{(\sin 2\theta)}$

Dashed loop in the third quadrant: $r = -c\sqrt{(\sin 2\theta)}$

7 (c)

$A = \frac{1}{2}\int_0^{\frac{\pi}{2}} r^2\, d\theta = \frac{1}{2}\int_0^{\frac{\pi}{2}} c^2 \sin 2\theta\, d\theta$

$= \frac{1}{2}c^2\left[-\frac{1}{2}\cos 2\theta\right]$ (from 0 to ½π)

$= \frac{1}{2}c^2\left(\frac{1}{2} + \frac{1}{2}\right)$

$= \frac{1}{2}c^2$

8 (a)

Product of roots: $\alpha \times (1/\alpha) \times \beta = -(-2)/1 = 2$

$\Rightarrow \beta = 2$

Sum of roots: $\alpha + \frac{1}{\alpha} + \beta = -(-1)/1 = 1$

$\alpha + \frac{1}{\alpha} + 2 = 1 \Rightarrow \alpha + \frac{1}{\alpha} = -1$

$\alpha^2 + \alpha + 1 = 0$

Question number

$\alpha = \frac{-1 \pm \sqrt{1-4}}{2} = \frac{-1 \pm \sqrt{3}i}{2}$

Roots are $2, \ -\frac{1}{2} + \frac{\sqrt{3}}{2}i, \ -\frac{1}{2} - \frac{\sqrt{3}}{2}i$

8 (b) k = sum of products in pairs $= \alpha \times \frac{1}{\alpha} + \alpha\beta + \frac{\beta}{\alpha}$

$= 1 + 2(\alpha + \frac{1}{\alpha})$

$= 1 + 2(-1) = -1$

$k = -1$

9 Let $u_n = 5^n + 2 \times 11^n$

When $n = 1$: $u_1 = 5 + 22 = 27 = 3 \times 9$, divisible by 3, so true for $n = 1$

Assume true for $n = k$: $5^k + 2 \times 11^k = 3m$ for some integer m

$u_{k+1} = 5^{k+1} + 2 \times 11^{k+1}$

$= 5 \times 5^k + 22 \times 11^k$

$= 5(5^k + 2 \times 11^k) + 12 \times 11^k$

$= 5(3m) + 12 \times 11^k$

$= 3(5m + 4 \times 11^k)$, which is divisible by 3

So if the result is true for $n = k$, it is true for $n = k + 1$.

Since it is true for $n = 1$, by induction it is true for all positive integers n.

10 (a) $(-1 + i)^2 = 1 - 2i + i^2 = -2i$

$(-1 + i)^3 = -2i(-1 + i) = 2i - 2i^2 = 2 + 2i$

So $z^3 = 2 + 2i$: $a = 2, \ b = 2$

Question number

10 (b) $|2 + 2i| = \sqrt{8} = 2\sqrt{2}$, $\arg(2 + 2i) = \frac{\pi}{4}$

$z^3 = 2\sqrt{2}\,e^{i(\pi/4 + 2k\pi)}$, $k = 0, 1, -1$

$|z| = (2\sqrt{2})^{1/3} = (2^{3/2})^{1/3} = 2^{1/2} = \sqrt{2}$

$\arg z = \frac{\pi}{12} + \frac{2k\pi}{3}$: $\frac{\pi}{12}$, $\frac{3\pi}{4}$, $-\frac{7\pi}{12}$

$z = \sqrt{2}\,e^{i\pi/12}$, $\sqrt{2}\,e^{3i\pi/4}$, $\sqrt{2}\,e^{-7i\pi/12}$

10 (c) Roots only come in conjugate pairs when all the coefficients of the polynomial are real. Here $z^3 - (2 + 2i) = 0$ has a non-real coefficient, so the result does not apply.

11 (a) M_1: rotation about the origin, anticlockwise, through angle θ

where $\cos\theta = \frac{3}{5}$, $\sin\theta = \frac{4}{5}$: $\theta = 53.1°$ (0.927 rad)

M_2: reflection in the x-axis ($y = 0$)

11 (b) $M_3 = M_1M_2 = \frac{1}{5}\begin{pmatrix}3 & -4\\ 4 & 3\end{pmatrix}\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix} = \frac{1}{5}\begin{pmatrix}3 & 4\\ 4 & -3\end{pmatrix}$

Invariant points: $\frac{1}{5}\begin{pmatrix}3 & 4\\ 4 & -3\end{pmatrix}\begin{pmatrix}x\\ y\end{pmatrix} = \begin{pmatrix}x\\ y\end{pmatrix}$

$\frac{3}{5}x + \frac{4}{5}y = x \Rightarrow 4y = 2x \Rightarrow y = \frac{1}{2}x$

$\frac{4}{5}x - \frac{3}{5}y = y \Rightarrow 4x = 8y \Rightarrow y = \frac{1}{2}x$

Mirror line of R: $y = \frac{1}{2}x$

Question number

11 (c) $M_4 = M_2M_1 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \times \frac{1}{5}\begin{pmatrix} 3 & -4 \\ 4 & 3 \end{pmatrix} = \frac{1}{5}\begin{pmatrix} 3 & -4 \\ -4 & -3 \end{pmatrix}$

$M_4 \neq M_3$, so it is a different reflection (a reflection is fixed by its mirror line).

Invariant points: $\frac{3}{5}x - \frac{4}{5}y = x \Rightarrow y = -\frac{1}{2}x$

Mirror line is $y = -\frac{1}{2}x$, not $y = \frac{1}{2}x$, so the claim is incorrect.

12 L_1: $r = \lambda(2, 3, 1)$, L_2: $r = \mu(1, 2, -4)$, L_3: $r = (1, 2, -4) + \nu(1, 1, 5)$

L_1 and L_2 both pass through O, so they meet at O(0, 0, 0)

L_1 and L_3: $2\lambda = 1 + \nu$, $3\lambda = 2 + \nu \Rightarrow \lambda = 1$, $\nu = 1$

check z: $\lambda = 1$ and $-4 + 5(1) = 1$ ✓ meet at A(2, 3, 1)

L_2 and L_3: $\mu = 1 + \nu$, $2\mu = 2 + \nu \Rightarrow \nu = 0$, $\mu = 1$

check z: $-4(1) = -4 + 0$ ✓ meet at B(1, 2, −4)

$\overrightarrow{OA} \times \overrightarrow{OB} = (2, 3, 1) \times (1, 2, -4)$

$= (3(-4) - 1(2),\ 1(1) - 2(-4),\ 2(2) - 3(1))$

$= (-14, 9, 1)$

$|\overrightarrow{OA} \times \overrightarrow{OB}| = \sqrt{196 + 81 + 1} = \sqrt{278}$

Area $= \frac{1}{2}\sqrt{278} = 8.34$ units² (3 s.f.)

13 (a) $y = \operatorname{arcosh} x = \ln(x + \sqrt{x^2 - 1})$

$\frac{dy}{dx} = \frac{1 + x(x^2 - 1)^{-1/2}}{x + \sqrt{x^2 - 1}}$

$= \frac{1 + \frac{x}{\sqrt{x^2 - 1}}}{x + \sqrt{x^2 - 1}}$

Question number

$= \dfrac{\sqrt{x^2-1}+x}{\sqrt{x^2-1}(x+\sqrt{x^2-1})}$

$= \dfrac{1}{\sqrt{x^2-1}}$ as required

13 (b) By parts: $u = \operatorname{arcosh} x$, $u' = \dfrac{1}{\sqrt{x^2-1}}$, $v' = 1$, $v = x$

$\int_1^2 \operatorname{arcosh} x \, dx = [x \operatorname{arcosh} x]$ (from 1 to 2) $- \int_1^2 \dfrac{x}{\sqrt{x^2-1}} \, dx$

$= [x \operatorname{arcosh} x - \sqrt{x^2-1}]$ (from 1 to 2)

$= (2 \operatorname{arcosh} 2 - \sqrt{3}) - (\operatorname{arcosh} 1 - 0)$

$\operatorname{arcosh} 2 = \ln(2+\sqrt{3})$, $\operatorname{arcosh} 1 = \ln 1 = 0$

$= 2\ln(2+\sqrt{3}) - \sqrt{3}$

13 (c) arcosh x is only defined for $x \geq 1$ (for $x < 1$, $\sqrt{(x^2-1)}$ is not real),

so the integrand does not exist on most of the interval [0, 1].

14(a)(i) Coefficient matrix M has rows (−1, a, 0), (2, 3, 1), (1, b, 1)

$\det M = -1(3-b) - a(2-1) + 0 = b - 3 - a$

No unique point $\Rightarrow \det M = 0$

$b - 3 - a = 0 \Rightarrow b = a + 3$

14(a)(ii) Second equation − third equation: $x + (3-b)y = -3 - c$

With $b = a + 3$: $x - ay = -3 - c$

First equation: $-x + ay = 2 \Rightarrow x - ay = -2$

For a sheaf the equations must be consistent:

$-3 - c = -2 \Rightarrow c = -1$

Question number

14 (b)

$-x + ay = 2 \quad (1)$

$2x + 3y + z = -3 \quad (2)$

$x + ay + z = 1 \quad (3)$

$(2) - (3)$: $x + (3 - a)y = -4 \quad (4)$

$(1) + (4)$: $3y = -2 \Rightarrow y = -\frac{2}{3}$

From (1): $x = ay - 2 = -\frac{2a}{3} - 2 = -\frac{2a + 6}{3}$

From (3): $z = 1 - x - ay = 1 + \frac{2a + 6}{3} + \frac{2a}{3} = \frac{4a + 9}{3}$

Point of intersection: $\left(-\frac{2a + 6}{3}, -\frac{2}{3}, \frac{4a + 9}{3}\right)$

15

$4x^2 - 4x + 2 = (2x - 1)^2 + 1$

$\int_{\frac{3}{4}}^{\frac{3}{2}} \frac{1}{\sqrt{(2x - 1)^2 + 1}}\, dx = \left[\frac{1}{2}\operatorname{arsinh}(2x - 1)\right]$ (from ¾ to 3/2)

$= \frac{1}{2}\left(\operatorname{arsinh} 2 - \operatorname{arsinh}\frac{1}{2}\right)$

$\operatorname{arsinh} x = \ln\left(x + \sqrt{x^2 + 1}\right)$

$\operatorname{arsinh} 2 = \ln(2 + \sqrt{5}), \quad \operatorname{arsinh}\frac{1}{2} = \ln\left(\frac{1}{2} + \frac{\sqrt{5}}{2}\right) = \ln\left(\frac{1 + \sqrt{5}}{2}\right)$

$= \frac{1}{2}\ln\left(\frac{2(2 + \sqrt{5})}{1 + \sqrt{5}}\right)$

$= \frac{1}{2}\ln\left(\frac{2(2 + \sqrt{5})(\sqrt{5} - 1)}{(\sqrt{5} + 1)(\sqrt{5} - 1)}\right)$

$= \frac{1}{2}\ln\left(\frac{2(2\sqrt{5} - 2 + 5 - \sqrt{5})}{4}\right)$

Question number

$= \frac{1}{2}\ln\left(\frac{3+\sqrt{5}}{2}\right)$ as required

16 (a) $(2 - e^{i\theta})(2 - e^{-i\theta}) = 4 - 2e^{-i\theta} - 2e^{i\theta} + e^{i\theta}e^{-i\theta}$

$= 4 - 2(e^{i\theta} + e^{-i\theta}) + 1$

$= 5 - 2(2\cos\theta)$

$= 5 - 4\cos\theta$ as required

16 (b) $C + iS = \frac{1}{2}e^{i\theta} + \frac{1}{4}e^{2i\theta} + \frac{1}{8}e^{3i\theta} + \ldots + \frac{1}{2^n}e^{ni\theta}$

Geometric series, first term $\frac{1}{2}e^{i\theta}$, common ratio $\frac{1}{2}e^{i\theta}$

$$C + iS = \frac{\frac{1}{2}e^{i\theta}\left(1 - \frac{1}{2^n}e^{ni\theta}\right)}{1 - \frac{1}{2}e^{i\theta}} = \frac{e^{i\theta}\left(1 - \frac{1}{2^n}e^{ni\theta}\right)}{2 - e^{i\theta}}$$

$$= \frac{e^{i\theta}\left(1 - \frac{1}{2^n}e^{ni\theta}\right)(2 - e^{-i\theta})}{(2 - e^{i\theta})(2 - e^{-i\theta})}$$

$$= \frac{2e^{i\theta} - 1 - \frac{2}{2^n}e^{(n+1)i\theta} + \frac{1}{2^n}e^{ni\theta}}{5 - 4\cos\theta}$$

$$= \frac{2^{n+1}e^{i\theta} - 2^n - 2e^{(n+1)i\theta} + e^{ni\theta}}{2^n(5 - 4\cos\theta)}$$

$$C = \mathrm{Re}(C + iS) = \frac{2^{n+1}\cos\theta - 2^n - 2\cos(n+1)\theta + \cos n\theta}{2^n(5 - 4\cos\theta)}$$

$$C = \frac{2^n(2\cos\theta - 1) - 2\cos(n+1)\theta + \cos n\theta}{2^n(5 - 4\cos\theta)}$$ as required

Question number

17 (a) Air resistance is likely to increase as the speed increases,

so a force proportional to v is more realistic than a constant one.

17 (b) Newton's second law: $m\frac{dv}{dt} = -2m - 0.1mv$

$\div m$: $\frac{dv}{dt} + 0.1v = -2$ as required

17(c)(i) Integrating factor $= e^{\int 0.1\,dt} = e^{0.1t}$

$\frac{d}{dt}(ve^{0.1t}) = -2e^{0.1t}$

$ve^{0.1t} = \int -2e^{0.1t}\,dt = -20e^{0.1t} + A$

At $t = 10$, $v = 0$: $0 = -20e + A \Rightarrow A = 20e$

$ve^{0.1t} = -20e^{0.1t} + 20e$

$v = 20(e^{1 - 0.1t} - 1)$

17(c)(ii) $t = 5$: $v = 20(e^{0.5} - 1) = 12.97\ m\ s^{-1}$ $(13.0\ m\ s^{-1}$ to 3 s.f.)

17 (d) Driving force $= Ft$, where F is a positive constant

$m\frac{dv}{dt} = Ft - 0.1mv$

$\frac{dv}{dt} + 0.1v = \lambda t$, where $\lambda = \frac{F}{m} > 0$ as required

17(e)(i) Integrating factor $e^{0.1t}$: $\frac{d}{dt}(ve^{0.1t}) = \lambda te^{0.1t}$

$ve^{0.1t} = \int \lambda te^{0.1t}\,dt$

Question number

By parts: $= 10\lambda te^{0.1t} - \int 10\lambda e^{0.1t}\,dt$

$ve^{0.1t} = 10\lambda te^{0.1t} - 100\lambda e^{0.1t} + B$

At $t = 0$, $v = 0$: $0 = 0 - 100\lambda + B \Rightarrow B = 100\lambda$

$v = 10\lambda t - 100\lambda + 100\lambda e^{-0.1t}$

$v = 10\lambda(t - 10 + 10e^{-0.1t})$ as required

17(e)(ii) Velocities agree at $t = 5$:

$10\lambda(5 - 10 + 10e^{-0.5}) = 20(e^{0.5} - 1)$

$10.653...\lambda = 12.974...$

$\lambda = 1.22$ (3 s.f.)

17 (f) $s_1 = \int_0^5 10\lambda(t - 10 + 10e^{-0.1t})\,dt$

$= 10\lambda[\frac{1}{2}t^2 - 10t - 100e^{-0.1t}]$ (from 0 to 5)

$= 10\lambda(12.5 - 50 - 100e^{-0.5} + 100)$

$= 10 \times 1.2179... \times 1.8469... = 22.49$ m

$s_2 = \int_5^{10} 20(e^{1 - 0.1t} - 1)\,dt$

$= 20[-10e^{1 - 0.1t} - t]$ (from 5 to 10)

$= 20[(-10 - 10) - (-10e^{0.5} - 5)]$

$= 20(10e^{0.5} - 15) = 29.74$ m

Total distance $= 22.49 + 29.74 = 52.2... \approx 52$ m