OCR
Oxford Cambridge and RSA

Wednesday 25 May 2022 – Afternoon

A Level Further Mathematics B (MEI)

Y420/01 Core Pure

Time allowed: 2 hours 40 minutes

94514909091

You must have:
- the Printed Answer Booklet
- the Formulae Booklet for Further Mathematics B (MEI)
- a scientific or graphical calculator

INSTRUCTIONS
- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided in the **Printed Answer Booklet**. If you need extra space use the lined pages at the end of the Printed Answer Booklet. The question numbers must be clearly shown.
- Fill in the boxes on the front of the Printed Answer Booklet.
- Answer **all** the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.
- Give your final answers to a degree of accuracy that is appropriate to the context.
- Do **not** send this Question Paper for marking. Keep it in the centre or recycle it.

INFORMATION
- The total mark for this paper is **144**.
- The marks for each question are shown in brackets **[ ]**.
- This document has **8** pages.

ADVICE
- Read each question carefully before you start your answer.

© OCR 2022 [Y/508/5592]
DC (ST/SG) 301212/4
OCR is an exempt Charity

stemup.co.uk

Answer **all** the questions.

Section A (37 marks)

1 **(a)** By considering $(r+1)^3 - r^3$, find $\sum_{r=1}^{n}(3r^2+3r+1)$. **[3]**

(b) Use this result to find $\sum_{r=1}^{n} r(r+1)$, expressing your answer in fully factorised form. **[4]**

2 **In this question you must show detailed reasoning.**

Find the exact value of $\int_3^{\infty} \frac{1}{x^2-4x+5}\,\mathrm{d}x$. **[5]**

3 **In this question you must show detailed reasoning.**

Solve the equation $3\cosh x = 2\sinh^2 x$, giving your solutions in exact logarithmic form. **[6]**

4 **(a)** A transformation with associated matrix $\begin{pmatrix} m & 2 & 1 \\ 0 & 1 & -2 \\ 2 & 0 & 3 \end{pmatrix}$, where m is a constant, maps the vertices of a cube to points that all lie in a plane.

Find m. **[3]**

(b) The transformations S and T of the plane have associated matrices **M** and **N** respectively, where $\mathbf{M} = \begin{pmatrix} k & 1 \\ -3 & 4 \end{pmatrix}$ and the determinant of **N** is $3k+1$. The transformation U is equivalent to the combined transformation consisting of S followed by T.

Given that U preserves orientation and has an area scale factor 2, find the possible values of k. **[4]**

5 **(a)** Sketch the polar curve $r = a(1-\cos\theta)$, $0 \leqslant \theta < 2\pi$, where a is a positive constant. **[2]**

(b) Determine the exact area of the region enclosed by the curve. **[5]**

6 Prove by mathematical induction that $\begin{pmatrix} 2 & 0 \\ -1 & 1 \end{pmatrix}^n = \begin{pmatrix} 2^n & 0 \\ 1-2^n & 1 \end{pmatrix}$ for all positive integers n. **[5]**

© OCR 2022

stemup.co.uk

Answer **all** the questions.

Section B (107 marks)

7 **In this question you must show detailed reasoning.**

Show that $\displaystyle\int_2^3 \frac{x+1}{(x-1)(x^2+1)}\,\mathrm{d}x = \tfrac{1}{2}\ln 2$. **[9]**

8 Two sets of complex numbers are given by $\left\{z : \arg(z-10) = \frac{3}{4}\pi\right\}$ and $\{z : |z-3-6\mathrm{i}| = k\}$, where k is a positive constant. In an Argand diagram, one of the points of intersection of the two loci representing these sets lies on the imaginary axis.

(a) Sketch the loci on an Argand diagram. **[4]**

(b) **In this question you must show detailed reasoning.**

Find the complex numbers represented by the points of intersection. **[7]**

9 The function $\mathrm{f}(x)$ is defined by $\mathrm{f}(x) = \ln(1+\sinh x)$.

(a) Given that k lies in the domain of this function, explain why k must be greater than $\ln(\sqrt{2}-1)$. **[2]**

(b) **(i)** Find $\mathrm{f}'(x)$. **[2]**

(ii) Show that $\mathrm{f}''(x) = \dfrac{a\sinh x + b}{(1+\sinh x)^2}$, where a and b are integers to be determined. **[3]**

(c) Hence find a quadratic approximation to $\mathrm{f}(x)$ for small values of x. **[3]**

(d) Find the percentage error in this approximation when $x = 0.1$. **[2]**

stemup.co.uk

10 The equation


$4x^4 + 16x^3 + ax^2 + bx + 6 = 0,$

where a and b are real, has roots α, $\frac{2}{\alpha}$, β and 3β.

(a) Given that $\beta < 0$, determine all 4 roots. **[6]**

(b) Determine the values of a and b. **[4]**

11 An Argand diagram with the point A representing a complex number z_1 is shown below.



The complex numbers z_2 and z_3 are $z_1 e^{\frac{2}{3}i\pi}$ and $z_1 e^{\frac{4}{3}i\pi}$ respectively.

(a) **(i)** On the copy of the Argand diagram in the Printed Answer Booklet, mark the points B and C representing the complex numbers z_2 and z_3. **[2]**

(ii) Show that $z_1 + z_2 + z_3 = 0$. **[2]**

(b) Given now that z_1, z_2 and z_3 are roots of the equation $z^3 = 8i$, find these three roots, giving your answers in the form $a + ib$, where a and b are real and exact. **[4]**

© OCR 2022

stemup.co.uk

12 Solve the differential equation $(4-x^2)\frac{dy}{dx} - xy = 1$, given that $y = 1$ when $x = 0$, giving your answer in the form $y = f(x)$. **[9]**

13 The points A and B have coordinates (4, 0, −1) and (10, 4, −3) respectively. The planes Π_1 and Π_2 have equations $x - 2y = 5$ and $2x + 3y - z = -4$ respectively.

(a) Find the acute angle between the line AB and the plane Π_1. **[4]**

(b) Show that the line AB meets Π_1 and Π_2 at the same point, whose coordinates should be specified. **[5]**

(c) (i) Find $(\mathbf{i} - 2\mathbf{j}) \times (2\mathbf{i} + 3\mathbf{j} - \mathbf{k})$. **[1]**

(ii) Hence find the acute angle between the planes Π_1 and Π_2. **[3]**

(iii) Find the shortest distance between the point A and the line of intersection of the planes Π_1 and Π_2. **[4]**

14 (a) Find $(3 - e^{2i\theta})(3 - e^{-2i\theta})$ in terms of $\cos 2\theta$. **[2]**

(b) Hence show that the sum of the infinite series


$$\sin\theta + \frac{1}{3}\sin 3\theta + \frac{1}{9}\sin 5\theta + \frac{1}{27}\sin 7\theta + \ldots$$

can be expressed as $\frac{6\sin\theta}{5 - 3\cos 2\theta}$. **[6]**

© OCR 2022

stemup.co.uk

15 In an oscillating system, a particle of mass m kg moves in a horizontal line. Its displacement from its equilibrium position O at time t seconds is x metres, its velocity is $v\,\mathrm{m\,s^{-1}}$, and it is acted on by a force $2mx$ newtons acting towards O as shown in the diagram.



Initially, the particle is projected away from O with speed $1\,\mathrm{m\,s^{-1}}$ from a point 2 m from O in the positive direction.

(a) (i) Show that the motion is modelled by the differential equation $\dfrac{\mathrm{d}^2x}{\mathrm{d}t^2} + 2x = 0$. **[1]**

(ii) State the type of motion. **[1]**

(iii) Write down the period of the motion. **[1]**

(iv) Find x in terms of t. **[4]**

(v) Find the amplitude of the motion. **[2]**

(b) The motion is now damped by a force $2mv$ newtons.

(i) Show that $\dfrac{\mathrm{d}^2x}{\mathrm{d}t^2} + 2\dfrac{\mathrm{d}x}{\mathrm{d}t} + 2x = 0$. **[1]**

(ii) State, giving a reason, whether the system is under-damped, critically damped or over-damped. **[1]**

(iii) Determine the general solution of this differential equation. **[3]**

(c) Finally, a variable force $2m\cos 2t$ newtons is added, so that the motion is now modelled by the differential equation

$$\frac{\mathrm{d}^2x}{\mathrm{d}t^2} + 2\frac{\mathrm{d}x}{\mathrm{d}t} + 2x = 2\cos 2t.$$

(i) Find x in terms of t. **[7]**

In the long term, the particle is seen to perform simple harmonic motion with a period of just over 3 seconds.

(ii) Verify that this behaviour is consistent with the answer to part **(c)(i)**. **[2]**

END OF QUESTION PAPER

© OCR 2022

stemup.co.uk

BLANK PAGE

© OCR 2022

stemup.co.uk

OCR
Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.

© OCR 2022

stemup.co.uk

Surname	StemUp Model Solution
Answer booklet	Write the question number in the left-hand margin at the start of each answer.

Question number

1 (a)

$(r+1)^3 - r^3 = r^3 + 3r^2 + 3r + 1 - r^3 = 3r^2 + 3r + 1$

$\sum_{r=1}^{n} (3r^2 + 3r + 1) = \sum_{r=1}^{n} [(r+1)^3 - r^3]$

$= (2^3 - 1^3) + (3^3 - 2^3) + (4^3 - 3^3) + \ldots + ((n+1)^3 - n^3)$

All terms cancel except the last and the first:

$= (n+1)^3 - 1$

1 (b)

$\sum_{r=1}^{n} (3r^2 + 3r + 1) = 3\sum_{r=1}^{n} r(r+1) + \sum_{r=1}^{n} 1 = 3\sum_{r=1}^{n} r(r+1) + n$

$\Rightarrow 3\sum_{r=1}^{n} r(r+1) + n = (n+1)^3 - 1$

$\sum_{r=1}^{n} r(r+1) = \frac{1}{3}[(n+1)^3 - 1 - n] = \frac{1}{3}[(n+1)^3 - (n+1)]$

$= \frac{1}{3}(n+1)[(n+1)^2 - 1]$

$= \frac{1}{3}(n+1)(n+1-1)(n+1+1)$

$= \frac{1}{3}n(n+1)(n+2)$

2

$x^2 - 4x + 5 = (x-2)^2 + 1$

$\int_3^t \frac{1}{(x-2)^2+1}\,dx = [\arctan(x-2)]_3^t$

$= \arctan(t-2) - \arctan(1) = \arctan(t-2) - \frac{\pi}{4}$

Question number

As $t \to \infty$, $t - 2 \to \infty$ so $\arctan(t-2) \to \frac{\pi}{2}$

$\int_3^\infty \frac{1}{x^2 - 4x + 5}\,dx = \lim_{t\to\infty}[\arctan(t-2) - \frac{\pi}{4}] = \frac{\pi}{2} - \frac{\pi}{4}$

$= \frac{\pi}{4}$

3

$3\cosh x = 2\sinh^2 x = 2(\cosh^2 x - 1)$

$2\cosh^2 x - 3\cosh x - 2 = 0$

$(2\cosh x + 1)(\cosh x - 2) = 0$

$\cosh x = -\frac{1}{2}$ (reject, since $\cosh x \geq 1$) or $\cosh x = 2$

$\cosh x = 2 \Rightarrow e^x + e^{-x} = 4 \Rightarrow e^{2x} - 4e^x + 1 = 0$

$e^x = \frac{4 \pm \sqrt{16-4}}{2} = 2 \pm \sqrt{3}$

$x = \ln(2+\sqrt{3})$ or $x = \ln(2-\sqrt{3}) = \ln\frac{1}{2+\sqrt{3}} = -\ln(2+\sqrt{3})$

$x = \pm\ln(2+\sqrt{3})$

4 (a)

The image of the cube has zero volume, so the determinant is 0.

$\det = m(1\times 3 - (-2)\times 0) - 2(0\times 3 - (-2)\times 2) + 1(0\times 0 - 1\times 2)$

$= 3m - 8 - 2 = 3m - 10$

$3m - 10 = 0 \Rightarrow m = \frac{10}{3}$

4 (b)

$\det M = 4k - (1)(-3) = 4k + 3$

U is S followed by T, so U has matrix NM

$\det(NM) = \det N \times \det M = (3k+1)(4k+3)$

Question number

U preserves orientation with area scale factor 2, so $\det(NM) = 2$


$(3k + 1)(4k + 3) = 2$

$12k^2 + 13k + 3 = 2 \Rightarrow 12k^2 + 13k + 1 = 0$

$(12k + 1)(k + 1) = 0$

$k = -1 \quad \text{or} \quad k = -\frac{1}{12}$

5 (a)



Cardioid with cusp at 0, symmetrical about the initial line,

furthest point $(2a, \pi)$, crossing $\theta = \frac{1}{2}\pi$ and $\theta = 3\pi/2$ at $r = a$.

5 (b)

$A = \frac{1}{2}\int_0^{2\pi} a^2(1 - \cos\theta)^2 \, d\theta = \frac{1}{2}a^2\int_0^{2\pi} (1 - 2\cos\theta + \cos^2\theta) \, d\theta$

$\cos^2\theta = \frac{1}{2}(1 + \cos 2\theta)$

$A = \frac{1}{2}a^2\int_0^{2\pi} (\frac{3}{2} - 2\cos\theta + \frac{1}{2}\cos 2\theta) \, d\theta$

$= \frac{1}{2}a^2[\frac{3}{2}\theta - 2\sin\theta + \frac{1}{4}\sin 2\theta]_0^{2\pi}$

$= \frac{1}{2}a^2[(3\pi - 0 + 0) - 0]$

Question number

$= \frac{3}{2}\pi a^2$

6 $n = 1$: $LHS = \begin{pmatrix} 2 & 0 \\ -1 & 1 \end{pmatrix}$, $RHS = \begin{pmatrix} 2^1 & 0 \\ 1 - 2^1 & 1 \end{pmatrix} = \begin{pmatrix} 2 & 0 \\ -1 & 1 \end{pmatrix}$

So the result is true for $n = 1$.

Assume true for $n = k$: $\begin{pmatrix} 2 & 0 \\ -1 & 1 \end{pmatrix}^k = \begin{pmatrix} 2^k & 0 \\ 1 - 2^k & 1 \end{pmatrix}$

Then $\begin{pmatrix} 2 & 0 \\ -1 & 1 \end{pmatrix}^{k+1} = \begin{pmatrix} 2^k & 0 \\ 1 - 2^k & 1 \end{pmatrix}\begin{pmatrix} 2 & 0 \\ -1 & 1 \end{pmatrix}$

$= \begin{pmatrix} 2 \times 2^k + 0 & 0 + 0 \\ 2(1 - 2^k) - 1 & 0 + 1 \end{pmatrix} = \begin{pmatrix} 2^{k+1} & 0 \\ 2 - 2^{k+1} - 1 & 1 \end{pmatrix}$

$= \begin{pmatrix} 2^{k+1} & 0 \\ 1 - 2^{k+1} & 1 \end{pmatrix}$, which is the result for $n = k + 1$.

So if the result is true for $n = k$, it is true for $n = k + 1$.

Since it is true for $n = 1$, by induction it is true for all positive integers n.

7 $\frac{x + 1}{(x - 1)(x^2 + 1)} = \frac{A}{x - 1} + \frac{Bx + C}{x^2 + 1}$

$x + 1 = A(x^2 + 1) + (Bx + C)(x - 1)$

$x = 1$: $2 = 2A \Rightarrow A = 1$

Coefficient of x^2: $0 = A + B \Rightarrow B = -1$

Constant term: $1 = A - C \Rightarrow C = 0$

$\int_2^3 \frac{x + 1}{(x - 1)(x^2 + 1)}\,dx = \int_2^3 \left(\frac{1}{x - 1} - \frac{x}{x^2 + 1}\right) dx$

$= \left[\ln(x - 1) - \frac{1}{2}\ln(x^2 + 1)\right]_2^3$


Question number

$= (\ln 2 - \frac{1}{2}\ln 10) - (\ln 1 - \frac{1}{2}\ln 5)$

$= \ln 2 - \frac{1}{2}\ln 10 + \frac{1}{2}\ln 5 = \ln 2 - \frac{1}{2}\ln\frac{10}{5}$

$= \ln 2 - \frac{1}{2}\ln 2 = \frac{1}{2}\ln 2$ as required

8 (a)



Half-line from 10 at angle ¾π to the positive real axis;

circle centre 3 + 6i passing through the point where it meets the Im axis.

8 (b)

Half-line: $y = -(x - 10)$, i.e. $x + y = 10$, with $x < 10$

It meets the imaginary axis when $x = 0$, so $y = 10$: the point $10i$

$10i$ lies on the circle, so $k = |10i - 3 - 6i| = |-3 + 4i|$

$k = \sqrt{(-3)^2 + 4^2} = 5$

Circle: $(x - 3)^2 + (y - 6)^2 = 25$

Substitute $y = 10 - x$: $(x - 3)^2 + (4 - x)^2 = 25$

$x^2 - 6x + 9 + 16 - 8x + x^2 = 25$

Question number

$2x^2 - 14x = 0 \Rightarrow 2x(x - 7) = 0$

$x = 0, y = 10$ or $x = 7, y = 3$ ($x < 10$ so on the half-line)

The points of intersection represent $10i$ and $7 + 3i$

9 (a) ln is only defined for positive values, so $1 + \sinh k > 0$

$\sinh k > -1$, and sinh is an increasing function, so

$k > \text{arsinh}(-1) = \ln(-1 + \sqrt{(-1)^2 + 1}) = \ln(\sqrt{2} - 1)$

9 (b)(i) $f'(x) = \dfrac{\cosh x}{1 + \sinh x}$

9 (b)(ii) $f''(x) = \dfrac{(1 + \sinh x)\sinh x - \cosh x \times \cosh x}{(1 + \sinh x)^2}$

$= \dfrac{\sinh x + \sinh^2 x - \cosh^2 x}{(1 + \sinh x)^2}$

$\cosh^2 x - \sinh^2 x = 1 \Rightarrow f'(x) = \dfrac{\sinh x - 1}{(1 + \sinh x)^2}$

$a = 1, \ b = -1$

9 (c) $f(0) = \ln(1 + 0) = 0$

$f'(0) = \dfrac{1}{1 + 0} = 1, \quad f''(0) = \dfrac{0 - 1}{(1 + 0)^2} = -1$

$f(x) \approx f(0) + f'(0)x + \dfrac{f''(0)}{2!}x^2 = 0 + x - \dfrac{1}{2}x^2$

$f(x) \approx x - \dfrac{1}{2}x^2$

9 (d) Exact: $f(0.1) = \ln(1 + \sinh 0.1) = 0.095462 \ldots$

Question number

Approximation: $0.1 - \frac{1}{2}(0.1)^2 = 0.095$

Percentage error $= \frac{0.095 - 0.095462}{0.095462} \times 100 = -0.48\%$

(the approximation is 0.48% too small)

10 (a)

Product of roots: $\alpha \times \frac{2}{\alpha} \times \beta \times 3\beta = \frac{6}{4}$

$6\beta^2 = \frac{3}{2} \Rightarrow \beta^2 = \frac{1}{4} \Rightarrow \beta = -\frac{1}{2}$ (since $\beta < 0$)

Sum of roots: $\alpha + \frac{2}{\alpha} + \beta + 3\beta = -\frac{16}{4} = -4$

$\alpha + \frac{2}{\alpha} - 2 = -4 \Rightarrow \alpha + \frac{2}{\alpha} = -2$

$\alpha^2 + 2\alpha + 2 = 0$

$\alpha = \frac{-2 \pm \sqrt{4 - 8}}{2} = -1 \pm i$

If $\alpha = -1 + i$ then $\frac{2}{\alpha} = \frac{2(-1 - i)}{2} = -1 - i$ (and vice versa)

Roots: $-1 + i$, $-1 - i$, $-\frac{1}{2}$, $-\frac{3}{2}$

10 (b)

$(x - (-1 + i))(x - (-1 - i)) = (x + 1)^2 + 1 = x^2 + 2x + 2$

$(2x + 1)(2x + 3) = 4x^2 + 8x + 3$

$4x^4 + 16x^3 + ax^2 + bx + 6 = (x^2 + 2x + 2)(4x^2 + 8x + 3)$


$= 4x^4 + 8x^3 + 3x^2 + 8x^3 + 16x^2 + 6x + 8x^2 + 16x + 6$

$= 4x^4 + 16x^3 + 27x^2 + 22x + 6$

$a = 27, \quad b = 22$

Question number

11 (a)(i)



B and C are A rotated by $\frac{2}{3}\pi$ and $4\pi/3$ about O (same modulus as z_1),

so A, B, C form an equilateral triangle on a circle centre O.

11 (a)(ii) $z_1 + z_2 + z_3 = z_1(1 + e^{2\pi i/3} + e^{4\pi i/3})$

$e^{2\pi i/3} = -\frac{1}{2} + \frac{\sqrt{3}}{2}i, \quad e^{4\pi i/3} = -\frac{1}{2} - \frac{\sqrt{3}}{2}i$

$1 + e^{2\pi i/3} + e^{4\pi i/3} = 1 - \frac{1}{2} - \frac{1}{2} + (\frac{\sqrt{3}}{2} - \frac{\sqrt{3}}{2})i = 0$

So $z_1 + z_2 + z_3 = z_1 \times 0 = 0$ as required

11 (b) $z^3 = 8i = 8e^{\pi i/2}$ (modulus 8, argument $\frac{1}{2}\pi$)

$|z| = \sqrt[3]{8} = 2$

$\arg z = \frac{1}{3}(\frac{\pi}{2} + 2k\pi) = \frac{\pi}{6}, \ \frac{5\pi}{6}, \ \frac{3\pi}{2}$

$z = 2e^{\pi i/6}, \ 2e^{5\pi i/6}, \ 2e^{3\pi i/2}$

$z = \sqrt{3} + i, \quad -\sqrt{3} + i, \quad -2i$

12 $\frac{dy}{dx} - \frac{x}{4 - x^2}y = \frac{1}{4 - x^2}$

Question number

Integrating factor $= e^{\int -x/(4-x^2)\,dx} = e^{\frac{1}{2}\ln(4-x^2)} = \sqrt{4-x^2}$

$$\sqrt{4-x^2}\frac{dy}{dx} - \frac{x}{\sqrt{4-x^2}}y = \frac{1}{\sqrt{4-x^2}}$$

$$\frac{d}{dx}\left(y\sqrt{4-x^2}\right) = \frac{1}{\sqrt{4-x^2}}$$

$$y\sqrt{4-x^2} = \int \frac{1}{\sqrt{4-x^2}}\,dx = \arcsin\frac{x}{2} + c$$

$x = 0, y = 1$: $1 \times 2 = \arcsin 0 + c \Rightarrow c = 2$

$$y = \frac{\arcsin\left(\frac{x}{2}\right) + 2}{\sqrt{4-x^2}}$$

13 (a)

vector $AB = \begin{pmatrix} 10-4 \\ 4-0 \\ -3-(-1) \end{pmatrix} = \begin{pmatrix} 6 \\ 4 \\ -2 \end{pmatrix}$, direction $\begin{pmatrix} 3 \\ 2 \\ -1 \end{pmatrix}$

Normal to Π_1: $n_1 = \begin{pmatrix} 1 \\ -2 \\ 0 \end{pmatrix}$

$$\begin{pmatrix} 3 \\ 2 \\ -1 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ -2 \\ 0 \end{pmatrix} = 3 - 4 + 0 = -1$$

$$\sin\varphi = \frac{|-1|}{\sqrt{14}\sqrt{5}} = \frac{1}{\sqrt{70}} \quad (\varphi = \text{angle between } AB \text{ and } \Pi_1)$$

$\varphi = 6.86°$ (3 s.f.)

13 (b)

Line AB: $r = \begin{pmatrix} 4 \\ 0 \\ -1 \end{pmatrix} + \lambda\begin{pmatrix} 3 \\ 2 \\ -1 \end{pmatrix}$, so $x = 4 + 3\lambda$, $y = 2\lambda$, $z = -1 - \lambda$

Question number

Π_1: $(4 + 3\lambda) - 2(2\lambda) = 5 \Rightarrow 4 - \lambda = 5 \Rightarrow \lambda = -1$

Π_2: $2(4 + 3\lambda) + 3(2\lambda) - (-1 - \lambda) = -4$

$8 + 6\lambda + 6\lambda + 1 + \lambda = -4 \Rightarrow 13\lambda = -13 \Rightarrow \lambda = -1$

Same value of λ, so AB meets Π_1 and Π_2 at the same point:

$\lambda = -1$: $(4 - 3, -2, -1 + 1) = (1, -2, 0)$

13 (c)(i) $(i - 2j) \times (2i + 3j - k) = \begin{pmatrix} (-2)(-1) - 0 \times 3 \\ 0 \times 2 - 1 \times (-1) \\ 1 \times 3 - (-2) \times 2 \end{pmatrix} = 2i + j + 7k$

13 (c)(ii) $|n_1 \times n_2| = \sqrt{2^2 + 1^2 + 7^2} = \sqrt{54}$

$|n_1 \times n_2| = |n_1||n_2| \sin\theta \Rightarrow \sqrt{54} = \sqrt{5}\sqrt{14} \sin\theta$

$\sin\theta = \frac{\sqrt{54}}{\sqrt{70}} \Rightarrow \theta = 61.4°$ (3 s.f.)

13 (c)(iii) Line of intersection passes through $C(1, -2, 0)$ (on both planes from (b))

and has direction $d = 2i + j + 7k$ (perpendicular to both normals).

vector $CA = \begin{pmatrix} 4 - 1 \\ 0 + 2 \\ -1 - 0 \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ -1 \end{pmatrix}$

$CA \times d = \begin{pmatrix} 3 \\ 2 \\ -1 \end{pmatrix} \times \begin{pmatrix} 2 \\ 1 \\ 7 \end{pmatrix} = \begin{pmatrix} 2 \times 7 - (-1) \times 1 \\ (-1) \times 2 - 3 \times 7 \\ 3 \times 1 - 2 \times 2 \end{pmatrix} = \begin{pmatrix} 15 \\ -23 \\ -1 \end{pmatrix}$

distance $= \frac{|CA \times d|}{|d|} = \frac{\sqrt{15^2 + (-23)^2 + (-1)^2}}{\sqrt{2^2 + 1^2 + 7^2}} = \frac{\sqrt{755}}{\sqrt{54}}$

$= 3.74$ (3 s.f.)

14 (a) $(3 - e^{2i\theta})(3 - e^{-2i\theta}) = 9 - 3e^{-2i\theta} - 3e^{2i\theta} + e^0$

Question number

$= 10 - 3(e^{2i\theta} + e^{-2i\theta}) = 10 - 3(2\cos 2\theta)$

$= 10 - 6\cos 2\theta$

14 (b) Let $S = \sin\theta + \frac{1}{3}\sin 3\theta + \frac{1}{9}\sin 5\theta + \frac{1}{27}\sin 7\theta + \ldots$

and $C = \cos\theta + \frac{1}{3}\cos 3\theta + \frac{1}{9}\cos 5\theta + \frac{1}{27}\cos 7\theta + \ldots$

$C + iS = e^{i\theta} + \frac{1}{3}e^{3i\theta} + \frac{1}{9}e^{5i\theta} + \frac{1}{27}e^{7i\theta} + \ldots$

Geometric series, first term $e^{i\theta}$, ratio $\frac{1}{3}e^{2i\theta}$, $|\text{ratio}| = \frac{1}{3} < 1$ so it converges

$$C + iS = \frac{e^{i\theta}}{1 - \frac{1}{3}e^{2i\theta}} = \frac{3e^{i\theta}}{3 - e^{2i\theta}}$$

$$= \frac{3e^{i\theta}(3 - e^{-2i\theta})}{(3 - e^{2i\theta})(3 - e^{-2i\theta})} = \frac{9e^{i\theta} - 3e^{-i\theta}}{10 - 6\cos 2\theta}$$

$$= \frac{9(\cos\theta + i\sin\theta) - 3(\cos\theta - i\sin\theta)}{10 - 6\cos 2\theta}$$

$$S = \text{Im}(C + iS) = \frac{9\sin\theta + 3\sin\theta}{10 - 6\cos 2\theta} = \frac{12\sin\theta}{10 - 6\cos 2\theta}$$

$$= \frac{6\sin\theta}{5 - 3\cos 2\theta} \quad \text{as required}$$

15 (a)(i) Newton II (force towards O): $m\frac{d^2x}{dt^2} = -2mx \Rightarrow \frac{d^2x}{dt^2} + 2x = 0$

15 (a)(ii) Simple harmonic motion

Question number

15 (a)(iii) $\omega = \sqrt{2}$, period $= \frac{2\pi}{\sqrt{2}} = \sqrt{2}\,\pi \approx 4.44$ s

15 (a)(iv) Auxiliary equation $\lambda^2 + 2 = 0 \Rightarrow \lambda = \pm\sqrt{2}\,i$

$x = A\cos\sqrt{2}t + B\sin\sqrt{2}t$

$t = 0,\ x = 2 \Rightarrow A = 2$

$\frac{dx}{dt} = -\sqrt{2}A\sin\sqrt{2}t + \sqrt{2}B\cos\sqrt{2}t$

$t = 0,\ \frac{dx}{dt} = 1 \Rightarrow \sqrt{2}B = 1 \Rightarrow B = \frac{\sqrt{2}}{2}$

$x = 2\cos\sqrt{2}t + \frac{\sqrt{2}}{2}\sin\sqrt{2}t$

15 (a)(v) Amplitude $= \sqrt{2^2 + (\frac{\sqrt{2}}{2})^2} = \sqrt{\frac{9}{2}} = \frac{3\sqrt{2}}{2} \approx 2.12$ m

15 (b)(i) $m\frac{d^2x}{dt^2} = -2mx - 2mv = -2mx - 2m\frac{dx}{dt}$

$\Rightarrow \frac{d^2x}{dt^2} + 2\frac{dx}{dt} + 2x = 0$ as required

15 (b)(ii) Auxiliary equation $\lambda^2 + 2\lambda + 2 = 0$ has discriminant $2^2 - 4 \times 1 \times 2 = -4 < 0$ (complex roots), so the system is under-damped.

15 (b)(iii) $\lambda^2 + 2\lambda + 2 = 0 \Rightarrow \lambda = \frac{-2 \pm \sqrt{-4}}{2} = -1 \pm i$

$x = e^{-t}(A\cos t + B\sin t)$

Question number

15 (c)(i) Particular integral: $x = C\cos 2t + D\sin 2t$

$\frac{dx}{dt} = -2C\sin 2t + 2D\cos 2t, \quad \frac{d^2x}{dt^2} = -4C\cos 2t - 4D\sin 2t$

Substitute: $(-4C + 4D + 2C)\cos 2t + (-4D - 4C + 2D)\sin 2t = 2\cos 2t$

cos 2t: $-2C + 4D = 2$ sin 2t: $-4C - 2D = 0 \Rightarrow D = -2C$

$-2C - 8C = 2 \Rightarrow C = -0.2, \quad D = 0.4$

$x = e^{-t}(A\cos t + B\sin t) - 0.2\cos 2t + 0.4\sin 2t$

$t = 0, x = 2$: $A - 0.2 = 2 \Rightarrow A = 2.2$

$\frac{dx}{dt} = -e^{-t}(A\cos t + B\sin t) + e^{-t}(-A\sin t + B\cos t)$

$+ 0.4\sin 2t + 0.8\cos 2t$

$t = 0, \frac{dx}{dt} = 1$: $1 = -A + B + 0.8 \Rightarrow B = 1 + 2.2 - 0.8 = 2.4$

$x = e^{-t}(2.2\cos t + 2.4\sin t) - 0.2\cos 2t + 0.4\sin 2t$

15 (c)(ii) As $t \to \infty$, $e^{-t} \to 0$, so $x \to -0.2\cos 2t + 0.4\sin 2t$

This is simple harmonic motion with $\omega = 2$,

period $= \frac{2\pi}{2} = \pi \approx 3.14$ s, which is just over 3 seconds, so it is consistent.