OCR
Oxford Cambridge and RSA

Wednesday 8 June 2022 – Afternoon

A Level Further Mathematics B (MEI)

Y433/01 Modelling with Algorithms

Time allowed: 1 hour 15 minutes

9453418660

You must have:
- the Printed Answer Booklet
- the Formulae Booklet for Further Mathematics B (MEI)
- a scientific or graphical calculator

INSTRUCTIONS
- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided in the **Printed Answer Booklet**. If you need extra space use the lined pages at the end of the Printed Answer Booklet. The question numbers must be clearly shown.
- Fill in the boxes on the front of the Printed Answer Booklet.
- Answer **all** the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.
- Give your final answers to a degree of accuracy that is appropriate to the context.
- Do **not** send this Question Paper for marking. Keep it in the centre or recycle it.

INFORMATION
- The total mark for this paper is **60**.
- The marks for each question are shown in brackets **[ ]**.
- This document has **12** pages.

ADVICE
- Read each question carefully before you start your answer.

© OCR 2022 [T/508/5597]
DC (RW/SW) 312288/4

OCR is an exempt Charity

stemup.co.uk

Answer **all** the questions.

1




Fig. 1.1

The diagram in **Fig. 1.1** represents a system of pipes through which a fluid can flow from three sources to three sinks. The weights on the arcs show the capacities of the pipes in litres per minute.

(a) Add a supersource S and a supersink T to the network in the Printed Answer Booklet. Give appropriate weightings and directions to the connecting arcs. **[2]**

(b) The cut α partitions the vertices into sets {S, A, B, C, D},{E, F, G, H, I, J, K, T}. Calculate the capacity of cut α. **[1]**

(c) Explain why a flow of 34 litres per minute along AE cannot be achieved. **[1]**

(d) An LP formulation is set up to find the maximum flow through the network. Write down a suitable objective function for the LP formulation. **[1]**

© OCR 2022

stemup.co.uk





Fig. 1.2

The diagram in **Fig. 1.2** shows a feasible flow through the network.

(e) Show that this is a maximum flow. **[2]**

stemup.co.uk

2



The flow chart shows an algorithm that gives a numerical approximation for the real root of the cubic equation $x^3 - 11x - 21 = 0$.

(a) Work through the algorithm using the input $A = 5$. Record the values of A and B correct to **7** decimal places every time they change. Give the final output correct to **3** decimal places. **[3]**

(b) Show that the algorithm does give the real root of $x^3 - 11x - 21 = 0$ correct to **3** decimal places. **[2]**

(c) Explain how to adapt the algorithm so that a numerical approximation for the real root of the cubic equation $x^3 - 12x - 23 = 0$ can be found. **[1]**

© OCR 2022

stemup.co.uk

3




Fig. 3.1

Activity	Independent Float
A	0
B	7
C	0
D	0
E	0
F	2
G	2
H	0
I	0
J	1
K	0

Fig. 3.2

Fig. 3.1 shows an activity network for a project. Each activity is represented by an arc. The duration of each activity is measured in hours. The early event times and the late event times are shown at each vertex.

Fig. 3.2 shows the independent float for each activity.

(a) State which activities are critical. **[1]**

(b) Complete the table in the Printed Answer Booklet showing the duration of each activity. **[3]**

Each activity requires one worker. Once a worker has started an activity they continue with that activity until it is complete.

(c) Draw a schedule to show how three workers can complete the project in the minimum completion time. Each box in the Printed Answer Booklet represents 1 hour. For each worker, write the letter of the activity they are doing in each box, or leave the box blank if the worker is not required for that 1 hour. **[2]**

© OCR 2022

stemup.co.uk

4 A list of n numbers is to be sorted into descending order using the quick sort algorithm.

The number of comparisons made in each pass of the algorithm is used as a measure of the complexity of the quick sort algorithm.

(a) Show that, in the worst case, the quick sort algorithm has complexity $O(n^2)$. **[2]**

(b) A computer takes 2.3×10^{-7} seconds to sort the list of 100 ascending integers 1, 2, 3…, 99, 100 into descending order using the quick sort algorithm.

Calculate approximately how long it will take the computer to sort the list of 500 000 ascending integers 1, 2, 3,…, 499 999, 500 000 into descending order using the quick sort algorithm. You should assume that the computer uses the first value as the pivot for each sublist. **[2]**

A student attempts to use the quick sort algorithm to sort a list of seven random integers into descending order. After two passes through the list the student produces the following list.

6 8 10 5 7 11 4

(c) Explain how you know that the student has made at least one error in these two passes. **[1]**

© OCR 2022

stemup.co.uk

5 Consider the following LP problem.

Maximise $P = 2x + 3y - z$

Subject to

$3x + y - 4z \leqslant 70$
$5x + 4y \leqslant 60$
$x \geqslant 4,\ y \geqslant 2,\ z \leqslant -2$

(a) Explain why the simplex algorithm cannot be used to solve this LP problem. **[1]**

(b) Use the substitutions $x = X + 4,\ y = Y + 2,\ z = -Z - 2$ and $P = Q + 16$ to reformulate this LP problem into standard form. **[2]**

(c) Represent the reformulated problem as an initial simplex tableau. **[2]**

After a first iteration of the simplex algorithm the tableau below is produced.

Q	X	Y	Z	s_1	s_2	RHS
1	$\frac{7}{4}$	0	-1	0	$\frac{3}{4}$	24
0	$\frac{7}{4}$	0	4	1	$-\frac{1}{4}$	40
0	$\frac{5}{4}$	1	0	0	$\frac{1}{4}$	8

(d) Perform a second iteration of the simplex algorithm. **[3]**

(e) Find the maximum value of P, and the corresponding values of x, y and z. **[2]**

stemup.co.uk

6 **Fig. 6.1** shows a weighted graph. The weights represent arc lengths.




Fig. 6.1

(a) Apply Dijkstra's algorithm to the copy of the network in the Printed Answer Booklet to find the shortest path from A to G. **[5]**

An LP formulation is set up to find the longest route between A and G in **Fig. 6.1** subject to the conditions that no arc can be repeated but nodes B, C, D, E and F can be passed through more than once.

Maximise $15AB + 32AC + 29AD + 37AE + 16BC + 11BD + 20BE$
$+ 16CB + 12CE + 10CF + 11DB + 8DE + 17DF + 12EC$
$+ 16EG + 10FC + 17FD + 6FE + 7FG$

Subject to

$AB + AC + AD + AE = 1$
$EG + FG = 1$
$AB + CB + DB - BC - BD - BE = 0$
$AC + BC + EC + FC - CB - CE - CF = 0$
$AD + BD + FD - DB - DE - DF = 0$
$AE + BE + CE + DE + FE - EC - EG = 0$
$CF + DF - FC - FD - FE - FG = 0$
$BD + DB \leqslant 1$
$BC + CB \leqslant 1$
$CE + EC \leqslant 1$
$CF + FC \leqslant 1$
$DF + FD \leqslant 1$
$BE \leqslant 1$
$DE \leqslant 1$
$FE \leqslant 1$

© OCR 2022

stemup.co.uk

(b) Explain the purpose of the following lines in the LP formulation

(i) $AB + AC + AD + AE = 1$ **[1]**

(ii) $CF + DF - FC - FD - FE - FG = 0$ **[1]**

(iii) $CE + EC \leqslant 1$ **[1]**

The LP is run in a solver and the following output is shown in the table in **Fig. 6.2**.

VARIABLE	VALUE
AB	0.000000
AC	1.000000
AD	0.000000
AE	0.000000
BC	0.000000
BD	0.000000
BE	1.000000
CB	1.000000
CE	0.000000
CF	1.000000
DB	0.000000
DE	1.000000
DF	0.000000
EC	1.000000
EG	1.000000
FC	0.000000
FD	1.000000
FE	0.000000
FG	0.000000

Fig. 6.2

(c) **(i)** Find the length of the longest route from A to G given by the output in **Fig. 6.2**. **[1]**

(ii) State all possible corresponding longest routes from A to G. **[2]**

© OCR 2022

stemup.co.uk

7 Kai makes three sizes of model car, small, medium and large, to sell each year at a fair.

Kai has enough material to make either 250 small cars or 175 medium cars or 150 large cars. Alternatively, he could make a combination of the three sizes.

Each small car requires 40 minutes to make, each medium car requires 70 minutes to make, and each large car requires 140 minutes to make.

The total time that Kai spends making all the cars must not exceed 220 hours.

From experience, small cars sell particularly well so Kai decides to make at least three small cars for every one large car.

Furthermore, Kai decides to make exactly 210 cars.

Kai plans on selling each small car for £5, each medium car for £8 and each large car for £12. Kai wants to maximise the total income from the sale of the cars at the fair (that is, money made from selling cars, ignoring the cost of materials).

Let x, y and z represent the number of small, medium and large cars respectively that Kai makes.

(a) Explain why the maximum total income is achieved when $7x + 4y$ is minimised. **[2]**

(b) Determine the constraints of this ILP problem by listing them as simplified inequalities with integer coefficients in x and y only. **[6]**

(c) By representing the feasible region for x and y graphically, determine how many of each size of car Kai should make to maximise the total income from the fair. You must show all your working. **[6]**

(d) Using the solution found in part **(c)**, find the maximum possible total income from the sales. **[1]**

END OF QUESTION PAPER

© OCR 2022

stemup.co.uk

BLANK PAGE

© OCR 2022

stemup.co.uk

OCR
Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.


© OCR 2022

stemup.co.uk

Surname	StemUp Model Solution
Answer booklet	Write the question number in the left-hand margin at the start of each answer.

Question number

1 (a)



SA = 15 + 34 = 49, SB = 15, SC = 10 + 39 = 49

HT = 25, IT = 3 + 25 = 28, KT = 17 + 29 = 46

1 (b)

Arcs from {S, A, B, C, D} to the other set: AF, AE, BE, CG, DF

(ED goes from E to D, backwards across the cut, so it is not counted)

Capacity of cut α = 15 + 34 + 15 + 39 + 24 = 127 litres per minute

1 (c)

The only arcs out of E are ED (8) and EG (24).

So at most 8 + 24 = 32 litres per minute can leave E.

Flow in = flow out at E, so at most 32 < 34 can flow along AE.

1 (d)

Maximise AF + AE + BE + CD + CG

(the total flow out of the three sources)

Question number

1 (e)

Flow in Fig. 1.2 = flow into the sinks

$= GH + FI + JI + FK + GK = 25 + 3 + 20 + 10 + 29 = 87$

Consider the cut $\{S, A, B, C, E, G\} \mid \{D, F, H, I, J, K, T\}$

Arcs crossing forwards: AF, CD, ED, GH, GK

(FG and JG go backwards across the cut)

Capacity $= 15 + 10 + 8 + 25 + 29 = 87$

By the maximum flow, minimum cut theorem, flow ≤ 87 for any flow,

and the flow shown is 87, so 87 litres per minute is a maximum flow.

2 (a)

A	B	$\lvert A - B \rvert < 0.001$?
S	4.2358236	No
4.2358236	4.0735168	No
4.0735168	4.0373314	No
4.0373314	4.0291751	No
4.0291751	4.0273321	No
4.0273321	4.0269154	Yes

Output B = 4.027 (3 d.p.)

2 (b)

Let $f(x) = x^3 - 11x - 21$

$f(4.0265) = 4.0265^3 - 11(4.0265) - 21 = -0.0111 < 0$

$f(4.0275) = 4.0275^3 - 11(4.0275) - 21 = 0.0266 > 0$

Change of sign and f is continuous, so there is a root in

$4.0265 < x < 4.0275$, so the root is 4.027 correct to 3 d.p.

This agrees with the output, so the algorithm gives the real root.

2 (c)

Change the box "Let $B = \sqrt[3]{(11A + 21)}$"

to "Let $B = \sqrt[3]{(12A + 23)}$" (everything else stays the same).


Question number

3 (a) Critical activities: C, E, H and K

3 (b)

Activity	Duration (hours)
A	5
B	7
C	10
D	2
E	4
F	5
G	4
H	3
I	3
J	5
K	3

3 (c)



C 0–10, E 10–14, H 14–17, K 17–20 (critical path, worker 1)

A 0–5, D 5–7, I 7–10, F 10–15, G 15–19 (worker 2)

B 0–7, J 7–12 (worker 3). Project completed in 20 hours.

4 (a) Worst case: each pivot ends up at one end of its sublist, so each pass only fixes one value. With n values the passes need at most

Question number

$(n-1), (n-2), \ldots, 2, 1$ comparisons.

Total $= (n-1) + (n-2) + \ldots + 1 = \sum_{r=1}^{n-1} r = \frac{1}{2}n(n-1)$

$= \frac{1}{2}n^2 - \frac{1}{2}n$, which is quadratic in n, so the complexity is $O(n^2)$

4 (b) Ascending list into descending order with the first value as pivot

is the worst case, so the time is proportional to n^2.

Time $\approx 2.3 \times 10^{-7} \times (500000 \div 100)^2$

$= 2.3 \times 10^{-7} \times 2.5 \times 10^{7} = 5.75$ seconds (about 5.8 s)

4 (c) Correct final order: 11 10 8 7 6 5 4

After two passes at least two pivots must be in their final positions.

In 6 8 10 5 7 11 4 only the 4 is in its correct position

(e.g. 11 should be at the far left), so there must be an error.

5 (a) The origin is not in the feasible region ($x \geq 4$, $y \geq 2$, $z \leq -2$),

so there is no initial basic feasible solution at $x = y = z = 0$.

5 (b) $Q + 16 = 2(X + 4) + 3(Y + 2) - (-2 - 2) = 2X + 3Y + 2 + 16$

$3(X + 4) + (Y + 2) - 4(-2 - 2) \leq 70 \Rightarrow 3X + Y + 42 + 22 \leq 70$

$5(X + 4) + 4(Y + 2) \leq 60 \Rightarrow 5X + 4Y + 28 \leq 60$

Maximise $Q = 2X + 3Y + 2$

subject to $3X + Y + 42 \leq 48$

$5X + 4Y \leq 32$

$X \geq 0, \quad Y \geq 0, \quad 2 \geq 0$

5 (c)

Q	X	Y	Z	s_1	s_2	RHS

Question number

1	−2	−3	−1	0	0	0
0	3	1	4	1	0	48
0	5	4	0	0	1	32

5 (d)

Only negative entry in the objective row is −1 in the Z column.

Ratio test: row 2: $40 \div 4 = 10$ (row 3 has 0 in Z column, ignore)

Pivot on 4 in row 2.

R2 new = R2 ÷ 4, R1 new = R1 + R2 new, R3 new = R3

Q	X	Y	Z	s_1	s_2	RHS
1	$\frac{35}{16}$	0	0	$\frac{1}{4}$	$\frac{11}{16}$	34
0	$\frac{7}{16}$	0	1	$\frac{1}{4}$	$-\frac{1}{16}$	10
0	$\frac{5}{4}$	1	0	0	$\frac{1}{4}$	8

No negative entries in the objective row, so this is optimal.

5 (e)

Q = 34, X = 0, Y = 8, Z = 10


P = Q + 16 = 50

x = 0 + 4 = 4, y = 8 + 2 = 10, z = −10 − 2 = −12

Maximum P = 50 when x = 4, y = 10, z = −12

Question number

6 (a)



Vertex	Order	Final label	Working values
A	1	0	0
B	2	15	15
D	3	26	29 26
C	4	31	32 31
E	5	34	37 35 34
F	6	41	43 41
G	7	48	50 48

Shortest path from A to G: A → B → C → F → G, length 48


6(b)(i) A is the start vertex: exactly one of the arcs out of A is used, so the route leaves A once and never comes back to A.

6(b)(ii) The number of arcs used going into F equals the number of arcs used coming out of F: if the route enters F it must also leave F.

Question number	
6(b)(iii)	The edge between C and E can be used at most once, in one direction or the other (CE or EC or neither, but not both).
6(c)(i)	Arcs with value 1: AC, BE, CB, CF, DE, EC, EG, FD Length = 32 + 20 + 16 + 10 + 8 + 12 + 16 + 17 = 131
6(c)(ii)	A → C → B → E → C → F → D → E → G A → C → F → D → E → C → B → E → G
7 (a)	Income $P = 5x + 8y + 12z$ and $x + y + z = 210 \Rightarrow z = 210 - x - y$ $P = 5x + 8y + 12(210 - x - y) = 2520 - 7x - 4y$ 2520 is fixed, so P is greatest when $7x + 4y$ is as small as possible.
7 (b)	Material: $\frac{x}{250} + \frac{y}{175} + \frac{z}{150} \leq 1 \Rightarrow 21x + 30y + 35z \leq 5250$ Time (minutes): $40x + 70y + 140z \leq 220 \times 60 = 13200$ $\Rightarrow 4x + 7y + 14z \leq 1320$ Demand: $x \geq 3z$ Substitute $z = 210 - x - y$: $21x + 30y + 35(210 - x - y) \leq 5250 \Rightarrow 14x + 5y \geq 2100$ $4x + 7y + 14(210 - x - y) \leq 1320 \Rightarrow 10x + 7y \geq 1620$ $x \geq 3(210 - x - y) \Rightarrow 4x + 3y \geq 630$ $z \geq 0 \Rightarrow x + y \leq 210$ $x \geq 0$, $y \geq 0$, x, y integers

Question number

7 (c)



Feasible region (FR) is hatched: below $x + y = 210$ and above the other lines. Vertices: (162, 0), (210, 0), (116.7, 93.3), (137.5, 35)

$7x + 4y$ at vertices: 1134, 1470, 1190, 1102.5

Minimum at intersection of $14x + 5y = 2100$ and $10x + 7y = 1620$:

$98x + 35y = 14700$, $50x + 35y = 8100 \Rightarrow 48x = 6600$

$x = 137.5$, $y = 35$ (not integer), so test nearby integer points:

(x, y)	$14x + 5y \geq 2100$	$10x + 7y \geq 1620$	$7x + 4y$
(137, 37)	2103 ✓	1629 ✓	1107
(138, 35)	2107 ✓	1625 ✓	1106
(139, 33)	2111 ✓	1621 ✓	1105
(140, 31)	2115 ✓	1617 ✗	(not feasible)
(140, 32)	2120 ✓	1624 ✓	1108
(141, 30)	2124 ✓	1620 ✓	1107

(139, 33) also satisfies $4x + 3y = 655 \geq 630$ and $x + y = 172 \leq 210$

Minimum $7x + 4y = 1105$ at $x = 139$, $y = 33$, so $z = 210 - 139 - 33 = 38$

Question number

Kai should make 139 small, 33 medium and 38 large cars.

7 (d) Income = 5(139) + 8(33) + 12(38) = 695 + 264 + 456

= £1415 (check: 2520 − 1105 = 1415)