OCR
Oxford Cambridge and RSA

Thursday 23 June 2022 – Afternoon

A Level Further Mathematics B (MEI)

Y435/01 Extra Pure

Time allowed: 1 hour 15 minutes

9453736284

You must have:
- the Printed Answer Booklet
- the Formulae Booklet for Further Mathematics B (MEI)
- a scientific or graphical calculator

INSTRUCTIONS
- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided in the **Printed Answer Booklet**. If you need extra space use the lined pages at the end of the Printed Answer Booklet. The question numbers must be clearly shown.
- Fill in the boxes on the front of the Printed Answer Booklet.
- Answer **all** the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.
- Give your final answers to a degree of accuracy that is appropriate to the context.
- Do **not** send this Question Paper for marking. Keep it in the centre or recycle it.

INFORMATION
- The total mark for this paper is **60**.
- The marks for each question are shown in brackets **[ ]**.
- This document has **8** pages.

ADVICE
- Read each question carefully before you start your answer.

© OCR 2022 [F/508/5599]
DC (ST/CT) 301215/4
OCR is an exempt Charity

stemup.co.uk

Answer **all** the questions.

1 Three sequences, a_n, b_n and c_n, are defined for $n \geqslant 1$ by the following recurrence relations.

$(a_{n+1} - 2)(2 - a_n) = 3$ with $a_1 = 3$

$b_{n+1} = -\frac{1}{2}b_n + 3$ with $b_1 = 1.5$

$c_{n+1} - \frac{c_n^2}{n} = 1$ with $c_1 = 2.5$

The output from a spreadsheet which presents the first 10 terms of a_n, b_n and c_n, is shown below.

	A	B	C	D
1	n	a_n	b_n	c_n
2	1	3	1.5	2.5
3	2	–1	2.25	7.25
4	3	3	1.875	27.28125
5	4	–1	2.0625	249.0889
6	5	3	1.96875	15512.32
7	6	–1	2.01563	48126390
8	7	3	1.99219	3.86E+14
9	8	–1	2.00391	2.13E+28
10	9	3	1.99805	5.66E+55
11	10	–1	2.00098	3.6E+110

Without attempting to solve any recurrence relations, describe the apparent behaviour, including as $n \to \infty$, of

- a_n
- b_n
- c_n

[7]

© OCR 2022

stemup.co.uk

2 The matrix $\mathbf{A}$ is given by $\mathbf{A} = \begin{pmatrix} 10 & 12 & -8 \\ -1 & 2 & 4 \\ 3 & 6 & 2 \end{pmatrix}$.

(a) In this question you must show detailed reasoning.

Show that the characteristic equation of $\mathbf{A}$ is $-\lambda^3 + 14\lambda^2 - 56\lambda + 64 = 0$. **[3]**

(b) Use the Cayley-Hamilton theorem to determine $\mathbf{A}^{-1}$. **[5]**

A matrix $\mathbf{E}$ and a diagonal matrix $\mathbf{D}$ are such that $\mathbf{A} = \mathbf{EDE}^{-1}$. The elements in the diagonal of $\mathbf{D}$ increase from top left to bottom right.

(c) Determine the matrix $\mathbf{D}$. **[4]**

3 A sequence is defined by the recurrence relation $5t_{n+1} - 4t_n = 3n^2 + 28n + 6$, for $n \geqslant 0$, with $t_0 = 7$.

(a) Find an expression for t_n in terms of n. **[6]**

Another sequence is defined by $v_n = \dfrac{t_n}{n^m}$, for $n \geqslant 1$, where m is a constant.

(b) In each of the following cases determine $\lim_{n\to\infty} v_n$, if it exists, or show that the sequence is divergent.

(i) $m = 3$ **[1]**

(ii) $m = 2$ **[1]**

(iii) $m = 1$ **[1]**

stemup.co.uk

4 A binary operation, $\circ$, is defined on a set of numbers, A, in the following way.

$a \circ b = k_1 a - k_2 b + k_3$, for $a, b \in A$,

where k_1, k_2 and k_3 are constants (which are not necessarily in A) and the operations addition, subtraction and multiplication of numbers have their usual notation and meaning.

You are initially given the following information about $\circ$ and A.

- $A = \mathbb{R}$
- $0 \circ 0 = 2$
- An identity element, e, exists for $\circ$ in A

(a) Show that $a \circ b = a + b + 2$. **[5]**

(b) State the value of e. **[1]**

(c) Explain whether $\circ$ is commutative over A. **[1]**

(d) Determine whether or not $(A, \circ)$ is a group. **[6]**

(e) Explain whether your answer to part **(d)** would change in each of the following cases, giving details of any change.

(i) $A = \mathbb{Z}$ **[1]**

(ii) $A = \{2m : m \in \mathbb{Z}\}$ **[1]**

(iii) $A = \{n : n \in \mathbb{Z}, n \geqslant -2\}$ **[1]**

© OCR 2022

stemup.co.uk

5 A surface S is defined by $z = \mathrm{f}(x, y)$, where $\mathrm{f}(x, y) = y\,\mathrm{e}^{-(x^2+2x+2)y}$.

(a) (i) Find $\dfrac{\partial \mathrm{f}}{\partial x}$. **[1]**

(ii) Show that $\dfrac{\partial \mathrm{f}}{\partial y} = -(x^2y + 2xy + 2y - 1)\,\mathrm{e}^{-(x^2+2x+2)y}$. **[1]**

(iii) Determine the coordinates of any stationary points on S. **[4]**

Fig. 5.1 shows the graph of $z = \mathrm{e}^{-x^2}$ and **Fig. 5.2** shows the contour of S defined by $z = 0.25$.




Fig. 5.1




Fig. 5.2

(b) Specify a sequence of transformations which transforms the graph of $z = \mathrm{e}^{-x^2}$ onto the graph of the section defined by $z = \mathrm{f}(x, 1)$. **[2]**

(c) Hence, or otherwise, sketch the section defined by $z = \mathrm{f}(x, 1)$. **[1]**

(d) Using **Fig. 5.2** and your answer to part **(c)**, classify any stationary points on S, justifying your answer. **[2]**

You are given that P is a point on S where $z = 0$.

(e) Find, in vector form, the equation of the tangent plane to S at P. **[4]**

The tangent plane found in part **(e)** intersects S in a straight line, L.

(f) Write down, in vector form, the equation of L. **[1]**

END OF QUESTION PAPER

© OCR 2022

stemup.co.uk

BLANK PAGE

© OCR 2022

stemup.co.uk

BLANK PAGE

© OCR 2022

stemup.co.uk

OCR
Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.

© OCR 2022

stemup.co.uk

Surname	StemUp Model Solution
Answer booklet	Write the question number in the left-hand margin at the start of each answer.

Question number

1

a_n: the terms alternate 3, −1, 3, −1, ...

so a_n is periodic with period 2 (repeating block 3, −1).

It oscillates and does not converge as $n \to \infty$.

b_n: the terms are alternately below and above 2, getting closer to 2,

so b_n is oscillating and convergent: $b_n \to 2$ as $n \to \infty$ (limit is 2).

c_n: the terms are increasing (and increasing very rapidly),

so c_n is divergent: $c_n \to \infty$ as $n \to \infty$.

2 (a)

$$\det(A - \lambda I) = \det \begin{pmatrix} 10-\lambda & 12 & -8 \\ -1 & 2-\lambda & 4 \\ 3 & 6 & 2-\lambda \end{pmatrix}$$

$$= (10-\lambda)[(2-\lambda)^2 - 4\times 6] - 12[-(2-\lambda) - 4\times 3] - 8[-1\times 6 - 3(2-\lambda)]$$

$$= (10-\lambda)(\lambda^2 - 4\lambda + 4 - 24) - 12(-2 + \lambda - 12) - 8(-6 - 6 + 3\lambda)$$

$$= (10-\lambda)(\lambda^2 - 4\lambda - 20) - 12(\lambda - 14) - 8(3\lambda - 12)$$

$$= 10\lambda^2 - 40\lambda - 200 - \lambda^3 + 4\lambda^2 + 20\lambda - 12\lambda + 168 - 24\lambda + 96$$

$$= -\lambda^3 + 14\lambda^2 - 56\lambda + 64$$

So the characteristic equation is $-\lambda^3 + 14\lambda^2 - 56\lambda + 64 = 0$, as required

2 (b)

By the Cayley-Hamilton theorem: $-A^3 + 14A^2 - 56A + 64I = 0$

Multiply by A^{-1}: $-A^2 + 14A - 56I + 64A^{-1} = 0$

$$A^{-1} = \frac{1}{64}(A^2 - 14A + 56I)$$

$$A^2 = \begin{pmatrix} 10 & 12 & -8 \\ -1 & 2 & 4 \\ 3 & 6 & 2 \end{pmatrix}\begin{pmatrix} 10 & 12 & -8 \\ -1 & 2 & 4 \\ 3 & 6 & 2 \end{pmatrix} = \begin{pmatrix} 64 & 96 & -48 \\ 0 & 16 & 24 \\ 30 & 60 & 4 \end{pmatrix}$$

Question number

$$14A = \begin{pmatrix} 140 & 168 & -112 \\ -14 & 28 & 56 \\ 42 & 84 & 28 \end{pmatrix}$$

$$A^2 - 14A + 56I = \begin{pmatrix} 64 - 140 + 56 & 96 - 168 & -48 + 112 \\ 0 + 14 & 16 - 28 + 56 & 24 - 56 \\ 30 - 42 & 60 - 84 & 4 - 28 + 56 \end{pmatrix}$$

$$A^{-1} = \frac{1}{64}\begin{pmatrix} -20 & -72 & 64 \\ 14 & 44 & -32 \\ -12 & -24 & 32 \end{pmatrix}$$

2 (c) $A = EDE^{-1}$, so D is the diagonal matrix of the eigenvalues of A.

$\lambda = 2$: $-8 + 56 - 112 + 64 = 0 \Rightarrow (\lambda - 2)$ is a factor

$-\lambda^3 + 14\lambda^2 - 56\lambda + 64 = \lambda^2(2 - \lambda) - 12\lambda(2 - \lambda) + 32(2 - \lambda)$

$= (2 - \lambda)(\lambda^2 - 12\lambda + 32) = (2 - \lambda)(\lambda - 4)(\lambda - 8)$

Eigenvalues are 2, 4 and 8

Diagonal increasing: $D = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 8 \end{pmatrix}$

3 (a) Complementary function: $5t_{n+1} - 4t_n = 0 \Rightarrow t_n = \alpha(0.8)^n$

Particular solution: try $t_n = an^2 + bn + c$

$5[a(n + 1)^2 + b(n + 1) + c] - 4[an^2 + bn + c] = 3n^2 + 28n + 6$

$5an^2 + 10an + 5a + 5bn + 5b + 5c - 4an^2 - 4bn - 4c = 3n^2 + 28n + 6$

$an^2 + (10a + b)n + (5a + 5b + c) = 3n^2 + 28n + 6$

n^2: $a = 3$ $\quad$ n: $10a + b = 28 \Rightarrow b = -2$

constant: $5a + 5b + c = 6 \Rightarrow 15 - 10 + c = 6 \Rightarrow c = 1$

General solution: $t_n = \alpha(0.8)^n + 3n^2 - 2n + 1$

$t_0 = 7$: $\alpha + 1 = 7 \Rightarrow \alpha = 6$

Question number	
	$t_n = 6(0.8)^n + 3n^2 - 2n + 1$
3 (b)(i)	$m = 3$: $v_n = \frac{6(0.8)^n}{n^3} + \frac{3}{n} - \frac{2}{n^2} + \frac{1}{n^3}$
	every term $\to 0$ as $n \to \infty$, so $\lim_{n\to\infty} v_n = 0$
3 (b)(ii)	$m = 2$: $v_n = 3 + \frac{6(0.8)^n}{n^2} - \frac{2}{n} + \frac{1}{n^2}$
	all terms except 3 tend to 0, so $\lim_{n\to\infty} v_n = 3$
3 (b)(iii)	$m = 1$: $v_n = 3n - 2 + \frac{6(0.8)^n}{n} + \frac{1}{n}$
	$3n - 2 \to \infty$ and the other terms $\to 0$, so $v_n \to \infty$
	The sequence is divergent (the limit does not exist).
4 (a)	$0 \circ 0 = k_1(0) - k_2(0) + k_3 = 2 \Rightarrow k_3 = 2$
	Identity: $a \circ e = a$ for all $a \in \mathbb{R}$
	$k_1 a - k_2 e + 2 = a$ for all $a \Rightarrow k_1 = 1$ and $-k_2 e + 2 = 0$
	Also $e \circ a = a$ for all $a \in \mathbb{R}$
	$k_1 e - k_2 a + 2 = a$ for all $a \Rightarrow -k_2 = 1 \Rightarrow k_2 = -1$
	(and then $e + 2 = 0$ from both equations, which is consistent)
	$k_1 = 1$, $k_2 = -1$, $k_3 = 2 \Rightarrow a \circ b = a + b + 2$, as required
4 (b)	$e + 2 = 0 \Rightarrow e = -2$
	(check: $a \circ (-2) = a - 2 + 2 = a$ and $(-2) \circ a = a$)
4 (c)	$b \circ a = b + a + 2 = a + b + 2 = a \circ b$ for all $a, b \in \mathbb{R}$,
	so $\circ$ is commutative over A.

Question number	
4 (d)	Closure: $a, b \in \mathbb{R} \Rightarrow a + b + 2 \in \mathbb{R}$, so A is closed under $\circ$
	Associativity:
	$(a \circ b) \circ c = (a + b + 2) + c + 2 = a + b + c + 4$
	$a \circ (b \circ c) = a + (b + c + 2) + 2 = a + b + c + 4$
	equal, so $\circ$ is associative
	Identity: $e = -2 \in \mathbb{R}$ exists (given, found in (b))
	Inverse: need $a \circ a^{-1} = -2 \Rightarrow a + a^{-1} + 2 = -2$
	$a^{-1} = -4 - a$, which is in $\mathbb{R}$ for every $a \in \mathbb{R}$ (and $a^{-1} \circ a = -2$ too)
	All four group axioms are satisfied, so $(A, \circ)$ is a group.
4 (e)(i)	$A = \mathbb{Z}$: $a + b + 2 \in \mathbb{Z}$, $e = -2 \in \mathbb{Z}$, $a^{-1} = -4 - a \in \mathbb{Z}$, associativity
	still holds, so the arguments in (d) still apply: no change, still a group.
4 (e)(ii)	A = even integers: $a + b + 2$ is even, $e = -2$ is even, $-4 - a$ is even,
	so the arguments in (d) still apply: no change, still a group.
4 (e)(iii)	$A = \{n \in \mathbb{Z} : n \geq -2\}$: closure, associativity and the identity -2 still hold,
	but the inverse property fails, eg $2^{-1} = -4 - 2 = -6 \notin A$.
	So the answer changes: $(A, \circ)$ is not a group.
5 (a)(i)	$\frac{\partial f}{\partial x} = y\, e^{-(x^2 + 2x + 2)y} \times (-y(2x + 2)) = -2y^2(x + 1)e^{-(x^2 + 2x + 2)y}$
5 (a)(ii)	Product rule (with the chain rule):
	$\frac{\partial f}{\partial y} = 1 \times e^{-(x^2 + 2x + 2)y} + y \times (-(x^2 + 2x + 2))e^{-(x^2 + 2x + 2)y}$
	$= (1 - y(x^2 + 2x + 2))e^{-(x^2 + 2x + 2)y}$

Question number

$= -(x^2y + 2xy + 2y - 1)e^{-(x^2 + 2x + 2)y}$, as required

5 (a)(iii) $\frac{\partial f}{\partial x} = 0 \Rightarrow -2y^2(x + 1) = 0 \Rightarrow y = 0$ or $x = -1$

$y = 0$: $\frac{\partial f}{\partial y} = -(-1)e^0 = 1 \neq 0$, so no stationary point

$x = -1$: $\frac{\partial f}{\partial y} = -(y - 2y + 2y - 1)e^{-y} = -(y - 1)e^{-y} = 0 \Rightarrow y = 1$

$z = f(-1, 1) = 1 \times e^{-(1 - 2 + 2)} = e^{-1}$


Only stationary point: $(-1, 1, e^{-1})$

5 (b) $f(x, 1) = e^{-(x^2 + 2x + 2)} = e^{-1}e^{-(x + 1)^2}$

Translation by 1 unit in the negative x-direction,

followed by a stretch parallel to the z-axis, scale factor e^{-1}.

5 (c) $z = e^{-1}e^{-(x + 1)^2}$



5 (d) The stationary point $(-1, 1, e^{-1})$ is a maximum point.

The section $y = 1$ in (c) has a maximum at $x = -1$, with $z = e^{-1} \approx 0.368$.

Since $e^{-1} > 0.25$ and the contour $z = 0.25$ is a closed curve around

Question number

$(-1, 1)$, moving from the stationary point in any direction we must go down to $z = 0.25$ (it is the only stationary point), so it is a maximum.

5 (e) $z = 0 \Rightarrow y\,e^{-(x^2 + 2x + 2)y} = 0 \Rightarrow y = 0$ (exponential > 0)

So $P = (a, 0, 0)$ for some a.

At $y = 0$: $\partial f/\partial x = -2(0)^2(a + 1) = 0$, $\partial f/\partial y = -(-1)e^0 = 1$

$$\text{Normal } n = \begin{pmatrix} \partial f/\partial x \\ \partial f/\partial y \\ -1 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}$$

$$\begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} \cdot \begin{pmatrix} a \\ 0 \\ 0 \end{pmatrix} = 0 \text{ for any } a, \text{ so the plane passes through } O$$

$$\text{Tangent plane: } r \cdot \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} = 0 \quad (\text{ie } z = y)$$

5 (f) On the plane $z = y$, so $y = y\,e^{-(x^2 + 2x + 2)y} \Rightarrow y = 0$ $(x^2 + 2x + 2 > 0)$

$$L: \quad r = \lambda \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$