OCR
Oxford Cambridge and RSA

Monday 24 June 2024 – Afternoon

A Level Further Mathematics B (MEI)

Y436/01 Further Pure with Technology

Time allowed: 1 hour 45 minutes

141904481 5

You must have:
- the Printed Answer Booklet
- the Formulae Booklet for Further Mathematics B (MEI)
- a computer with appropriate software
- a scientific or graphical calculator

QP

INSTRUCTIONS
- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided in the **Printed Answer Booklet**. If you need extra space use the lined pages at the end of the Printed Answer Booklet. The question numbers must be clearly shown.
- Fill in the boxes on the front of the Printed Answer Booklet.
- Answer **all** the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.
- Give your final answers to a degree of accuracy that is appropriate to the context.
- Do **not** send this Question Paper for marking. Keep it in the centre or recycle it.

INFORMATION
- The total mark for this paper is **60**.
- The marks for each question are shown in brackets **[ ]**.
- This document has **8** pages.

ADVICE
- Read each question carefully before you start your answer.

© OCR 2024 [K/508/5600]

OCR is an exempt Charity

stemup.co.uk

1 A family of curves is given by the equation

$$y = \frac{x^2 - x + a^2 - a}{x - 1} \quad (*)$$

where the parameter a is a real number.

(a) (i) On the axes in the Printed Answer Booklet, sketch the curve in each of these cases.

- $a = -0.5$
- $a = -0.1$
- $a = 0.5$

[3]

(ii) State **one** feature of the curve for the cases $a = -0.5$ and $a = -0.1$ that is **not** a feature of the curve in the case $a = 0.5$. **[1]**

(iii) By using a slider for a, or otherwise, write down the **non-zero** value of a for which the points on the curve (*) all lie on a straight line. **[1]**

(iv) Write down the equation of the vertical asymptote of the curve (*). **[1]**

The equation of the curve (*) can be written in the form $y = x + A + \frac{a^2 - a}{x - 1}$, where A is a constant.

(v) Show that $A = 0$. **[2]**

(vi) Hence, or otherwise, find the value of

$$\lim_{x \to \infty} \left(\frac{x^2 - x + a^2 - a}{x - 1} - x \right).$$ **[2]**

(vii) Explain the significance of the result in part **(a)(vi)** in terms of a feature of the curve (*). **[1]**

(b) In this part of the question the value of the parameter a satisfies $0 < a < 1$. For values of a in this range the curve intersects the x-axis at points X and Y. The point Z has coordinates $(0, -1)$. These three points form a triangle XYZ.

(i) Determine, in terms of a, the area of the triangle XYZ. **[4]**

(ii) Find the maximum area of the triangle XYZ. **[2]**

© OCR 2024

stemup.co.uk

2 Wilson's theorem states that a positive integer $n > 1$ is prime if and only if $(n-1)! \equiv n-1 \pmod{n}$.

(a) (i) Calculate $136! \pmod{137}$. **[1]**

(ii) Hence, determine if the integer 137 is prime. **[1]**

(b) (i) Create a program that uses Wilson's theorem to find all prime numbers less than or equal to n.
Write down your program in the Printed Answer Booklet. **[3]**

(ii) Using part **(b)(i)**, write down **all** prime numbers x, where $260 \leqslant x \leqslant 300$. **[1]**

(c) (i) Explain why there is exactly **one** prime number congruent to $2 \pmod{4}$. **[1]**

(ii) Explain why **no** prime number is congruent to $0 \pmod{4}$. **[1]**

(iii) Using part **(b)(ii)**, write down the **three** prime numbers y, where $260 \leqslant y \leqslant 300$, that are congruent to $3 \pmod{4}$. **[1]**

Label the three prime numbers in part **(c)(iii)** c_1, c_2 and c_3. Define the integer N by

$N = 4c_1c_2c_3 + 3$.

(iv) Explain why N is **not** divisible by c_1, c_2 or c_3. **[1]**

(v) Write down the value of $N \pmod{4}$. **[1]**

(d) The fundamental theorem of arithmetic states that every positive integer can be written uniquely as the product of powers of prime numbers.

Suppose there are finitely many prime numbers congruent to $3 \pmod{4}$. Label these prime numbers $p_1, \ldots, p_n$, where $p_1 = 3$. Using the $n-1$ integers $p_2, \ldots, p_n$, define the integer M by

$M = 4p_2p_3 \ldots p_n + 3$.

(i) Write down the value of $M \pmod{4}$. **[1]**

(ii) Explain why $M = q_1q_2 \ldots q_r$, where the integers $q_1, \ldots, q_r$ are all prime. **[2]**

(iii) Prove that there is at least one integer i, where $1 \leqslant i \leqslant n$, such that $q_i \equiv 3 \pmod{4}$. **[4]**

(iv) Hence, deduce that there are infinitely many prime numbers congruent to $3 \pmod{4}$. **[5]**

stemup.co.uk

3 This question concerns the family of differential equations

$$\frac{dy}{dx} = x^2 - y + a\cos(x)\cos(y) \quad (**)$$

where a is a constant, $x \geqslant 0$ and $y > 0$.

(a) In this part of the question $a = 0$.

(i) Find the solution to (**) in which $y = 1$ when $x = 0$. **[1]**

(ii) In this part of the question m is a real number. Show that the equation of the isocline $\frac{dy}{dx} = m$ is a parabola. **[1]**

(iii) Using the result given in part **(a)(ii)**, or otherwise, sketch the tangent field for (**) on the axes in the Printed Answer Booklet. **[2]**

(b) **Fig. 3.1** and **Fig. 3.2** show the tangent fields for two distinct and unspecified values of a. In each case, a sketch of the solution curve $y = g(x)$ which passes through the point (0, 2) is shown for $0 \leqslant x \leqslant \frac{1}{2}$.

Fig. 3.1





Fig. 3.2



(i) In each case, continue the sketch of the solution curve for $\frac{1}{2} \leqslant x \leqslant 3$ on the axes in the Printed Answer Booklet. **[2]**

(ii) State **one** feature which is present in the continued solution curve for **Fig. 3.1** that is **not** a feature of the continued solution curve for **Fig. 3.2**. **[1]**

(iii) Using a slider for a, or otherwise, estimate the value of a for the solution curve shown in **Fig. 3.2**. **[1]**

© OCR 2024

stemup.co.uk

(c) The Euler method for the solution of the differential equation $\frac{dy}{dx} = f(x, y)$ is as follows.

$$y_{n+1} = y_n + hf(x_n, y_n)$$
$$x_{n+1} = x_n + h$$

(i) Construct a spreadsheet to solve (**), so that the value of a and the value of h can be varied, in the case $x_0 = 0$ and $y_0 = 1$. State the formulae you have used in your spreadsheet. **[2]**

(ii) In this part of the question $a = 0$. Use your spreadsheet with $h = 0.1$ to approximate the value of y when $x = 0.5$ for the solution to (**) in which $y = 1$ when $x = 0$. **[1]**

(iii) Using part **(a)(i)**, state the accuracy of the approximate value of y given in part **(c)(ii)**. **[1]**

(iv) State **one** change to your spreadsheet that could improve the accuracy of the approximate value of y found in part **(c)(ii)**. **[1]**

(d) The modified Euler method for the solution of the differential equation $\frac{dy}{dx} = f(x, y)$ is as follows.

$$k_1 = hf(x_n, y_n)$$
$$k_2 = hf(x_n + h, y_n + k_1)$$
$$y_{n+1} = y_n + \frac{1}{2}(k_1 + k_2)$$
$$x_{n+1} = x_n + h$$

(i) Adapt your spreadsheet from part **(c)(i)** to a spreadsheet to solve (**), so that the value of a and the value of h can be varied, in the case $x_0 = 0$ and $y_0 = 1$. State the formulae you have used in your spreadsheet. **[2]**

(ii) In this part of the question $a = -0.5$. Use the spreadsheet from part **(d)(i)** with $h = 0.1$ to approximate the value of y when $x = 0.5$ for the solution to (**) in which $y = 1$ when $x = 0$. **[1]**

In this part of the question $a = -0.5$. The solution to (**) in which $y = 1$ when $x = 0$ has a turning point with coordinates (c, d) where $0 < c < 1$.

(iii) Use the spreadsheet in part **(d)(i)** to determine the value of c correct to **1** decimal place. **[2]**

(iv) Use the spreadsheet in part **(d)(i)** to determine the value of d correct to **3** decimal places. **[2]**

END OF QUESTION PAPER

© OCR 2024

stemup.co.uk

BLANK PAGE

© OCR 2024

stemup.co.uk

BLANK PAGE

© OCR 2024

stemup.co.uk

OCR
Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.


© OCR 2024

stemup.co.uk

Surname	StemUp Model Solution
Answer booklet	Write the question number in the left-hand margin at the start of each answer.

Question number

1 (a)(i) $a = -0.5$ $a = -0.1$ $a = 0.5$



y-intercepts: $(0, -0.75)$, $(0, -0.11)$, $(0, 0.25)$ respectively; $x = 1$ asymptote

1 (a)(ii) For $a = -0.5$ and $a = -0.1$ the curve has turning points (a local maximum and a local minimum); for $a = 0.5$ it has no turning points.

1 (a)(iii) $a = 1$ (then $a^2 - a = 0$ and $y = x$, for $x \neq 1$)

1 (a)(iv) $x = 1$

1 (a)(v) $y = x + A + \dfrac{a^2 - a}{x - 1} = \dfrac{(x + A)(x - 1) + a^2 - a}{x - 1}$

$= \dfrac{x^2 + (A - 1)x - A + a^2 - a}{x - 1}$

Comparing with $x^2 - x + a^2 - a$, the coefficient of x gives

$A - 1 = -1 \Rightarrow A = 0$, as required.

Question number

1 (a)(vi) By (a)(v), $\frac{x^2 - x + a^2 - a}{x - 1} = x + \frac{a^2 - a}{x - 1}$

$\lim_{x\to\infty} \left(\frac{x^2 - x + a^2 - a}{x - 1} - x\right) = \lim_{x\to\infty} \frac{a^2 - a}{x - 1}$

$= 0$

1 (a)(vii) The line $y = x$ is an oblique asymptote of the curve:

the curve tends towards $y = x$ as $x \to \infty$.

1 (b)(i) Curve meets the x-axis when $\frac{x^2 - x + a^2 - a}{x - 1} = 0$

$\Rightarrow x^2 - x + a^2 - a = 0$

$x = \frac{1 \pm \sqrt{1 - 4(a^2 - a)}}{2} = \frac{1 \pm \sqrt{1 + 4a - 4a^2}}{2}$

X: $\left(\frac{1}{2} - \frac{\sqrt{1 + 4a - 4a^2}}{2}, 0\right)$ Y: $\left(\frac{1}{2} + \frac{\sqrt{1 + 4a - 4a^2}}{2}, 0\right)$

(for $0 < a < 1$, $1 + 4a - 4a^2 > 0$, so X and Y are distinct)

Base XY = $\sqrt{(1 + 4a - 4a^2)}$, height = distance of Z from x-axis = 1

Area $= \frac{1}{2} \times \sqrt{1 + 4a - 4a^2} \times 1 = \frac{1}{2}\sqrt{1 + 4a - 4a^2}$

1 (b)(ii) $\frac{d}{da}\left(\frac{1}{2}\sqrt{1 + 4a - 4a^2}\right) = \frac{1 - 2a}{\sqrt{1 + 4a - 4a^2}} = 0 \quad \Rightarrow \quad a = \frac{1}{2}$

Maximum area $= \frac{1}{2}\sqrt{1 + 2 - 1} = \frac{\sqrt{2}}{2}$ (≈ 0.707)

2 (a)(i) $136! \pmod{137} = 136$

Question number	
2 (a)(ii)	$136! \equiv 136 \equiv -1 \pmod{137}$, so by Wilson's theorem 137 is prime.

2 (b)(i)	(Python)
	`import math`
	`def wlistprimes(n):`
	`  for j in range(2, n + 1):`
	`  if (math.factorial(j - 1) + 1) % j == 0:`
	`  print(j)`

2 (b)(ii)	263, 269, 271, 277, 281, 283, 293
2 (c)(i)	A number $\equiv 2 \pmod 4$ is even. 2 is the only even prime, so 2 is the only prime congruent to 2 (mod 4).
2 (c)(ii)	A number $\equiv 0 \pmod 4$ is divisible by 4 (and is at least 4), so it is not prime.
2 (c)(iii)	263, 271, 283
2 (c)(iv)	$N = 4c_1c_2c_3 + 3$, and $4c_1c_2c_3$ is a multiple of each of c_1, c_2, c_3, so N leaves remainder 3 when divided by c_1, c_2 or c_3 (as each $c_i > 3$), so N is not divisible by c_1, c_2 or c_3.
2 (c)(v)	$N \equiv 3 \pmod 4$
2 (d)(i)	$M \equiv 3 \pmod 4$

Question number	
2 (d)(ii)	M is a positive integer ($M > 1$), so by the fundamental theorem of arithmetic M can be written uniquely as a product of powers of primes. Writing each prime power as a repeated product of that prime gives $M = q_1 q_2 \dots q_r$, where the primes q_i are not necessarily distinct.
2 (d)(iii)	M is odd, so no $q_i = 2$; by (c)(ii) no prime is $\equiv 0 \pmod 4$, so each $q_i \equiv 1$ or $3 \pmod 4$. For a contradiction, suppose $q_i \equiv 1 \pmod 4$ for all i, $1 \leq i \leq r$. Then $M \equiv q_1 q_2 \dots q_r \pmod 4$ $\equiv 1 \times 1 \times \dots \times 1 \pmod 4$ $\equiv 1 \pmod 4$ But $M \equiv 3 \pmod 4$ by (d)(i). This is a contradiction, so $q_i \not\equiv 1 \pmod 4$ for some i, i.e. there is at least one i with $q_i \equiv 3 \pmod 4$.
2 (d)(iv)	By (d)(iii), some $q_i \equiv 3 \pmod 4$, and by (d)(ii) q_i divides M, i.e. $M \equiv 0 \pmod{q_i}$. By assumption $p_1, \dots, p_n$ is the complete list of primes $\equiv 3 \pmod 4$, so q_i is one of $p_1, p_2, \dots, p_n$. $q_i \neq 3 = p_1$, since $M = 4p_2 \dots p_n + 3$ with $3 \nmid 4p_2 \dots p_n$, so $3 \nmid M$. Hence q_i is one of $p_2, \dots, p_n$. Then q_i divides $4p_2 \dots p_n$, so $M \equiv 4p_2 \dots p_n + 3 \equiv 0 + 3 \equiv 3 \pmod{q_i}$ This contradicts $M \equiv 0 \pmod{q_i}$ (as $q_i > 3$). So the assumption is false: there are infinitely many prime numbers congruent to $3 \pmod 4$.

Question number

3 (a)(i) $a = 0$: $\frac{dy}{dx} + y = x^2 \Rightarrow y = x^2 - 2x + 2 + Ce^{-x}$


$y = 1$ when $x = 0$: $1 = 2 + C \Rightarrow C = -1$

$y = x^2 - 2x + 2 - e^{-x}$

3 (a)(ii) $\frac{dy}{dx} = m \Leftrightarrow x^2 - y = m \Leftrightarrow y = x^2 - m$

which is the equation of a parabola.


3 (a)(iii)



(gradient is constant along each parabola $y = x^2 - m$)

3 (b)(i) Fig. 3.1 Fig. 3.2

Question number



3 (b)(ii) The curve for Fig. 3.1 has a local minimum close to $x = 1$

(Fig. 3.2 has a local maximum near $x = 1$ instead).

3 (b)(iii) $a \approx -5.5$

3 (c)(i) I1: 0.1 (value of h) K1: 0 (value of a)

A2: 0 (x_0) B2: 1 (y_0)

C2: =A2^2−B2+$K$1∗COS(A2)∗COS(B2) (dy/dx)

D2: =$I$1∗C2 (h f(x_n, y_n))

A3: =A2+$I$1

B3: =B2+D2

then fill C2:D2 and A3:B3 down the columns.

3 (c)(ii) $h = 0.1$, $a = 0$: $y(0.5) \approx 0.618559$ (0.6186 to 4 d.p.)

3 (c)(iii) Exact: $y(0.5) = 0.25 - 1 + 2 - e^{-0.5} = 0.643469...$

Error = 0.643469 − 0.618559 = 0.0249 (about 4%)

So the approximation is accurate to 1 decimal place (0.6).

Question number	
3 (c)(iv)	Use a smaller value of h (e.g. h = 0.01).
3 (d)(i)	I1: 0.1 (h) K1: −0.5 (a) A2: 0 B2: 1
	C2: =A2^2−B2+$K$1∗COS(A2)∗COS(B2)
	D2: =$I$1∗C2 (k_1)
	E2: =A2+$I$1 ($x_n + h$)
	F2: =B2+D2 ($y_n + k_1$)
	G2: =$I$1∗(E2^2−F2+$K$1∗COS(E2)∗COS(F2)) (k_2)
	A3: =A2+$I$1
	B3: =B2+(1/2)∗(D2+G2)
	then fill C2:G2 and A3:B3 down the columns.
3 (d)(ii)	h = 0.1, a = −0.5: y(0.5) ≈ 0.504227 (0.5042 to 4 d.p.)
3 (d)(iii)	With h = 0.1:

x	y	dy/dx
0.8	0.396189	−0.0776
0.9	0.399006	0.1246

Sign change in dy/dx, so 0.8 < c < 0.9.

Reducing to h = 0.01:

0.83	0.392233	−0.0151
0.84	0.392183	0.0050

Sign change, so 0.83 < c < 0.84

∴ c = 0.8 (1 d.p.)

3 (d)(iv) From the spreadsheet with h = 0.01, near the minimum:

y(0.83) = 0.392233, y(0.84) = 0.392183, y(0.85) = 0.392335

so the least value of y is about 0.39218 (just below 0.392183).

Question number

Reducing h to 0.001 gives a minimum $y \approx 0.392151$ (at $x \approx 0.838$).

Both estimates agree to 3 d.p.

$\therefore\ d = 0.392$ (3 d.p.)